

1

2

3 **Spatial and temporal variation in detection probability**

4 **and abundance of freshwater turtles in a peri-urban area**

5 **of Midwestern Brazil**

6

7 ELIZÂNGELA S. BRITO, NICOLLE B. F. PRADO, KLYSMAN F. F. ALMEIDA, LUCIANA M.

8 VALÉRIO,CHRISTINE STRÜSSMANN, TAINÁ F. DORADO-RODRIGUES

9

10

11

12 **This article has been accepted for publication and undergone full peer review but has not**

13 **been through the copyediting, typesetting, pagination and proofreading process, which**

14 **may lead to differences between this version and the Version of Record.**

15 **Please cite this article as:**

16

17 Brito, E. S., Prado, N. B. F., Almeida, K. F. F., Valério, L. M., Strüssmann, C., Dorado-Rodrigues, T. F. (2025).

18 Spatial and temporal variation in detection probability and abundance of freshwater turtles in a peri-urban

19 area of Midwestern Brazil. *Acta Herpetol.* **21**. doi: 10.36253/a_h-17006.

20

21

22

**Spatial and Temporal Variation in Detection Probability and Abundance of Freshwater
Turtles in a Peri-Urban Area of Midwestern Brazil**

ELIZÂNGELA S. BRITO^{1,2,*}, NICOLLE B. F. PRADO³, KLYSMAN F. F. ALMEIDA⁴, LUCIANA M.
VALÉRIO^{4,5}, CHRISTINE STRÜSSMANN^{2,6}, TAINÁ F. DORADO-RODRIGUES^{1,2}

¹*Programa de Capacitação Institucional, Instituto Nacional de Pesquisas do Pantanal, Av.
Fernando Correa da Costa 2367, 78060-900 Cuiabá, Mato Grosso, Brazil*

²*Laboratório de Herpetologia, Centro de Biodiversidade, Universidade Federal de Mato
Grosso, Av. Fernando Correa da Costa 2367, 78060-900 Cuiabá, Mato Grosso, Brazil*

³*Programa de Pós-graduação em Biologia Animal, Instituto de Biociências, Universidade
Federal do Mato Grosso do Sul, Av. Costa e Silva 8000, 79070-900 Campo Grande, Mato
Grosso do Sul, Brazil*

⁴*Departamento de Ciências Biológicas, Universidade Católica Dom Bosco, Av. Tamandaré
6000, 79117-900 Campo Grande, Mato Grosso do Sul, Brazil*

⁵*Instituto de Biociências, Letras e Ciências Exatas, Universidade Estadual Paulista, Rua
Cristóvão Colombo 2265, 15054-000 São José do Rio Preto, São Paulo, Brazil*

⁶*Faculdade de Medicina Veterinária, Universidade Federal de Mato Grosso, Av. Fernando
Correa da Costa 2367, 78060-900 Cuiabá, Mato Grosso, Brazil*

**Corresponding author. Email: eliz.chelidae@gmail.com*

Submitted on: 2024, 11th November; revised on: 2025, 11th May; accepted on: 2025, 23th July.

Editor: Jairo Mendoza Roldan

Short running title: Spatial and Temporal Variation in a Freshwater Turtle Assemblage

46

47 **Abstract.** This study assessed spatial (across water bodies) and temporal (over months and
48 years) variation in the detection probability and abundance of freshwater turtles, along with the
49 effects of climatic variables such as precipitation and temperature. Fieldwork was conducted in
50 four water bodies located at the headwaters of a stream in the peri-urban area of Campo Grande,
51 Mato Grosso do Sul, in Midwestern Brazil. Over 38 months of sampling (2014–2017), we
52 captured 94 individuals: 73 *Phrynops geoffroanus*, 19 *Mesoclemmys vanderhaegei*, and two
53 *Trachemys dorbigni*. The detection probability of *M. vanderhaegei* was influenced by both
54 precipitation and temperature, whereas detection of *P. geoffroanus* was affected only by
55 temperature. The abundance of both species varied across water body types and sampling
56 months, with *P. geoffroanus* being more abundant in water bodies where *M. vanderhaegei* was
57 less common. *P. geoffroanus* reached its highest abundance during the rainy season (November
58 to March), with fluctuations in the subsequent months. In contrast, *M. vanderhaegei* maintained
59 low abundance throughout most of the year, with a peak in December. To our knowledge, this
60 is the first study to examine population patterns of these two syntopic species. *Phrynops*
61 *geoffroanus*, the more abundant species, is often recorded in anthropogenically altered
62 environments and appears to possess greater competitive ability than *M. vanderhaegei*, which
63 is less abundant and likely more sensitive to environmental degradation. Despite the urban
64 context and the presence of an invasive species (*T. dorbigni*), the occurrence of *M.*
65 *vanderhaegei* suggests that the study area may still retain important elements of habitat quality.

66

67 **Keywords.** Chelidae, chelonians, Testudines, urban ecology, *Phrynops geoffroanus*,
68 *Mesoclemmys vanderhaegei*, *Trachemys dorbigni*.

69

INTRODUCTION

Knowledge about chelonian ecology is still incipient in South America. Population studies are now more frequent, but studies on the structure of assemblages or on the temporal partition of resources of the different species of freshwater turtles are rare, even in the Amazon basin, where the diversity of these reptiles is high (Mittermeier et al., 2015). This highlights the challenge of conducting studies on freshwater turtles in natural environments.

Due to their ease of access, water bodies in urban areas can be considered excellent sites for obtaining information on different aspects of the ecology of freshwater turtles (Dorado-Rodrigues et al., 2024; Rocha et al., 2025). On the other hand, urbanization imposes significant pressure on wildlife, altering natural environments through deforestation, which leads to vegetation fragmentation, species loss, and biodiversity decline (Guzy et al., 2013; Liu et al., 2016; Rocha et al., 2025). Nevertheless, many turtles tolerate, adapt to, and survive in urbanized environments, even those of poor quality, as long as they have access to resources essential for their survival (e.g., food, shelter, and reproductive sites) (Gibbons et al., 2000; Luiselli, 2003; Stokeld et al., 2014; Spencer et al., 2018; Rocha et al., 2025).

Freshwater turtles inhabit different types of habitats, terrestrial and aquatic, which vary according to their basic ecological needs as a function of their semi-aquatic habit (Bodie and Semlitsch, 2000). Terrestrial and aquatic habitats (whether lentic or lotic) undergo changes due to variations promoted by the hydrological cycle which, in turn, affect the availability of resources, and consequently the quality of the habitat (Alho and Pádua, 1982; Fachín-Terán et al., 2006). As a result of these environmental variations, freshwater turtles adjust their preexisting strategies—such as feeding, thermoregulation, and reproduction—to habitat alterations to ensure their survival (Moll and Moll, 2004). Thus, habitat use by freshwater turtles may vary according to periods of drought and rainfall, since these aspects directly influence the

spatial distribution and structure of assemblages by affecting the distribution and supply of resources in the environment (Plummer et al., 1997; Luiselli, 2008).

In this study, we evaluated the effects of population parameters, such as detectability and abundance, for two species of freshwater turtles in relation to temporal (sampling period, rainfall and temperature) and spatial (water bodies) characteristics. We tested the following hypotheses: (1) the detection probability of freshwater turtles varies with monthly precipitation and temperature; (2) The abundance of freshwater turtles varies spatially among water bodies and temporally with sampling months and temperature, as well as with the interaction between water bodies and sampling months; and (3) the abundance of a species varies with the area of the water body and/or the abundance of syntopic species.

MATERIALS AND METHODS

Study area

The study area is located in the urban perimeter of the municipality of Campo Grande (Mato Grosso do Sul, Midwestern Brazil), situated in the Cerrado biogeographic domain and the Upper Paraguay river basin (20° 23'00" S and 54° 36' 00" W; Datum WGS84; Fig. 1). The study was conducted in a vegetation fragment of 191 hectares of which 20 hectares are designated for the protection area called legal reserve in Brazil (Cheung et al., 2016; Corral et al., 2021). The fragment is located in an area of urban expansion and is part of a region that includes both forest and open formations. The forest formations range from *cerradão*—a dense forest formation characterized by tall trees (8–15 m), a closed canopy, a sparse grassy understory, and occurrence in fertile, humid soils—to riparian vegetation, including gallery and riparian forests (Sano et al., 2008; Cheung et al., 2016). The open formations include *Cerrado stricto sensu* and

117 *campo sujo*. *Campo sujo* refers to herbaceous-shrubby habitats dominated by grasses, with few
118 trees and shrubs (<2% of the land cover; Furley, 1999).

119 The climate in the region is of type Aw (tropical climate with a dry winter season),
120 according to the Köppen classification (Peel et al., 2007). The rainy (October to March) and
121 dry (April to September) seasons are well defined (see Marcuzzo and Mello, 2011). The average
122 annual temperature is 24°C and the annual rainfall is between 1800 and 2000 mm.

123 *Data collection*. We sampled four water bodies located at the headwaters of the Segredo stream,
124 300 meters apart (Table 1), in a landscape with fragmented Cerrado vegetation (Fig. 1). We
125 collected information on the freshwater turtle assemblage between March 2014 and July 2017,
126 using funnel traps 1.2 m in length (see description in Brito et al., 2009; 2018; Balestra et al.,
127 2016). The traps were baited with ground beef and remained in the water diurnally, being
128 reviewed once a day over five consecutive days per month. The number of traps ranged from
129 two to four traps per water body (Table 1). The minimum distance between traps within the
130 same water body was 100 meters, while the maximum distance varied up to 150 meters in water
131 body A, the largest of the four. The traps were installed along the margins of each water body.

132 Considering the possibility of imperfect detection, we performed five repeated
133 samplings per month (five consecutive days), necessary to estimate the probability of detection.
134 Since each species could be detected (1) or not detected (0), we constructed a detection history
135 denoting the sequence of detections and non-detections for each species (e.g., 10100 - five
136 sampling occasions within a session, where the species was detected on two and not detected
137 on three). The assumption was that the site occupancy status for each species did not change
138 during the period, or that the changes occurred completely at random (i.e., the species present
139 at the sites were constant during sampling) (MacKenzie et al., 2006).

We identified, marked, sexed, and performed biometrics on all of the captured turtles. We marked each captured turtle individually using a system of rectangular cuts in marginal scutes of the carapace, adapted from Cagle (1939). We determined the sex of the captured individuals by examining secondary sexual characteristics (males having a more elongated tail than females) and measured carapace length (CL; to the nearest 0.05 mm) with a 150 mm or 300 mm Vernier caliper. The sexed individuals were classified as either female or male, but their sexual maturity status was not determined, as this information is unknown for *M. vanderhaegei* and *P. geoffroanus*. We could not determine the sex of some individuals smaller than 100 mm CL for both species investigated. Therefore, we classified them as juveniles. We obtained the body mass with Pesola® spring balances of the following capacities: 2.5 kg (50g precision) and 5.000 g (100g precision).

Finally, we obtained information on monthly precipitation and temperature from the INMET (National Institute of Meteorology) database obtained from the meteorological station in Campo Grande, Mato Grosso do Sul (20° 27'00" S; 54° 43' 12" W), located 25 km from the study area.

Characterization of the water bodies

Among the four water bodies sampled at the headwaters of the Segredo stream, three are artificial dams (A, B and C) and one is a natural pond (D) that remained with water during the entire study period. The area of the water bodies, estimated using Google® Earth Pro software, ranged from 6,542 m² to 9,171 m², are located in the *Cerrado*. The surrounding matrix is formed by floodable riparian forest and *campo sujo*, with tree and shrub vegetation, also containing exotic plant species (e.g., bamboo, eucalyptus, and mango trees). The margin and the interior of the water bodies present vegetation cover by herbaceous and aquatic plants and the substrate

is sandy or clay (Table 1). Despite being located in an area of urban expansion, none of the water bodies receive sewage effluents.

Data analysis

Sex ratio. We used a chi-square test to determine whether the sex ratio differed from 1:1 in each of the turtle populations, employing only those individuals sexed and captured for the first time.

Detectability, temporal and spatial variations. To assess the probability of detection of each species of freshwater turtles, we used the "single-season, single-species" occupancy model, which explicitly accounts for imperfect detection [$\psi(\cdot)\rho(\cdot)$]. Since our goal is focused on comparing only how much the detection pattern changes according to the temporal characteristics of rainfall and daily temperature occurrence, the detection probability was estimated without the effect of environmental variables on occupancy (ψ Naïve). Thus, we tested the effect of rainfall occurrence and daily temperature on the detection probability (from the history of detection [1] and non-detection [0]), considering the other parameters constant [$\psi(\cdot)\rho(\text{variable})$] (MacKenzie et al., 2006). We considered the most parsimonious detection probability model, with the smoothing method with 100 Bootstrap interactions (to compute the standard error of the estimated derived parameters) (MacKenzie et al., 2006). We obtained the parameter estimates by means of maximum likelihood estimation (MLEs).

We used AIC (Akaike Information Criterion; Burnham and Anderson, 2002) to rank and compare the candidate models for each species. We selected the best models (i.e., the model with the parameter that best describes the data as a function of the variable) by considering those with ΔAIC values ≤ 2 . We examined the relevance of the variables in the best selected models for uncertainty from analysis of the 95% confidence interval of the parameter estimates (i.e., the overlap of the confidence interval to zero).

We assessed the normality of species abundance data in relation to water bodies and climatic variables using the Shapiro-Wilk test. To evaluate variations in freshwater turtle abundance (by species and total) in relation to temporal (month sampled, mean daily temperature) and spatial (water bodies) characteristics, we fitted Generalized Linear Mixed Models (GLMMs). The months sampled, mean daily temperature and water bodies, as well as the interaction between them, were considered to be fixed factors. To assess whether species abundance varied across sampling months and between water bodies, random factors, we fitted Generalized Linear Mixed Models (GLMMs) separately for each species and treated fixed and random factors in the same way. Due to the overdispersion in abundance data, we used the quasi-Poisson distribution, which has a better fit to data with inflated variance (Hoef and Boveng, 2007).

To assess whether the abundance of a freshwater turtle is affected by the abundance of another species and whether it varies with the area of the water bodies, we used Generalized Linear Models (GLMs). We evaluated the variation for each sampling year (2014, 2015, 2016, 2017) and considering the total catches taken over the four years. Generalized Linear Models present a major advantage over classical multiple regression approaches in that they integrate data from different statistical distributions with the appropriate modeling of statistical error. We verified by means of the Shapiro-Wilk test the absence of overdispersion in the abundance data, and thus we used the Poisson distribution to estimate abundance.

All statistical analyses were performed in the R 3.6.0 programming environment (R Development Core Team, 2019), using the Occu function in the unmarked package (Fiske and Chandler, 2011), in the AIC analysis, and the MASS package (Venables and Ripley, 2002), for the GLMMs analyses (Venables and Ripley, 2002). We adopted a significance level of $P < 0.05$.

RESULTS

We recorded three species of freshwater turtles in the four water bodies sampled: two native (*Phrynops geoffroanus* and *Mesoclemmys vanderhaegei*) and one introduced species (*Trachemys dorbigni*). In all, we captured 94 individuals: 73 were *P. geoffroanus* (78% of the total; 31 females, 30 males, and five juveniles), 19 *M. vanderhaegei* (20% of the total; 10 females, nine males, and one juvenile) and two *T. dorbigni* (2% of the total). Both *T. dorbigni* individuals were adults (156 and 190 mm CL), but their sexes were not determined. Sex ratio did not differ from 1:1 for either *P. geoffroanus* (0.96M:1F; $\chi^2 = 0.016$, $df = 1$, $P = 0.898$) or *M. vanderhaegei* (1M:1F; $\chi^2 = 0$, $df = 1$, $P = 1.000$). Juveniles represented a small proportion of captures for both *P. geoffroanus* and *M. vanderhaegei* (Fig. 2). Of the 94 individuals captured and marked, we recorded 49 (52%) recapture events— 55% involving *P. geoffroanus* and 47% involving *M. vanderhaegei*. We did not record recaptures of *T. dorbigni* (Table 2).

The detection probability of *Mesoclemmys vanderhaegei* did not include precipitation or temperature in the best model. However, the second-best model, which included precipitation ($\Delta AIC = 0.05$), and the third-best model, which incorporated temperature ($\Delta AIC = 1.99$), were equally informative for the species. On the other hand, *Phrynops geoffroanus* showed a higher detection probability only with increasing temperature, as both the naïve model and the model including precipitation had $\Delta AIC > 3$, indicating lower support compared to the temperature model ($\Delta AIC = 0$; Table 3; Fig. 3).

The abundance of *M. vanderhaegei* and *P. geoffroanus* varied among water bodies, across sampling months, and also based on the interaction between these two factors (Table 4; Fig. 4, 5). However, the average daily temperature influenced abundance only for *M. vanderhaegei* ($X^2 = 22.676$, $df = 2$, $p < 0.001$; Table 4). When both species were modeled together, the average daily temperature also did not have a significant effect on species abundance ($X^2 = 3.3153$, $df = 1$, $p = 0.068$; Table 4). The variation in abundance over the sampling months was different for *P. geoffroanus* and *M. vanderhaegei*, with high capture rates

in the wettest and warmest months of the year for *P. geoffroanus* (November, December, January, February and March) (Fig. 4). For both species, no new individuals were captured in May (Fig. 4).

When analyzing, year by year (2014, 2015, and 2017), whether the water bodies (A, B, C, and D) and the abundance of *M. vanderhaegei* affected the abundance of *P. geoffroanus*, we observed a significant influence of the area only in 2014 ($X^2 = 1.17$, $df = 3$, $p < 0.001$). However, when all years were analyzed together, the sampled area did not affect the abundance of *P. geoffroanus*, but the abundance of *M. vanderhaegei* did ($X^2 = -0.15$, $df = 3$, $p < 0.001$; Table 5). Conversely, when examining whether the abundance of *P. geoffroanus* or the area affected the abundance of *M. vanderhaegei*, we found that both the area ($X^2 = 2.74$, $df = 3$, $p = 0.01$) and the abundance of *P. geoffroanus* ($X^2 = -0.34$, $df = 3$, $p = 0.01$) influenced *M. vanderhaegei* in 2014, but not in the other years when analyzed individually. When all years were considered together, the area did not significantly affect the abundance of *M. vanderhaegei* ($X^2 = -0.34$, $df = 3$, $p = 0.01$), but the abundance of *P. geoffroanus* did ($X^2 = -0.12$, $df = 3$, $p < 0.001$; Table 5). Abundance was not estimated for 2016, as no individuals of *M. vanderhaegei* were recorded that year. Although the area was not significantly related to the abundance of either species, graphical analysis showed a trend: *P. geoffroanus* tended to be more abundant in water bodies with fewer individuals of *M. vanderhaegei*, and similarly, *M. vanderhaegei* was more abundant where *P. geoffroanus* occurred at lower abundance (Figure 5).

DISCUSSION

In the present study, we found a higher abundance of *Phrynosoma geoffroanus* compared to *Mesoclemmys vanderhaegei*. The abundance of *P. geoffroanus* was higher from November to March, coinciding with the rainy season, while *M. vanderhaegei* showed a peak in abundance in December and remained at low levels during the other months. The probability of detection

was influenced by climatic factors such as precipitation and temperature, but the effect differed between the two most abundant species. Overall, considering all sampling years, the abundance of the two species varied in relation to each other; however, water bodies and sampling months also influenced the individual abundance of each species.

Phrynops geoffroanus was the dominant species in the study. This pattern was expected, since *P. geoffroanus* seems to be more adapted to urban environments (e.g. Souza and Abe, 2000; Souza et al., 2008). *Phrynops geoffroanus* does not appear to be selective regarding habitat quality, using both lentic and lotic water bodies of various sizes (Souza et al., 2008; Magalhães et al., 2015; Silva et al., 2016; Rocha et al., 2025). *Mesoclemmys vanderhaegei*, on the other hand, is often found in oligotrophic, small, lentic and lotic water bodies, in conserved and higher altitude environments (Brito et al., 2009, 2018; Maffei et al., 2016) and, more rarely, in anthropic environments (Marques et al., 2013, 2014).

Precipitation and temperature affected the detection probability of *M. vanderhaegei*, while for *P. geoffroanus*, only temperature influenced its detection probability. However, when abundance was considered, average temperature affected only *M. vanderhaegei*. Freshwater turtles usually respond to climatic variables linked to temperature and humidity (Moll and Moll, 2004; Souza, 2004). Both variables are considered equally important for turtles, as demonstrated in other studies, such as those on flooding cycles for *Podocnemis expansa* (Eisemberg et al., 2016), or temperature for activity of the turtles in general (Plummer, 2003; Pittman and Dorcas, 2009; Tamplin and Cyr, 2011). Chelonians are ectothermic animals that depend directly on the temperature of the environment to regulate their internal temperature. Some species are more active when the environment is warmer (Tamplin and Cyr, 2011) or, less active during warmer periods of the day to avoid overheating (Hailey and Coulson, 1996). Our study showed that, although temperature and rainfall are equally important for turtle species, different responses to these variables may occur depending on the species analyzed.

286 Additionally, it is important to note that our study area is located in a tropical Aw climate zone,
287 characterized by consistently high temperatures and minimal seasonal variation, with dry
288 winters. Therefore, other environmental factors—such as aquatic habitat type and resource
289 availability—may play a more significant role in determining species abundance.

290 *Phrynops geoffroanus* was more abundant in water bodies with lower abundances of *M.*
291 *vanderhaegei*, particularly in water bodies A, B, and D. In contrast, *M. vanderhaegei* reached
292 its highest abundance in water body C, where only a few individuals of *P. geoffroanus* were
293 captured. Behavioral differences, resource use, or morphological traits between the two species
294 may lead to variation in habitat use. Larger species, such as *P. geoffroanus*, generally require
295 more resources and tend to occupy larger home ranges (Perry and Garland, 2002; Jaeger and
296 Cobb, 2012; Müller et al., 2019). Although the total area of these water bodies is relatively
297 similar, they may vary in depth, resource availability, and degree of eutrophication. For
298 instance, *M. vanderhaegei* is frequently recorded in small, oligotrophic water bodies, whereas
299 *P. geoffroanus* appears to be less selective in habitat use, occupying a variety of aquatic
300 environments (Brito et al., 2009, 2018; Souza and Abe, 2000, 2001). However, *P. geoffroanus*
301 is more commonly found in eutrophic rather than oligotrophic systems (Souza and Abe, 2000;
302 Souza et al., 2008; Ferronato et al., 2009; Rocha et al., 2025). In addition, both species are
303 omnivorous, with a tendency toward carnivory, feeding on aquatic insects, fish, and amphibians
304 (Martins et al., 2010; Brito et al., 2016). In general, species tend to partition resources to coexist,
305 and omnivores generally feed opportunistically, according to food availability. However,
306 interspecific competition can determine exclusion through aggressive interactions or through
307 the exploitation of resources with higher protein content by more competitive species (Luiselli,
308 2008).

309 Finally, the presence of *Trachemys dorbigni* does not seem to be a problem for native
310 species yet, due to its low abundance. However, it is an exotic species, found outside its natural

range, and with great potential for dispersal and adaptation, especially in human-altered environments (Rachemys, 2009; Molina et al., 2016; Ciccheto et al., 2018; Fonseca et al., 2021). *Trachemys dorbigni* occurs naturally in Brazil (Rio Grande do Sul), Uruguay, and Argentina, inhabiting large floodplains (Alcalde et al., 2012). If its population increases outside its natural range, it can quickly occupy the niche of native species, leading to a decrease in native species abundance or even local extinction if population control is not taken (Zalba and Ziller, 2007). Exotic species usually have greater niche width (Pérez-Santigosa et al., 2011), competing directly with native species and affecting the structure of assemblages with changes in the foraging process and the dissemination of pathogens (Lowe et al., 2000; Zalba and Ziller, 2007). In the case of *Phrynops geoffroanus*, a recent study conducted on a turtle assemblage in an urban area in southern Brazil indicates that the species may alter its activities in response to the coexistence with invasive turtles, such as *Trachemys dorbigni* and *T. scripta elegans* (Rocha et al., 2025). This adaptation may occur mainly due to its generalist habits and tolerance to anthropogenic pressure (Souza and Abe, 2001; Rocha et al., 2025). However, in the case of *M. vanderhaegei*, less abundant than *P. geoffroanus*, our study indicates that it may be less effective as a competitor and, consequently, less able to compete with *T. dorbigni*.

We show here the syntopic occurrence of *Phrynops geoffroanus* and *Mesoclemmys vanderhaegei* with different patterns of abundance. This varies for both species over time, and the abundance of *P. geoffroanus* is reflected in the abundance of *M. vanderhaegei*. Our findings suggest that *P. geoffroanus* is the dominant species in the study area and, according to other studies in urban areas (Souza and Abe, 2001; Souza et al., 2008; Ferronato et al., 2009; Rocha et al., 2025), should persist in this small protected fragment, as well as in other urban environments near the study area (Souza et al., 2008; Martins et al., 2010; Müller et al, 2019). Despite the challenges of living in fragmented urban environments, the presence of *M. vanderhaegei* suggests that the area still offers suitable conditions for species with higher

habitat quality requirements. Both species seem to have an overlapping trophic niche, and resource partitioning is a hypothesis to be investigated.

ACKNOWLEDGEMENTS

The authors would like to thank Kwok Chiu Cheung, colleagues of the Study Group on Fragment Ecology (EcoFrag) and all field assistants. We also thank the team of the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and the National Center for Research and Conservation of Reptiles and Amphibians (RAN) for the permits (SISBio #45297-1). ESB thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for a scholarship (process 300658/2019-2) to the Programa de Capacitação Institucional (PCI) of the Museu Paraense Emílio Goeldi (MPEG); TFDR thanks CNPq for a scholarship (process 302506/2024-1) to the PCI of the Instituto Nacional de Pesquisas do Pantanal (INPP); CS thanks CNPq for a research fellowship (process 2312303/2018-1).

- 352 Alcalde, L., Derocco, N.N., Rosset, S.D., Williams, J.D. (2012): Southern-most localities of
353 *Trachemys dorbigni* and first record of *Trachemys scripta elegans* for Argentina
354 (Cryptodira, Emydidae). Chelonian Conserv. Biol. **11**: 128-133.
- 355 Alho, C.J., Pádua, L.F. (1982): Sincronia entre regime de vazante do rio e comportamento de
356 nidificação da tartaruga da Amazônia *Podocnemis expansa* (Testudinata: Pelomedusidae).
357 Acta Amazon. **1**: 323-326.
- 358 Balestra, R.A.M., Valadão, R.M., Vogt, R.C., Bernhard, R., Ferrara, C.R., Brito, E.S., Arias,
359 R.B., Malvásio, A., Lustosa, A.P.G., Souza, F.L., Drummond, G.M., Bassetti, L.A.B.,
360 Coutinho, M.E., Ferreira-Junior, P.D., Campos, Z.M., Mendonça, S.H.S.T., Rocha, J.M.N.,
361 Luz, V.L.F. (2016): Roteiro para Inventários e Monitoramentos de Quelônios Continentais.
362 Biodiv. Bras. (BioBrasil). **1**: 114-152.
- 363 Brito, E.S.D., Strüssmann, C., Penha, J.M.F. (2009): Population structure of *Mesoclemmys*
364 *vanderhaegei* (Bour, 1973) (Testudines: Chelidae) in the Cerrado of Chapada dos
365 Guimarães, Mato Grosso, Brazil. Biota Neotrop. **9**: 245-248.
- 366 Brito, E., Souza, F., Strüssmann, C. (2016): Feeding habits of *Mesoclemmys vanderhaegei*
367 (Testudines: Chelidae). Acta Herpetol. **11**: 1-13.
- 368 Brito, E.S., Vogt, R.C., Valadão, R.M., Franco, L.F., Penha, J., Strüssmann, C. (2018):
369 Population ecology of the freshwater turtle *Mesoclemmys vanderhaegei* (testudines:
370 chelidae). Herpetol. Conserv. Biol. **13**: 355-365.
- 371 Bodie, J.R., Semlitsch, R.D. (2000): Spatial and temporal use of floodplain habitats by lentic
372 and lotic species of aquatic turtles. Oecologia **122**: 138-146.

373 Burnham, K.P., Anderson, D.R. (2002): Model Selection and Multi-Model Inference: a
 374 Practical Information- Theoretic Approach. New York, Springer.

375 Cagle, F.R. (1939): A system of marking turtles for future identification. *Copeia* **3**: 170-173.

376 Ciccheto, J.R.M., Grou, C.E.V., Rocha, S.B. (2018): Novos registros de ocorrência de
 377 *Trachemys dorbigni* no Brasil. In: Estudo da Herpetofauna Brasileira Ponta Grossa, pp. 42-
 378 50. Oldiges, D.P., Ed, Atena Editora.

379 Cheung, K.C., Koutchin-Reis, L., Jesus, C.C.C. (2016): Análise fitossociológica de um
 380 fragmento de cerrado em Campo Grande, MS. *Multitemas* **21**: 241-256.

381 Corral, A., Valério, L.M., Cheung, K.C., Ferreira, B.H.S., Guerra, A., Szabo, J.K., Koutchin-
 382 Reis, L. (2021): Plant-bird mutualistic interactions can contribute to the regeneration of
 383 forest and non-forest urban patches in the Brazilian Cerrado. *Urban Ecosyst.* **24**: 205-213.

384 Eiseberg, C.C., Balestra, R.A.M., Famelli, S., Pereira, F.F., Bernardes, V.C.D., Vogt, R.C.
 385 (2016): Vulnerability of Giant South American Turtle (*Podocnemis expansa*) nesting habitat
 386 to climate-change-induced alterations to fluvial cycles. *Trop. Conserv. Sci.* **9**: 1-12.

387 Fachín-Terán, A., Vogt, R.C., Thorbjarnarson, J.B. (2006): Seasonal movements of
 388 *Podocnemis sextuberculata* (Testudines: Podocnemididae) in the Mamirauá sustainable
 389 development reserve, Amazonas, Brazil. *Chelonian Conserv.Biol.* **5**: 18-24.

390 Ferronato, B.O., Merchant, M.E., Marques, T.S., Verdade, L.M. (2009): Characterization of
 391 innate immune activity in *Phrynops geoffroanus* (Testudines: Chelidae). *Zoologia* **26**: 747-
 392 752.

393 Fiske, I., Chanderler, R. (2011): Unmarked: an R package for fitting hierarchical models of
 394 wildlife occurrence and abundance. *J. Stat. Softw.* **43**: 1-23.

395 Fonseca, É., Both, C., Cechi, S.Z., Winck, G. (2021): Pet distribution modelling: Untangling
 396 the invasive potential of *Trachemys dorbigni* (Emydidae) in the Americas. PLoS ONE **16**:
 397 e0259626.

398 Gibbons, J.W., Scott, D.E., Ryan, T.J., Buhlmann, K.A., Tiuberville, T.D., Metts, B.S., Greene,
 399 J.L., Mills, T., Leiden, Y., Poppy S., Winne, C.T. (2000): The Global Decline of Reptiles,
 400 Déjà Vu Amphibians: Reptile species are declining on a global scale Six significant threats
 401 to reptile populations are habitat loss and degradation, introduced invasive species,
 402 environmental pollution, disease, unsustainable use, and global climate change. BioScience
 403 **50**: 6531-666.

404 Guzy, J.C., Price, S.J., Dorcas, M.E. (2013): The spatial configuration of greenspace affects
 405 semi-aquatic turtle occupancy and species richness in a suburban landscape. Landsc. Urban
 406 Plan. **117**: 46-56.

407 Hailey, A., Coulson, I.M. (1996): Temperature and the tropical tortoise *Kinixys spekii*:
 408 constraints on activity level and body temperature. J. Zool. **240**: 523-536.

409 Hoef, J.M., Boveng, P.L. (2007): Quasi-Poisson vs negative binomial regression: how should
 410 we model overdispersed count data? Ecology **88**: 27661-2772.

411 Jaeger, C.P., Cobb, V.A. (2012): Comparative spatial ecologies of female painted turtles
 412 (*Chrysemys picta*) and red-eared sliders (*Trachemys scripta*) at Reelfoot Lake, Tennessee.
 413 Chelonian Conserv. Biol. **11**: 591-67.

414 Liu, Z., He, C., Wu, J. (2016): The relationship between habitat loss and fragmentation during
 415 urbanization: an empirical evaluation from 16 world cities. PLoS ONE **11**(4): e0154613.

416 Lowe, S., Browne, M., Boudjelas, S., De Poorter, M. (2000): 100 of the world's worst invasive
 417 alien species: a selection from the global invasive species database (Vol 12) Auckland:
 418 Invasive Species Specialist Group.

419 Luiselli, L., Akani, G.C. (2003): An indirect assessment of the effects of oil pollution on the
 420 diversity and functioning of turtle communities in the Niger Delta, Nigeria. *Anim. Biodivers.*
 421 *Conserv.* **26**: 57-65.

422 Luiselli, L. (2008): Resource partitioning in freshwater turtle communities: A null model meta-
 423 analysis of available data. *Acta Oecol.* **34**: 80-88.

424 Mackenzie, D.I., Nichols, J.D., Royle, J.A., Pollock, K.H., Bailey, L.L., Hines, J.E. (2006):
 425 Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species
 426 Occurrence. Amsterdam, Elsevier/Academic Press.

427 Maffei, F., Do Nascimento, B.T.M., Moya, G.M., Donatelli, R.J. (2016): New distribution
 428 records of *Mesoclemmys vanderhaegei* (Testudines: Chelidae) from southeastern Brazil,
 429 including observations on reproduction. *J. Threat. Taxa* **8**: 9322-9326.

430 Magalhães, F.M., Laranjeiras, D.O., Costa, T.B., Juncá, F.A., Mesquita, D.O., Röhr, D.L.,
 431 Garda, A.A. (2015): Herpetofauna of protected areas in the Caatinga IV: Chapada
 432 Diamantina National Park, Bahia, Brazil. *Herpetol. Notes* **8**: 243-261.

433 Marques, T.S., Lara, N.R., Bassetti, L.A., Ferronato, B.O., Malvásio, A., Verdade, L.M. (2013):
 434 Population structure of *Mesoclemmys vanderhaegei* (Testudines, Chelidae) in a silvicultural
 435 system in southeastern Brazil. *Herpetol. Notes* **6**: 179-182.

436 Marques, T.S., Böhm, S., Brito, E.S., Cabrera, M.R., Verdade, L.M. (2014): *Mesoclemmys*
 437 *vanderhaegei* (Bour 1973) – Vanderhaege's Toad-headed Turtle, Karumbé-hy. In:
 438 Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the
 439 IUCN/SSC Tortoise and Freshwater Turtle Specialist Group. Rhodin, A.G.J., Pritchard,
 440 P.C.H., Van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., Iverson, J.B., Mittermeier, R.A.
 441 Chelonian Res. Monogr. **5**: 0831-8.

442 Marcuzzo, F.F.N., Melo, D.C.D.R., Rocha, H.M. (2011): Distribuição espaço-temporal e
 443 sazonalidade das chuvas no Estado do Mato Grosso. Rev. Bras. Recur. Híd. **16**: 157-167.

444 Martins, F.I., Souza, F.L., Da Costa, H.T.M. (2010): Feeding habits of *Phrynops geoffroanus*
 445 (Chelidae) in an urban river in Central Brazil. Chelonian Conserv. Biol. **9**: 294-297.

446 Mittermeier, R.A., Van Dijk, P.P., Rhodin, A. G., Nash, S.D. (2015): Turtle hotspots: an
 447 analysis of the occurrence of tortoises and freshwater turtles in biodiversity hotspots, high-
 448 biodiversity wilderness areas, and turtle priority areas. Chelonian Conserv. Biol. **14**: 2-10.

449 Moll, D., Moll, E.O. (2004): The Ecology, Exploitation, and Conservation of River Turtles.
 450 New York, Oxford University Press.

451 Molina, F.B., Ferronato, B.O., Souza, F.L. (2016): Freshwater Turtles in Natural, Rural and
 452 Urban Ecosystems in São Paulo State, Southeastern Brazil. In: Biodiversity in Agricultural
 453 Landscapes of Southeastern Brazil, Berlin. pp. 151-162. Gheler-Costa, C., Lyra-Jorge, M.C.,
 454 Verdade, L.M., Eds, De Gruyter Open.

455 Müller, M.M.P., Ortega, Z., Antunes, P.C., Seino, L.L., Hammarstron, M.J., Balbino-Silva,
 456 A.C.F., Oliveira-Santos, L.G.R. (2019): The home range of adult *Phrynops geoffroanus*
 457 (Testudines, Chelidae) in relation to sex and body mass. Herpetozoa **32**: 259-265.

458 Peel, M.C., Finlayson, B.L., McMahon, T.A. (2007): Updated world map of the Köppen-Geiger
 459 climate classification. Hydrol. Earth Syst. Sci. **11**: 1633-1644.

460 Perry, G., Garland, J.R.T. (2002): Lizard home ranges revisited: effects of sex, body size, diet,
 461 habitat, and phylogeny. Ecology **83**: 1870-1885.

462 Pérez-Santigosa, N., Florencio, M., Hidalgo-Vila, J., Días-Paniagua, C. (2011): Does the exotic
 463 invader turtle, *Trachemys scripta elegans*, compete for food with coexisting native turtles?
 464 Amphibia-Reptilia **32**: 167-175.

465 Pittman, S.E., Dorcas, M.E. (2009): Movements, habitat use, and thermal ecology of an isolated
 466 population of bog turtles (*Glyptemys hlenbergii*). *Copeia* **2009**: 781-790.

467 Plummer, M.V., Mills, N.E., Allen, S.L. (1997): Activity, habitat, and movement patterns of
 468 softshell turtles (*Trionyx spiniferus*) in a small stream. *Chelonian Conserv. Biol.* **2**:514-520.

469 Plummer, M.V. (2003): Activity and thermal ecology of the box turtle, *Terrapene ornata*, at its
 470 southwestern range limit in Arizona. *Chelonian Conserv. Biol.* **4**: 569-577.

471 R DEVELOPMENT CORE TEAM. (2019): R: a language and environment for statistical
 472 computing. Available at [http:// wwwR-projectorg/](http://www.R-project.org/)

473 Rachemys, T. (2009): First record of *Trachemys dorbigni* (Duméril & Bibron, 1835)
 474 (Testudines, Emydidae) in a remnant of Mesophytic Semideciduous Forest of São Paulo
 475 State, southeastern. *Herpetol. Bull.* **108**: 27-30.

476 Rocha, S.B., Rouco, C., Grou, C.E.V., Ferreira, V.L. (2025): Freshwater turtle (Reptilia:
 477 Testudines) assemblage from an urban protected Atlantic Forest remnant: population
 478 density, movement and habitat preference. *Zoologia (Curitiba)* **42**: e24009.

479 Silva, M.B.D., Da Rocha, W.A., Nogueira-Paranhos, J.D. (2016): Checklist of reptiles of the
 480 Amazonia-Caatinga-Cerrado ecotonal zone in eastern Maranhão, Brazil. *Herpetol. Notes* **9**:
 481 7-14.

482 Sano, S.M., Almeida, S.P., Ribeiro, J.F. (2008): Cerrado: ecologia e flora. Brasília, DF:
 483 Embrapa Informação Tecnológica. Planaltina, Distrito Federal, Embrapa Cerrados.

484 Souza, F.L., Abe, A.S. (2000): Feeding ecology, density and biomass of the freshwater turtle,
 485 *Phrynops Geoffroyanus*, inhabiting a polluted urban river in south-eastern Brazil. *J. Zool.* **252**:
 486 437-446.

487 Souza, F.L., Abe, A.S. (2001): Population structure and reproductive aspects of the freshwater
 488 turtle, *Phrynops geoffroanus*, inhabiting an urban river in southeastern Brazil. Stud. Neotrop.
 489 Fauna Environ. **36**: 57-62.

490 Souza, F.L. (2004): Uma revisão sobre padrões de atividade, reprodução e alimentação de
 491 cágados brasileiros (Testudines, Chelidae). Phyllomedusa **3**: 15-27.

492 Souza, F.L., Raizer, J., Costa, H.Y.M., Martins, F.I. (2008): Dispersal of *Phrynops geoffroanus*
 493 (Chelidae) in an urban river in central Brazil. Chelonian Conserv. Biol **7**: 257-61.

494 Spencer, R.J., Dykea, J.V., Petrova, K., Ferronato, B., Mcdougalla, F., Austina, M., Keiteld, C.,
 495 Georges, A. (2018): Profiling a possible rapid extinction event in a long-lived species. Biol.
 496 Conserv. **221**: 190-197.

497 Stokeld, D., Hamer, A.J., Van Der Ree, R., Pettigrove, V., Gillespie, G. (2014): Factors
 498 influencing occurrence of a freshwater turtle in an urban landscape: a resilient species?
 499 Wildl. Res. **41**: 163-171.

500 Tamplin, J.W., Cyr, A.B. (2011): Effects of acclimation and egg-incubation temperature on
 501 selected temperature by hatchling western painted turtles (*Chrysemys picta bellii*). J. Therm.
 502 Biol. **36**: 507-514.

503 Venables, W.N., Ripley, B.D. (2002): Modern applied statistics with S-Plus. New York,
 504 Springer.

505 Zalba, S., Ziller, S.R. (2007): Adaptive management of alien invasive species: putting the
 506 theory into practice. Natureza & Conservação **5**: 86-92.

Table 1. Water bodies located at the Segredo stream spring sampled for freshwater turtle species in an urban area of the Cerrado region, located in Midwest Brazil.

Water bodies	Type	Habitat	Area (m ²)	Surrounding matrix	Vegetation cover	Substrate	Funnel traps
A	Dam	Lentic	7543,6	Riparian forest	Native arboreous plants	Sandy	4
B	Dam	Lentic	9171,6	Riparian forest with exotic vegetation	Exotic arboreous plants	Sandy	4
C	Dam	Lentic	8089,1	Grassland flooded	Herbaceous and aquatic plants	Clay	2
D	Lagoon	Lentic	6542,1	Riparian forest	Native arboreous plants	Sandy	2

Table 2. Total number of captures and recaptures of chelonians recorded in the Segredo stream, located in an urban area of the Cerrado region, located in Midwest Brazil.

Sampled water bodies	<i>Phrynops geoffoanus</i>		<i>Mesoclemmys vanderhaegei</i>		<i>Thachemys dorbigni</i>	
	Capture	Recapture	Capture	Recapture	Capture	Recapture
A	23	10	3	2	0	0
B	23	21	0	2	0	0
C	2	0	14	3	0	0
D	25	9	2	2	2	0
Total	73	40	19	9	2	0

Table 3. Probability models of detection of *Phrynosops geoffroanus* and *Mesoclemmys vanderhaegei* in relation to daily temperature and occurrence of precipitation, considering constant occupancy (Ψ).

Taxon	K	AIC	Δ AIC	AICw	Beta
<i>Mesoclemmys vanderhaegei</i>					
p() Ψ ()	2	198.54	0	0.46	-3.62
p(prec) Ψ ()	3	199.04	0.5	0.36	0.56
p(temp) Ψ ()	3	200.53	1.99	0.17	-0.0088
<i>Phrynosops geoffroanus</i>					
p(temp) Ψ ()	3	490.21	0	0.756	0.103
p() Ψ ()	2	493.44	3.22	0.151	-2.3
p(prec) Ψ ()	3	494.41	4.2	0.093	0.264

Table 4. Fixed-effect parameters in a Generalized Linear Mixed Model (GLMM) accounting for the abundance of *Phrynops geoffroanus*, *Mesoclemmys vanderhaegei* and total turtle abundance in urban vegetation fragment, in relation to spatial effect (water bodies sampled), temporal effect (month sampled and temperature) and their interaction (water bodies sampled*month sampled).

Variable	df	X ²	P
Abundance of <i>Phrynops geoffroanus</i>			
Water bodies	3	3458.4617	<0.001
Month	11	1123.1046	<0.001
Average temperature	1	7.2644	0.007
Interaction water body*month	33	4384.3083	<0.001
Abundance of <i>Mesoclemmys vanderhaegei</i>			
Water bodies	3	12840.363	<0.001
Month	11	11455.841	<0.001
Average temperature	2	22.676	<0.001
Interaction water body*month	33	44687.955	<0.001
Both species			
Water bodies	3	527.0880	<0.001
Month	11	1150.3408	<0.001
Average temperature	1	3.3153	0.068
Interaction water body*month	33	5100.7831	<0.001

Table 5. Fixed-effect parameters in a Generalized Linear Model (GLM) to assess whether the abundance of each turtle species (*Phrynops geoffroanus* and *Mesoclemmys vanderhaegei*) affects the abundance of the other and whether this variation is related to water body area.

Year	Variable	df	X ²	P
Abundance of <i>Phrynops geoffroanus</i>				
2014	Water body area	3	1.17	<0.001
	<i>Mesoclemmys vanderhaegei</i> abundance	3	0.02	0.81
2015	Water body area	3	0.31	0.36
	<i>Mesoclemmys vanderhaegei</i> abundance	3	-0.06	0.46
2017	Water body area	3	-0.66	0.16
	<i>Mesoclemmys vanderhaegei</i> abundance	3	0.03	0.92
Total	Water body area	3	-0.19	0.10
	<i>Mesoclemmys vanderhaegei</i> abundance	3	-0.15	<0.001
Abundance of <i>Mesoclemmys vanderhaegei</i>				
2014	Water body area	3	2.74	0.01
	<i>Phrynops geoffroanus</i> abundance	3	-0.34	0.01
2015	Water body area	3	0.39	0.70
	<i>Phrynops geoffroanus</i> abundance	3	-0.15	0.22
2017	Water body area	3	0.18	0.79
	<i>Phrynops geoffroanus</i> abundance	3	-0.06	0.77
Total	Water body area	3	-0.84	0.18
	<i>Phrynops geoffroanus</i> abundance	3	-0.12	<0.001

Figures legend

Fig. 1. Water bodies sampled, located at the Segredo stream spring, in forest fragments of an urban area of Midwest Brazil.

Fig. 2. Frequency distribution of size classes of *Phrynops geoffroanus* and *Mesoclemmys vanderhaegei* captured at the Segredo stream spring, in forest fragments of an urban area of Midwest Brazil.

Fig. 3. Detection estimates for *Mesoclemmys vanderhaegei* and *Phrynops geoffroanus* in relation to daily temperature and precipitation. Positive beta values indicate higher detections and negative lower detections of the species. When the confidence intervals do not overlap the dotted line, we consider that the environmental variables interfere with species detection.

Fig. 4. Abundance of *Phrynops geoffroanus* and *Mesoclemmys vanderhaegei* in the months sampled at the Segredo stream spring, Midwest, Brazil.

Fig. 5. Abundance of *Phrynops geoffroanus* and *Mesoclemmys vanderhaegei* in three artificial dams (A, B and C) and one natural pond (D) sampled at the Segredo stream spring, Midwest Brazil. Description of each water body (A, B, C, D) is shown in Table 1.

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

