

Age structure of a population of *Barbarophryne brongersmai* (Hoogmoed 1972) (Anura, Bufonidae) inhabiting an arid environment in the Central Jbilets (West-Morocco)

ABDERRAZZAK FATTAH¹, TAHAR SLIMANI¹, EL HASSAN EL MOUDEN¹, ODILE GROLET², PIERRE JOLY^{2,*}

¹Cadi Ayyad University, Faculty of Sciences Semlalia, Biodiversity and Ecosystems Dynamic Laboratory, PO Box 2390, Marrakech 40 000, Morocco

²Lyon1 University, UMR 5023 Ecology of Natural and anthropized Hydrosystems, ENTPE, CNRS, bât. Darwin C, 69622 Villeurbanne, France. *Corresponding author. E-mail: pierre.joly@univ-lyon1.fr

Submitted on 2014, 7th February; revised on 2014, 14th May; accepted on 2014, 18th August
Editor: Giovanni Scillitani

Abstract. We estimated age structure and some life-history traits in a population of *Barbarophryne brongersmai* at the northern border of the distribution area of this species by using skeletochronology to establish individual age and age at sexual maturity. Maturity was reached at 2-4 years in males and 3-5 years in females, which was late relatively to small body size. Longevity was estimated to be 12 and 8 years for males and females, respectively, whereas mean body size is 45.6 mm in both sexes. However, this estimate suffers from small sample size in females. We did not detect a positive relationship between age and size, suggesting that adult individuals cease growing after maturity. Conversely, body size varied within an age class, suggesting great variability of growth rate among individuals during the juvenile stage. Collectively, these traits characterize a life history strategy that could be an evolutionary response to habitat unpredictability in arid regions.

Keywords. Endemic species, skeletochronology, life history traits, arid environments, Morocco

In arid zones, the reproduction of the amphibians is exposed to high environmental stochasticity because both the amount of rainfall and the hydroperiod of the breeding sites are usually variable and unpredictable. We assume that increased probability of reproduction failure due to pond drying is compensated by higher longevity as suggested by life history theory (Roff, 1992) and already shown in amphibians (Joly and Morand, 1994; Church et al., 2007; Cayuela et al., 2014). In the perspective of increasing the data set about this issue, we investigated the age structure of a small-bodied toad, *Barbarophryne brongersmai*, at the northern edge of the Sahara desert to compare it with those established in other Bufonidae breeding in less unpredictable environments.

Described and separated from *Bufo viridis* by Hoogmoed (1972), *Barbarophryne brongersmai* is a small-

bodied toad endemic to south-western Morocco (in the present paper, we consider recent taxonomic revisions of the *Bufo* genus by Pyron and Wiens, 2011, and Beukema et al., 2013). The studied population inhabits an arid grassland with sparsely scattered jujube trees (*Ziziphus lotus*), located 25 km north of Marrakech in the central Jbilets (31°37'N, 8°02'W), a range of low hills near the northern border of the distribution area of the species (Guillon et al., 2004). In this region, annual precipitation rarely reaches 300 mm (Le Houérou, 2001) and rare rain events of short duration and high intensity provoke sudden water discharges in the wadis (temporary rivers), usually occurring from the end of February to mid-April. The studied area is heavily grazed by large herds of sheep and goats in permanent movement that keep the grass very short, even during the rainy season. Our popula-

tion breeds at a temporary pond (15 m in diameter and 80 cm depth) that appears in the bed of a temporary river (Defla wadi), the hydroperiod of which did not exceed 18 days from 2009 to 2012 because of strong evaporation and intensive use for livestock watering.

Because the toads come to the pond immediately after it fills with water (Guillon et al., 2004; Fattah, 2008; Slimani et al., 2012), sampling occurred from February to April, at night (from 20 to 00h GMT) using hand nets. We measured the toads from snout to cloaca by means of a calliper and weighed them using a field balance (accuracy 0.1g). We determined gender by the presence of nuptial callosities and calling in males. After measurements, we collected the third toe of the hind leg and we immediately released the toad at the place of its capture. We sectioned phalanges in order to quantify numbers of lines of arrested growth (LAGs), following the method by Castanet and Smirina (1990) based on the detection of line of arrested growth generated by the cold and/or dry season (each LAG is interpreted as 1 year of age). The studies by Barbault et al. (1979) on "*Bufo*"*pentoni*, Esteban et al. (1999) on *Pelophylax saharicus*, and Doglio et al. (2008) on *B. brongersmai* have confirmed that long bones reliably record yearly growth arrest in arid regions bordering Sahara desert. However, this method is suspected to regularly underestimate age because of bone remodelling and rapprochement of peripheral lines, especially in the oldest individuals (Wagner et al., 2011).

We made sections in the medium part (diaphysis) of the third phalanx because the two distal phalanges are known to exhibit strong medullary resorption (Francillon-Vieillot, 1987; Fretey and Le Garff, 1992). All the sections where age estimation was doubtful (because of medullary resorption or other cause) were discarded from the results. Three observers (A.F., O.G. and P.J.) independently counted the LAGs of each individual and re-examined the specimen and discussed the results when their estimations diverged. When the result remained uncertain, the specimen was not taken into account. Age at maturity was estimated by the sudden and permanent rapprochement of the LAGs that reflects the slowing of growth rate after sexual maturity is reached.

Data normality was first assessed (Kolmogorov-Smirnov test) before testing for difference between means with Mann-Whitney U test, while correlation between age and body size was assessed by Spearman test. Means are reported with standard deviation and tests were considered significant at $\alpha = 0.05$.

During the 4-years survey, we sampled 80 individuals (71 males and 9 females, sex ratio = 0.89), but we succeeded in establishing age in only 37 males and 6 females. The bone sections show clear lines of arrested growth (LAGs) that are almost circular and continuous in most individuals (68.6%, Fig. 1). We considered the perimeter of the bone as corresponding to the last winter since the toads were sampled in February, March and




Fig. 1. Transverse sections within the diaphysal region of phalanx bones in the studied population. (A) Male with 6 visible LAGs (CM: medullary cavity; LM: line of metamorphosis). (B) Male with 9 visible LAGs. (C) Female with 8 visible LAGs.

April. We clearly detected the decrease in the distance that separates the LAGs suggesting that sexual maturity is reached at 2-4 years in males, and 3-5 years in females (Fig. 1B, C), although their low number prevents any firm conclusion. At the breeding site, we sampled some males younger than 4 years, although 4-year individuals were the most frequently observed (Fig. 2). Highest estimated age is 12 years in males and 8 years in females.

Body size varied from 35.5 to 52.2 mm in males (45.4 ± 2.9 mm, $n = 37$) and from 40.0 to 50.5 mm in females (45.6 ± 4.2 mm, $n = 6$), with a mass of 6.5 ± 1.2 g in males and 7.4 ± 1.9 g in females and the observed pattern of sexual dimorphism in body size was not significant (Mann-Whitney U -test: $U = 110.5$, $Z = 0.017$, $P = 0.10$). Because we did not detect a significant relationship between age and body size in males ($r = -0.073$, $P = 0.667$, $n = 37$), or in females ($r = 0.529$, $P = 0.279$, $n = 6$) (Fig. 3), the oldest individuals were not always the largest ones.

Our study provides data about population structure that brings new light on the adaptive strategy of this species, despite the low confidence of our conclusions about growth and longevity in females, because of the low numbers of sampled individuals. This apparent bias in sex ratio could be due to (i) higher detectability of males because of conspicuous breeding behaviour, (ii) longer stay at breeding site, (iii) visiting the breeding location too early after rainfalls for the females to be present, and (iv) the possibility that the reproductive cycle of females is biennial when that of males is annual. Sexual maturity is reached after about 3-4 years in most males and 5 years in females. However age structure in males

is skewed since we would expect the 2 and 3 years-old toads to be more numerous than those aged 4 as a result of mortality. A first hypothesis for explaining this gap in young males is that not all the males reach sexual maturity before reaching 4 years, whereas a second one is that some males reach sexual maturity when two years old. We do not have enough evidence to conclusively reject either of these hypotheses.

The one-year difference in age at maturity between males and females is commonly observed in the anurans, and is usually attributed to the higher investment of females in gametes (Miaud et al., 1999; Bastien and Leclair, 1992; Kyriakopoulou-Sklavounou and Grumiro, 2002; Eaton et al., 2005). However, males and females can reach sexual maturity at the same age in some toads and other anurans (Lykens and Forester, 1987; Friedl and Klump, 1997; Kyriakopoulou-Sklavounou et al., 2008), probably when resources are abundant. The late age at maturity in our population suggests a low growth rate in juveniles, strongly constrained by the shortness of the activity period and perhaps the scarcity of food. As a consequence, the difference in age at maturity between genders is not surprising in such an environment. However, the low number of female sampled prevents any firm conclusion about this supposed lack of difference between genders.

The absence of a relationship between age and body size is surprising because anurans often exhibit continuous growth, particularly females because of the strong relationship between fecundity and body size (Joly, 1991; Prado and Haddad, 2005; Lengagne et al., 2007). Increasing fecundity at the expense of reducing somatic growth is theoretically possible, leading thus females keeping a




Fig. 2. Frequency of *Barbarophryne brongersmai* at Defla wadi in 2009-2012 according to estimated age in the studied population. Filled bars: females; open bars: males.




Fig. 3. Relationship between estimated age (X-axis in year) and body size (Y-axis; snout-vent length in mm) in the studied population. Dark dots: females; open dots: males.

Table 1. Comparison of longevity and age at maturity in several Bufonids.

Species	Longevity (year)	Age at maturity (year)	Body size (average mass of the species/ average mass of <i>Barbarophryne brongersmai</i>)	Remarks	Data source
<i>Bufo bufo</i>	8	2-4	3 to 4	Higher at elevation >1800m	Hemelaar, 1988; Frazer, 1966; Fretey, 1995; Gittins et al., 1980; Gittins et al., 1982
<i>Epidalea calamita</i>	9	2-3	2	Higher at elevation > 2200 m	Stevens et al., 2003; Leskovar et al., 2006; Oromi et al., 2012
<i>Bufo viridis</i>	9	2-5	2	-	Sinsch et al., 2004
<i>Anaxyrus hemiophrys</i>	9	1-2	-	-	Eaton et al., 2005

constant body size while increasing reproductive investment when ageing (Castellano et al., 2004). However, such a theoretical option of resource allocation dynamics is not common in the amphibians that usually increase fecundity along life through indeterminate growth. Furthermore, the absence of sexual dimorphism in body size also suggests a selection for low annual fecundity that could be compensated by a longer lifespan. In males, the observed longevity exceeds by 4 years that estimated in Jbel Saghro (Anti Atlas, Morocco) by Doglio et al. (2008). This greater longevity in the Central Jbilets could be due to greater sampling effort in our population. By visually extrapolating the decreased number of individuals as age increases, we predict that life expectancy could reach 13-14 years in the studied population. Longevity (potential lifetime under optimal life conditions) should be longer first because skeletochronology usually underestimates current age due to bone remodelling and rapprochement of peripheral lines (Wagner et al., 2011) and second because natural populations are exposed to external causes of mortality (predation, disease, drought, decreased food availability) that reduce life expectancy. In other anurans, longevity estimated in the field is around threefold lower than that measured under optimal rearing conditions. In *Bombina variegata* for example, longevity reaches 29 years in artificially reared individuals when it does not exceed 12-13 years in the field (Plytycz and Bigaj, 1993; unpubl. res.). We thus suspect *B. brongersmai* to have a potential longevity much higher than expected on the basis of its body size. This hypothesis is supported by the fact that most toad species with body size larger than that of *B. brongersmai* exhibit shorter longevity and smaller age at maturity even in species with higher body size (Table 1). These comparisons suggest that small-bodied *B. brongersmai* is characterized by slow growth during the juvenile stage.

The interplay of high longevity, slow growth during the juvenile stage, absence of continued growth at

the adult stage, and small body size constitutes a specific demographic strategy. We assume that this life history style constitutes a response to the unpredictability of breeding habitats and the shortness of activity period by increasing the number of lifetime breeding occasions (Seger and Brockmann, 1987; Roff, 1992). The test of this hypothesis needs to establish age structure in other populations, including large numbers of females, and to investigate physiological mechanisms that support the great longevity of this species.

ACKNOWLEDGEMENTS.

This work has received the support of Hassan II Academy of Sciences and Technics (ICGVSA project). It was authorized by the High Commissariat for Water and Forest (Morocco), Scientific Permits 98/2009, 15/2010 and 234/2012 HCEFLCD/DLDCP/

REFERENCES

- Barbault, R., Castanet, J., Francillon, H., Ricqlès, A.D. (1979): Détermination de l'âge chez un Anoure déserticole *Bufo pentoni* Anderson, 1893. Rev. écol. (Terre Vie) **33**: 129-141.
- Bastien, H., Leclair, R. (1992): Aging wood frogs (*Rana sylvatica*) by skeletochronology. J. Herpetol. **26**: 222-225.
- Beukema, W., De Pous, P., Donaire-Barroso, D., Bogaerts, S., Garcia-Porta, J., Escoriza, D., Arribas, O.J., El Mouden, E.H., Carranza, S. (2013): Review of the systematics, distribution, biogeography and natural history of Moroccan amphibians. Zootaxa **3661**: 1-60.
- Castanet, J., Smirina, E.M. (1990): Introduction to the skeletochronological method in amphibians and reptiles. Ann. Sci. Nat, Zool. Biol. Anim. **11**: 191-196.

- Castellano, S., Cucco, M., Giacoma, C. (2004): Reproductive investment of female green toads (*Bufo viridis*). *Copeia* **2004**: 659-664.
- Cayuela, H., Besnard, A., Bonnaire, E., Perret, H., Rivoalen, J., Miaud, C., Joly, P., 2014. To breed or not to breed: past reproductive status and environmental cues drive current breeding decisions in a long-lived amphibian. *Oecologia* **176**: 107-116
- Church, D.R., Bailey, L.L., Wilbur, H.M., Kendall, W.L., Hines, J.E. (2007): Iteroparity in the variable environment of the salamander *Ambystoma tigrinum*. *Ecology* **88**: 891-903.
- Doglio, S., Seglie, D., Kabiri, L. (2008): First skeletochronology results of *Bufo (Epidalea) brongersmai* from the Moroccan Anti Atlas (Anura, Bufonidae). In: *Herpetologia Sardiniae*, pp. 226-230. Corti, C., Ed., Societas Herpetologica Italica, Edizioni Belvedere, Latina.
- Eaton, B.R., Paszkowski, C.A., Kristensen, K., Hiltz, M. (2005): Life-history variation among populations of Canadian toads in Alberta, Canada. *Can. J. Zool.* **83**: 1421-1430.
- Esteban, M., Garcia-Paris, M., Buckley, D., Castanet, J. (1999): Bone growth and age in *Rana saharica*, a water frog living in a desert environment. *Ann. Zool. Fenn.* **36**: 53-62.
- Fattah, A. (2008): Contribution à la connaissance de l'écologie de *Bufo brongersmai* (Amphibia, Anura, Bufonidae) dans les Jbilettes centrales (Maroc occidental). Implication en termes de conservation. Mémoire de Master, Faculté des Sciences Semlalia, Marrakech.
- Francillon-Vieillot, H. (1987): La croissance des os longs chez les Amphibiens et son utilisation comme critère d'âge. PhD thesis, Université Paris VII, France.
- Frazer, J.F.D. (1966): A breeding colony of toads (*Bufo bufo* L.) in Kent. *Brit. J. Herpetol.* **3**: 236-252.
- Fretey, T., Le Garff, B. (1992): Apports de la squeletochronologie dans la démographie du crapaud commun, *Bufo bufo* (L.) (Anura, Bufonidae) dans l'Ouest de la France. In: *Tissus durs et âge individuel des vertébrés*, pp. 409-419. Baglinière, J.L., Castanet, J., Conand, F., Meunier, F.J., Eds., ORSTOM-INRA, Paris, France.
- Fretey, T. (1995): Biologie d'une population reproductrice de crapaud commun, *Bufo bufo* (Linné, 1758) en forêt de Rennes (Ille-et-Vilaine, France). PhD thesis, Université de Rennes, France.
- Friedl, T.W.P., Klump, G.M. (1997): Some aspects of population biology in the European treefrog *Hyla arborea*. *Herpetologica* **53**: 321-330.
- Gittins, S.P., Parker, A.G., Slater, F.M. (1980): Population characteristics of the common toad (*Bufo bufo*) visiting a breeding site in mid-Wales. *J. Anim. Ecol.* **49**: 161-173.
- Gittins, S.P., Steeds, J.E., Williams, R. (1982): Population age-structure of the common toad (*Bufo bufo*) at a lake in mid-Wales determined from annual growth rings in the phalanges. *Brit. J. Herpetol.* **6**: 249-252.
- Guillon, M., Le Liard, G., Slimani, T. (2004): Nouvelles données sur la répartition et l'écologie de reproduction de *Bufo brongersmai*, *Bufo mauritanicus* et *Bufo viridis* (Anoura, Bufonidae) dans les Jbilettes centrales, Maroc occidental. *Bull. Soc. Herp. Fr.* **111-112**: 37-48.
- Hemelaar, A.S.M. (1988): Age, growth and other population characteristics of *Bufo bufo* from different latitudes and altitudes. *J. Herpetol.* **22**: 369-388.
- Hoogmoed, M.S. (1972): On a new species of toad from Southern Morocco. *Zool. Meded.* **47**: 49-67.
- Joly, P. (1991): Interdemic variations of size and fecundity in the common frog, *Rana temporaria*. *Alytes* **9**: 79-88.
- Joly, P., Morand, A. (1994): Theoretical habitat templates, species traits, and species richness: amphibian in the Upper Rhône and its floodplain. *Freshwater Biol.* **31**: 455-468.
- Kyriakopoulou-Sklavounou, P., Grumiro, I. (2002): Body size and age assessment among breeding populations of the tree frog *Hyla arborea* in northern Greece. *Amphibia-Reptilia* **23**: 219-224.
- Kyriakopoulou-Sklavounou, P., Stylianou, P., Tsiora, A. (2008): A skeletochronological study of age, growth and longevity in a population of the frog *Rana ridibunda* from southern Europe. *Zoology* **111**: 30-36.
- Le Houérou, H. N. (2001). Biogeography of the arid steppe-land north of the Sahara. *J. Arid Environ.*, **48**: 103-128
- Lengagne, T., Arthaud, F., Cormier, M., Joly, P. (2007): Cost of sexually embracing a large female offset by the number of eggs fertilised for small male *Bufo bufo* L. *Biol. J. Linn. Soc.* **92**: 755-762.
- Leskovar, C., Oromi, N., Sanuy, D., Sinsch, U. (2006): Demographic life history traits of reproductive natterjack toads (*Bufo calamita*) vary between northern and southern latitudes. *Amphibia-Reptilia* **27**: 365-375.
- Lykens, D.V., Forester, D.C. (1987): Age structure in the spring peeper: do males advertise longevity? *Herpetologica* **43**: 216-223.
- Miaud, C., Guyétant, R., Elmberg, J. (1999): Variations in life history traits in the common frog *Rana temporaria* (Amphibia, Anura): a literature review and new data from the French Alps. *J. Zool., London* **249**: 61-73.
- Oromi, N., Sanuy, D., Sinsch, U. (2012): Altitudinal variation of demographic life-history traits does not mimic latitudinal variation in natterjack toads (*Bufo calamita*). *Zoology* **115**: 30-37.

- Plytycz, B., Bigaj, J. (1993): Studies on the growth and longevity of the yellow-bellied toad, *Bombina variegata*, in natural environments. *Amphibia-Reptilia* **14**: 35-44.
- Prado, C.P.A., Haddad, C.F.B. (2005): Size-fecundity relationships and reproductive investment in female frogs in the Pantanal, south-western Braz. *Herp. J.* **15**: 181-189.
- Pyron, R.A., Wiens, J.J. (2011): A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of extant frogs, salamanders, and caecilians. *Mol. Phylogenet. Evol.* **61**: 543-583.
- Roff, D.A. (1992): The evolution of life histories: theory and analysis. Chapman and Hall, London.
- Seger, J., Brockmann H. J. (1987): What is bet-hedging? In: Oxford surveys in evolutionary biology, p. 182-211. Harvey, P. H., Partridge, L., Eds., Oxford University Press, Oxford, UK.
- Sinsch, U., Leskovar, C., Drobig, A., König, A., Grosse, W.R. (2004): Life-history traits in green toad (*Bufo viridis*) populations: indicators of habitat quality. *Can. J. Zool.* **85**: 665-673.
- Slimani, T., Fattah, A., El Mouden, E.H., Radi, M., Doglio, S. (2012): Ecology and population parameters of *Bufo (Pseudepidalea) brongersmai* (Hoogmoed, 1972) living in small temporary ponds in the arid area of Jbilets (Western Morocco). Summary in Seventh World Congress of Herpetology [WCH 7], p. 666-667. Vancouver, Canada, August 8-14.
- Stevens, V., Wesselingh, R., Baguette, M. (2003): Demographic processes in a small, isolated population of natterjack toads (*Bufo calamita*) in southern Belgium. *J. Herpetol.* **12**: 59-67.
- Wagner, A., Schabetsberger, R., Sztatecsny, M., Kaiser, R. (2011): Skeletochronology of phalanges underestimates the true age of long-lived Alpine newts (*Ichthyosaura alpestris*). *Herpetol. J.* **21**: 145-148.