

Herpetofaunal diversity in the Upper Guinean rainforest: A baseline survey from the Dugbe region, Liberia

RYAN VAN HUYSSTEEN, MELISSA PETFORD, MARIUS BURGER, MARK-
OLIVER RÖDEL, TROKON SUPERIOR GRIMES, AND JEROME GAUGRIS

THIS ARTICLE HAS BEEN ACCEPTED FOR PUBLICATION AND UNDERGONE FULL
PEER REVIEW BUT HAS NOT BEEN THROUGH THE COPYEDITING, TYPESETTING,
PAGINATION AND PROOFREADING PROCESS, WHICH MAY LEAD TO DIFFERENCES
BETWEEN THIS VERSION AND THE VERSION OF RECORD.

PLEASE CITE THIS ARTICLE AS:

Van Huyssteen, R., Petford, M., Burger, M., Rödel, M.-O., Superior Grimes, T., Gaugris, J. (2026): Herpetofaunal diversity in the Upper Guinean rainforest: A baseline survey from the Dugbe region, Liberia. *Acta Herpetol.* **21**. doi: 10.36253/a_h-18279.

Herpetofaunal diversity in the Upper Guinean rainforest: A baseline survey from the Dugbe region, Liberia

RYAN VAN HUYSSTEEN¹, MELISSA PETFORD¹, MARIUS BURGER^{1, 2}, MARK-OLIVER RÖDEL³, TROKON SUPERIOR GRIMES⁴, and JEROME GAUGRIS¹.

25 *I FLORA FAUNA & MAN, Ecological Services Ltd, Trident Chambers, P.O Box 146, Road*
26 *Town, 1110 – Tortola, British Virgin Islands.*

27 *2 African Amphibian Conservation Research Group, Unit for Environmental Sciences and*
28 *Management, North-West University, 2520 – Potchefstroom, South Africa.*

29 3 Museum für Naturkunde – Leibniz Institute for Evolution and Biodiversity Science,
30 Invalidenstrasse 43, 10115 Berlin, Germany

31 4 Forestry Development Authority, When Town, Mont. Barclay P. O. Box 10-3010 –
32 Monrovia, Liberia.

33 **Corresponding Author:** Ryan van Huyssteen, *Trident Chambers, Po Box 146, Road Town,*
34 *1110 – Tortola, British Virgin Islands* rv.huyssteen@florafaunaman.com

35 Submitted on: 2025, 22th July; revised on: 2025, 31th July accepted on: 2026, 16th January.

36 *Editor: David Andrew Beamer*

37

38 **Abstract.** The upper Guinean rainforest biome is a poorly studied, yet hyper-biodiverse
39 region facing severe fragmentation due to ongoing habitat transformation. We conducted a
40 herpetological baseline survey in the Dugbe region during 2021 using passive trapping and
41 active searching. Our survey resulted in 1145 herpetofauna observations, representing 71 taxa
42 (38 amphibians and 33 reptiles), with active searching yielding 47.8% of the species richness.
43 Nearly half (54.9%) of the documented species are West African endemics.
44 Rarefaction/extrapolation sampling curves indicated incomplete overall sampling of
45 herpetofauna diversity, though amphibian sampling completeness was high. Reptile diversity

46 metrics revealed significant sampling deficiencies, largely explained by the high proportion
47 of singleton observations (60.6% of reptile records). When compared to the IUCN predictive
48 distribution maps, our survey documented 51.2% of the 123 species predicted for the region,
49 with moderate overall Jaccard similarity (48.09%). Taxonomic groups showed varied patterns
50 of congruence: amphibians displayed relatively high Jaccard similarity (59.26%), while
51 reptiles showed lower similarity (40.26%). Notably, our survey documented substantially
52 higher herpetofauna species richness than benchmark surveys from nearby areas, particularly
53 for reptiles (33 species compared to 14 and 5 species in Krahn-Bassa proposed protected area
54 and Grebo National Forest surveys, respectively). Several amphibian observations were not
55 assignable to species using morphological characteristics alone, highlighting the need for
56 further taxonomic research on West African herpetofauna.

57 **Keywords.** Reptiles, Amphibians, West Africa, Field Survey, Herpetology, Tropical Forest
58 Ecology


INTRODUCTION

60 The Upper Guinean forests of West Africa are a global biodiversity hotspot with high faunal
61 and floral diversity and endemism (Myers et al., 2000; De Sousa et al., 2023; Lewin et al.,
62 2016; Ernst et al., 2025). These forests are important for amphibians, with high species richness
63 and taxonomic diversity (Penner et al., 2011, 2019; Nneji et al., 2023). The region's reptiles
64 are less well studied than amphibians, and there are significant gaps in our understanding of
65 their diversity and distribution (Tolley et al., 2016); this is a globally observed pattern in the
66 context of tropical forest herpetofauna (Deikumah et al., 2014; Tolley et al., 2016). The Upper
67 Guinean forests which once spanned continuously across West Africa have been fragmented
68 by deforestation, agricultural expansion, and mining (Sodhi et al., 2008, Hepner, 2025). Liberia
69 still holds some of the largest remaining tracts of these forests, but habitat fragmentation and
70 degradation have intensified in the last twenty years (De Sousa et al., 2023). This loss of habitat
71 threatens the region's herpetofauna (Rödel et al., 2021) and baseline surveys are needed to
72 document amphibian and reptile biodiversity, as loss or extinctions may be occurring without
73 our knowledge.

75 Although there have been numerous studies on Liberian herpetofauna (e.g., Barbour &
76 Loveridge, 1927; Loveridge, 1941, Angel, 1943a; Guibé & Lamotte, 1958;), particularly in the
77 Mount Nimba region (e.g., Xavier, 1978; Lamotte, 1998; Ineich, 2003; Rödel et al., 2009;
78 Sandberger et al., 2010), large portions of the country remain poorly surveyed. Our
79 understanding of herpetological species distributions is based on local surveys, species
80 descriptions, and taxonomic revisions (e.g., Trape & Mané, 2006; Hillers & Rödel, 2007;
81 Blackburn et al., 2008; Allen et al., 2019; Rödel & Glos, 2019). Comprehensive baseline

82 surveys from understudied regions can address these knowledge gaps, and guide conservation
83 efforts in response to ongoing habitat loss.

84
85 This study provides the first herpetological species inventory for the Dugbe region, located in
86 southern Liberia (Fig. 1). The survey was conducted in 2021 by Flora, Fauna and Man,
87 Ecological Services Ltd. (FFMES) and the aim was to establish a baseline of herpetological
88 diversity, identify species of conservation concern, and find potential novel or endemic species.
89 To contextualise our findings, we compare them with surveys conducted approximately 100
90 km away in Grebo National Forest (Hillers & Rödel, 2007) and the proposed Krahn-Bassa
91 proposed protected area (KBPPA) (Rödel & Glos, 2019). Both areas have similar ecological
92 and biogeographical features to the Dugbe study area and our comparisons focus on species
93 richness to provide an overview of the regional species diversity for this poorly studied area.



94
95 **Fig. 1.** Location of the Dugbe study area within Liberia and the locations of the herpetofauna
96 observations and trap arrays within the study area.

97 The Dugbe study area is located within the Upper Guinean Rainforest biome, near to Sapo
98 National Park and the Grand Kru–River Gee proposed protected area. Historically the
99 landscape was dominated by primary closed-canopy rainforest (Cushing et al., 2016; Ernst et
100 al., 2025). The landscape has high conservation significance and is currently characterised by
101 a mosaic of natural and transformed habitat types with most of the forested areas in various
102 stages of secondary succession. Over 14% of mature forest cover was lost between 2000 and
103 2018 due to ongoing habitat transformation and resource exploitation (De Sousa et al., 2023).
104 Several villages are present within the study area and there is a high reliance on subsistence
105 agriculture and artisanal mining. The study area consists of low hills with some rocky areas,
106 and the elevation ranges between 10–170 m above sea level. The region is drained by three
107 major rivers: the Boutu, Dugbe, and Geebo, as well as smaller streams.

108 **METHODS**

109 *Field surveys*

110 Surveys were conducted over two periods: an initial five-day scoping visit from 8–13 May
111 2021 during the wet season, and a follow-up 16-day survey from 20 October to 4 November
112 2021, during the transitional period between the wet and dry seasons. The first survey was
113 undertaken by a single herpetologist (Marius Burger) and involved diurnal and nocturnal active
114 searches to explore the terrain in preparation for a more detailed survey. The second survey
115 was conducted by two herpetologists (Marius Burger and Ryan van Huyssteen) and included
116 diurnal and nocturnal active searching and the use of passive trapping methods. Although the
117 second season was planned for the transitional period between the wet and dry seasons, there
118 was heavy rainfall for most of the survey period.

119

120 The sampling strategy followed a multistage stratified semi-random sampling (MSSRS)
121 approach (Bourgeron et al., 2001, Mottram et al., 2025). The study area was divided into 576
122 grid cells of 1 km² each. Within each grid cell, the MSSRS approach was used to semi-
123 randomly identify specific sampling sites through consideration of physiognomic and
124 physiographic elements (Mottram et al., 2025). In addition to predefined points, opportunistic
125 searches were conducted in microhabitats that are known to be ecological relevant for particular
126 species (e.g., ponds, streams, rocky outcrops). In total we investigated 64 grid cells and
127 sampled 133 sample points within those. Active searching involved looking for resting and
128 active herpetofauna, investigating likely hiding places (holes, under logs and other cover,
129 raking through leaf litter, etc.), and listening for vocalising frogs.

130

131 Acoustically detected amphibians were recorded as presence/absence data (i.e., single presence
132 per calling event regardless of the number of individuals estimated to be calling). This was a
133 methodological decision implemented to prevent potential bias in species accumulation
134 analyses when pooling acoustic-detected species and visual-detected species. However, for all
135 amphibians and reptiles directly observed during surveys, we maintained accurate count data
136 reflecting the true number of individuals encountered.

137

138 The trapping effort consisted of six trap arrays that each had four 25-litre bucket pitfall traps
139 and six double-ended funnel traps connected by drift fences assembled in a linear configuration
140 (Fig. 2). Traps arrays were operational for 13 nights (78 trap array nights/780 trap nights),
141 trapped specimens were recorded and removed daily.

142



143

144 **Fig. 2.** One of six trap arrays installed for the Dugbe herpetofauna survey. The trap array is
145 composed of six funnel traps (A) and four pitfall traps (B) that are installed along a 30-meter
146 drift fence (C) in a linear fashion.

147

148 Incidental observations, including photographs and specimens recorded by researchers of other
149 taxonomic groups and mining personnel during the survey period were also included and
150 documented. Most observations had a GPS accuracy of 5 m; however, several incidental
151 records had a lower resolution of 5000 m.

152

153 Reptiles and amphibians were identified based on the keys provided by Chippaux (2006), Trape
154 et al. (2012), Schiøtz (1999) and descriptions in Channing and Rödel (2019). In some instances,
155 specialists were consulted for advice on identifications (see acknowledgements). Where

156 possible observations were supported by photographic or sound-clip evidence and these were
157 deposited as digital vouchers onto iNaturalist.org (Supplementary material 1). Voucher
158 specimens and genetic samples were collected for most species encountered and deposited at
159 the American Museum of Natural History (AMNH) (Supplementary material 1). Amphibian
160 taxonomy follows Frost (2025), and common names following Channing and Rödel (2019),
161 while reptile taxonomy is based on Uetz et al. (2025).

162

163 *Survey evaluation*

164 We used three methods to evaluate our survey effort: firstly, we used rarefaction/extrapolation
165 (r/e) curves, secondly, we compared our survey to the IUCN (International Union for
166 Conservation of Nature) predictive species lists for the study area which are based on expert-
167 derived range maps, and finally we compared our survey species richness to two benchmark
168 surveys that were conducted within 100 km of our study area. All herpetofauna observations
169 made during the surveys, including incidental records, were included in our analyses.
170 Amphibian taxa that were identified as recognisable taxonomic units, but not confidently
171 assignable to currently known species, were also included in the sampling curves, diversity
172 estimates, and similarity analyses.

173

174 *Sampling curves.* To evaluate survey effort in the Dugbe area, r/e curves were generated using
175 the iNEXT package (Hsieh et al., 2016) in R (R Core Team, 2022) via the RStudio console
176 (RStudio Team, 2022). The curves were plotted and extrapolated to twice the original survey
177 effort, with observation sequences randomised 100 times using bootstrap resampling.
178 Estimated species richness, along with standard error and 95% confidence intervals were
179 calculated along with diversity indices (Species Richness, Shannon Diversity, and Simpson
180 Diversity) for all herpetofauna, as well as amphibians, and reptiles independently.

181

182 *IUCN predictive species lists.* To compare our observed local diversity based on the field
183 surveys, with the expected regional diversity, we extracted spatial distribution data for all
184 amphibian and reptile species with ranges overlapping our study site using the IUCN Red List
185 spatial database (IUCN, 2025). This predictive distribution list serves as a theoretical baseline
186 of expected species richness in the region, enabling the evaluation of our field surveys
187 performance, identifying potential sampling gaps, and contextualising our findings within the
188 broader regional herpetofaunal assemblage.

189

190 To assess the similarity between our field survey results and the IUCN predicted species
191 distribution, we calculated the Jaccard similarity index (J). This metric is based on
192 presence/absence and measures the proportion of shared species compared to the total species
193 pool (Magurran, 2004). The Jaccard index was calculated as $J = C/(A+B-C)$, where A is the
194 number of species observed through our surveys, B is the number of species from the IUCN
195 predicted distribution for the study area, and C is the number of species common to both lists.
196 The index ranges from 0, (no overlap) to 1 (complete similarity) and was used to quantify the
197 taxonomic overlap between our survey results and the predicted regional species assemblage
198 based on IUCN distribution maps. For clarity, we report both the proportion of IUCN-predicted

199 species we recorded in total (A/B) and the proportion of species our survey shared with IUCN
200 predictions (C/B), which is the numerator of the Jaccard index.

201

202 *Comparison to benchmark surveys.* To ground our findings within the broader context of
203 Liberian herpetofaunal research, we compared our survey results from the Dugbe study area
204 with two benchmark surveys conducted in southeastern Liberia: KBPPA by Rödel and Glos
205 (2019) and the Grebo National Forest survey by Hillers and Rödel (2007). These study sites
206 offer appropriate benchmarks for comparison based on their proximity (both located within
207 100 km of our Dugbe) and similar habitat types. For each survey, we compared species richness
208 for all herpetofauna, as well as amphibians and reptiles separately. As the focus and survey
209 methods of Rödel and Glos (2019) and Hillers and Rödel (2007) studies differed from ours, the
210 comparisons were restricted to species richness, providing an evaluation of our sampling effort
211 and general insight into the regional herpetofaunal assemblage.

212

RESULTS

213 During the Dugbe survey, we documented 1145 individual amphibians and reptiles
214 representing 71 distinct taxa (**Error! Reference source not found.**, Supplementary material
215 1). The assemblage comprised 38 amphibian species (982 individual observations) and 33
216 reptile species (156 individual observations). A comprehensive species inventory for the Dugbe
217 study area is presented in Table 1 and Table 2, detailing amphibians and reptiles respectively,
218 including common names, IUCN conservation status, observation frequencies, and West
219 African endemism status. A selection of photos of recorded species is presented in Fig. 3–4.

220

221 Active searching proved the most productive survey method, contributing 985 observations
222 (86%) and 34 unique species (47.8% of total richness). Trapping methods contributed 122

223 observations (10.6%) and four unique species (5.6%). Incidental records accounted for 38
224 observations (3.3%), including nine unique species (12.6%). Notably, 39 species (54.9%)
225 documented in the Dugbe study area are West African endemics. The regional distinctiveness
226 is particularly pronounced among amphibians (28 species, 73.6% are endemic) compared to
227 reptiles (11 species, 33.3%).

228



229

230 **Fig 3** .A) Seraphin's Caecilian (*Geotrypetes seraphini*); B) Mottled Squeaker (*Arthroleptis*
 231 *poecilonotus* complex); C) Western Night Frog (*Astylosternus occidentalis*); D) Large-eared
 232 Treefrog (*Leptopelis macrotis*); E) Western Treefrog (*Leptopelis occidentalis*); F) Ghostly Tree
 233 Frog (*Leptopelis spiritusnoctis*); G) Northern Flat-backed Toad Toad (*Sclerophrys maculata*);

234 H) Togo Toad (*Sclerophrys togoensis*); I) Allen's Giant Frog (*Conraua alleni*); J) African Tiger
235 Frog (*Hoplobatrachus occipitalis*); K) Brown Fishing Frog (*Aubria subsigillata*); L) Lesser
236 White-lipped Frog (*Amnirana parva*); M) Schiøtz's Grass Frog (*Ptychadena arnei*); N)
237 Bibron's Grass Frog (*Ptychadena cf. bibroni*); O) Snouted Grass Frog (*Ptychadena*
238 *longirostris*); P) Western Dwarf Grass Frog (*Ptychadena pumilo*); Q) Striped Spiny Reed Frog
239 (*Afrixalus dorsalis*); R) Nigerian Spiny Reed Frog (*Afrixalus nigeriensis*); S) Large Green Reed
240 Frog (*Hyperolius chlorosteus*); T) Uniform Reed Frog (*Hyperolius concolor*); U) Dark-bellied
241 Reed Frog (*Hyperolius fusciventralis*); V) Dotted Reed Frog (*Hyperolius guttulatus*); W) Painted
242 Reed Frog (*Hyperolius picturatus*); X) Soror Reed Frog (*Hyperolius soror*); Y) Boulenger's
243 Wot-wot (*Hylambates boulengeri*); Z) Allen's Puddle Frog (*Phrynobatrachus alleni*); AA)
244 Brother's Puddle Frog (*Phrynobatrachus fraterculus*); AB) Guinea Puddle Frog
245 (*Phrynobatrachus guineensis*); AC) Savanna Puddle Frog (*Phrynobatrachus latifrons*); AD)
246 Liberia Puddle Frog (*Phrynobatrachus liberiensis*); AE) Leaf-loving Puddle Frog
247 (*Phrynobatrachus phyllophilus*); AF) Ridged Puddle Frog (*Phrynobatrachus plicatus*). All
248 photographs by Marius Burger and Ryan van Huyssteen.

249



250

251 **Fig. 4.** A) Tokba Puddle Frog (*Phrynobatrachus tokba*); B) Villiers' Puddle Frog
 252 (*Phrynobatrachus villiersi*); C) Allen's Puddle Frog (*Phrynobatrachus allenii* complex); D)
 253 Common Toothed Frog (*Odontobatrachus natator*); E) Tropical Clawed Frog (*Xenopus*
 254 *tropicalis*); F) Western Foam-nest Frog (*Chiromantis rufescens*); G) Banded Leaf-toed Gecko

255 (Hemidactylus fasciatus); H) Guinea Leaf-toed Gecko (Hemidactylus muriceus); I) Keeled
256 Water Skink (Cophoscincopus durus); J) Fire Skink (Mochlus fernandi); K) Senegal Skink
257 (Trachylepis affinis); L) Speckle-lipped Skink (Trachylepis maculilabris); M) Tropical Skink
258 (Trachylepis paucisquamis); N) West African Agama (Agama africana); O) Water Monitor
259 (Varanus niloticus); P) Guinea Blind Snake (Afrotyphlops liberensis); Q) Calabar Ground
260 Python (Calabaria reinhardtii); R) Green Bush Viper (Atheris chlorechis); S) Rhinoceros Viper
261 (Bitis nasicornis); T) Spotted Night Adder (Causus maculatus); U) Western Forest Centipede-
262 eater (Aparallactus modestus); V) Red-Black Striped Snake (Bothrophthalmus lineatus); W)
263 Kling's File Snake (Gonionotophis klingi); X) Equatorial File Snake (Mehelya poensis); Y)
264 Ugandan House Snake (Hormonotus modestus); Z) Black Forest Cobra (Naja guineensis); AA)
265 Black Tree Cobra (Pseudohaje nigra); AB) African Brown Water Snake (Afronatrix
266 anoscopus); AC) Short-headed Tree Snake (Dipsadoboabrevirostris); AD) Black-lined Green
267 Snake (Hapsidophrys lineatus); AE) Variable Marsh Snake (Natriciteres variegata); AF) Forest
268 Vine Snake (Thelotornis kirtlandii). All photographs by Marius Burger and Ryan van
269 Huyssteen.


270

271 *Sampling curves*

272 The r/e curves (Fig. ; Table 1) reveal incomplete sampling of overall herpetofauna diversity
273 across the Dugbe study area. For combined herpetofauna, the model estimated total species
274 richness at 116.09 (upper confidence limit: 160.53) compared to the 71 observed. Shannon
275 diversity was 28.34 and predicted to be 29.78, while Simpson diversity was 15.30 and
276 estimated to be 15.49. For amphibians, sampling completeness was high. The observed species
277 richness of 38 and the estimated richness match (upper confidence limit: 40.61). This indicates
278 that the amphibian diversity was adequately sampled. Shannon diversity was 20.26 and closely
279 matched the estimated 20.66 and Simpson diversity was 11.91 and estimated to be 12.05 (Fig.

280 ; Table 13). In contrast, the species richness for reptiles of 33 was far less than the predicted
281 species richness of 122.67 (confidence intervals: 33–256.05), indicating a significant sampling
282 deficiency. The Shannon diversity of 12.69 was predicted to be 16.97, while Simpson diversity
283 of 6.57 was estimated to be 6.81. There were a high proportion of reptile species being recorded
284 only once, i.e. singletons (19 species; 57.57% of reptile records) (Fig. ; Table 1).

285



286

287 **Fig. 5.** The r/e sampling curves for all herpetofauna, amphibians and reptiles of the Dugbe
288 region, solid line shows rarified survey effort and stippled line shows extrapolated results.

289

290 *IUCN species list comparison*

291 When comparing the results of our Dugbe herpetofauna survey with the IUCN predictive
292 distribution maps, we found notable discrepancies. Our survey documented 71 species, 63 of
293 these were predicted by the IUCN predictive distribution maps. This represents 51.2% of the
294 123 species predicted by IUCN range maps for the region. Four species (*Afrixalus nigeriensis*
295 Schiøtz, 1963; *Hyperolius soror* (Chabanaud, 1921); *Leptopelis occidentalis* Schiøtz, 1967;
296 and *Phrynobatrachus gutturosus* (Chabanaud, 1921)) are not predicted to occur in the study
297 area by the IUCN, however they are predicted to occur nearby based on Channing and Rödel
298 (2019). Two reptile species (*Kinixys erosa* (Schweigger, 1812); and *Naja guineensis* Broadley
299 et al., 2018) are not mapped by the IUCN. Considering our survey results in relation to the
300 IUCN predicted list, we found a a moderate overall Jaccard similarity (48.09%) (Table 4).

301

302 Taxonomic groups showed differential patterns of congruence. For amphibians, 32 of the 38
303 observed species were predicted by IUCN (66.7% of 48 expected species). Amphibians showed
304 a relatively high Jaccard Similarity Index (59.26%). In contrast, 31 of the 33 observed reptile
305 species were predicted by the IUCN, representing just 41.3% of the 75 reptile species predicted
306 by IUCN maps, with a correspondingly lower Jaccard similarity (40.26%) (Table 4).

307

308 *Comparison to benchmark*

309 Our survey of the Dugbe region documented higher total herpetofauna species richness (71
310 species) compared to the KBPPA survey (48 species) and the Grebo National Forest survey (34
311 species). It is important to note that the two benchmark surveys employed different methods to
312 our survey, lasted less days, and were both focused on amphibians rather than all herpetofauna.
313 For amphibians, our survey recorded 38 species, compared to 34 species in the KBPPA survey
314 and 29 species in the Grebo National Forest survey. Reptiles were recorded opportunistically

315 by the benchmark studies, and therefore a pronounced difference can be observed in reptile
316 diversity, where our survey documented 33 species, while the KBPPA and Grebo National
317 Forest surveys recorded 14 and 5 species, respectively (Table 5).

318

319 Of the 48 species recorded in the KBPPA, 37 of these were also identified during our Dugbe
320 survey, while there was overlap between 27 species found in Grebo. Together, the KBPPA and
321 Grebo surveys documented 14 taxa which were not detected during our survey, including
322 *Trionyx triunguis* (Forskål, 1775) and *Osteolaemus tetraspis* (Cope, 1861).

323

324 The comparisons between the benchmark studies are restricted to species richness as the focus
325 and methods were different to ours. A comparative overview with the benchmark sites, KBPPA
326 (Rödel & Glos, 2019) and Grebo National Forest (Hillers & Rödel, 2007), is available as a
327 supplement which additionally serves as an evidence-based species list for the greater area
328 (Supplementary material 2).

329

330 **DISCUSSION**

331 Our survey provides 1145 herpetofauna observations comprising 71 species (38 amphibians
332 and 33 reptiles) from a previously understudied region of Liberia. For all herpetofauna, 39
333 (54.9%) of the species recorded are endemic to West Africa. For frogs this endemism was
334 73.6% and for reptiles 33.33%. One of the most notable species in terms of endemism is
335 *Odontobatrachus natator* (Boulenger, 1905) which belongs to West Africa's only endemic
336 vertebrate family, the Odontobatrachidae (Barej et al., 2014). This high degree of endemism
337 highlights the herpetological value of the study area, and the Upper Guinean forests in general
338 (Myers et al., 2000; Ernst et al., 2025). Regional comparisons with nearby surveys (KBPPA

339 and Grebo National Forest) contextualise our findings within the broader Upper Guinean forest
340 landscape.

341
342 Four species of conservation concern were recorded: *Leptopelis macrotis* Schiøtz, 1967 (Near
343 Threatened (NT)), *L. occidentalis* Schiøtz, 1967 (NT), *Bitis nasicornis* (Shaw & Nodder, 1792)
344 (Vulnerable (VU)), and *Kinixys erosa* (Data Deficient (DD)). The *Leptopelis* species rely on
345 primary forests and are threatened by human activities such as artisanal mining. *Bitis nasicornis*
346 is affected by habitat loss and hunting pressure (Penner et al., 2021). It is important to note that
347 the current conservation assessment for *K. erosa* is outdated, and preliminary assessments
348 suggest it could be categorised as Endangered (EN) due to the impacts of deforestation and
349 hunting (Luiselli & Diagne, 2014).

350
351 While sampling completeness metrics indicate comprehensive sampling for amphibians, reptile
352 sampling was likely incomplete, with r/e curves skewed by a high proportion of singletons
353 (60%). This difference in sampling success between amphibians and reptiles can be attributed
354 to several factors: our surveys coincided with the wet season when amphibian breeding activity
355 is at its peak. Amphibians are highly vocal during the breeding season and tend to aggregate at
356 breeding sites, increasing encounter probability (Rödel & Ernst, 2004; Heyer et al., 2014). In
357 terms of reptiles, while certain genera such as *Agama* and *Trachylepis* include relatively
358 conspicuous species, reptiles are generally secretive, cryptic, and difficult to detect, particularly
359 snakes (Durso & Seigel, 2015; Jordaan et al., 2021). For example, many reptile species exhibit
360 secretive lifestyles (i.e., nocturnal, fossorial), shy behaviour, cryptic coloration, and are often
361 sensitive to disturbances, all of which reduce detection probability during surveys.
362 Additionally, high rainfall has been found to negatively correlate with reptile detectability in

363 several studies (e.g., Eskew & Todd, 2017; Falaschi, 2021). The persistent heavy rainfall during
364 our surveys likely contributed to the under sampling of reptiles in our study.

365

366 Based on morphology, species level identifications were problematic for some amphibian taxa,
367 namely those in the *Arthroleptis poecilonotus* complex Peters, 1863, *Phrynobatrachus allenii*
368 complex Parker, 1936, and several *Ptychadena* specimens remained unidentified. New taxa
369 have been identified in these groups and are currently in the process of being described in the
370 study area (Rödel et al., in prep.). Currently, without a clear taxonomic framework, undescribed
371 species cannot be accurately classified under threat categories, impacting conservation efforts.
372 There is a need for additional taxonomic studies on West African herpetofauna (Luiselli et al.,
373 2019), particularly given the Upper Guinean forests' importance for amphibian diversification
374 and the increasing anthropogenic pressures on these ecosystems (Penner et al., 2011, Rödel et
375 al., 2021).

376

377 Our field survey documented fewer species than predicted by IUCN range maps, particularly
378 for reptiles (33 observed vs. 75 predicted), compared to amphibians (39 observed vs. 48
379 predicted). Considering the limited survey duration, it is likely that many of the additional
380 species predicted by the IUCN are present in the study area and were undetected. However,
381 while IUCN interpreted distributions provide valuable tools for conservation planners and
382 researchers, their accuracy varies significantly by region. In well-studied areas like South
383 Africa (Tolley et al. 2016), species ranges are well understood, resulting in highly reliable
384 distribution data (Tolley et al., 2023). However, knowledge of herpetofauna distributions in
385 Liberia remains largely in development. Therefore, it is possible that some of the IUCN
386 distribution maps are inaccurate. Baseline surveys, such as ours, are required to inform and

387 refine understanding of species distributions to improve the accuracy of future distribution
388 maps.

389
390 Active searching yielded the most records (>80%) and approximately half of the unique species
391 documented (47.8%). Passive trapping and incidental observations, accounted for only
392 approximately 10% and 3% of records respectively, however, they contributed significantly to
393 overall species richness by adding 5% and 11.5% of unique species. This highlights the
394 importance of diversifying survey techniques to maximise detection during baseline surveys
395 (McDiarmid et al., 2012; Ribeiro-Júnior et al., 2008).

396
397 The primary limitations of our survey were the climatic conditions and the challenges presented
398 by taxonomic resolution. We recommend that additional future surveys in the region take place
399 at the end of or at the start of the rain season, as that is when amphibian and reptile activity is
400 at its greatest. To overcome taxonomic uncertainties, researchers should maintain
401 collaborations with taxonomists specialising in West African herpetofauna, as this will improve
402 future species inventories and our understanding of the region's herpetofaunal diversity.

403

404 **ACKNOWLEDGEMENTS**

405 We thank Llyod Kirtley, the Dugbe camp manager. For their contributions in the field thanks
406 to: Marco Alexandre, Jean-Michel Blake, Phoebe Mottram, Edmond Miabangana, Lukas
407 Niemand, Ben Orban, Gina Walsh, and Graeme Wolfaard. For assistance with species
408 identification, we thank Kaitlin Allen, Uwe Fritz, Flora Ihlow, Olivier Pauwels, Johannes
409 Penner, Alexander Rebelo, Colin Tilbury, Krystal Tolley, and Jean-Francois Trappe. Permits to
410 conduct the study were granted by Forestry Development Authority (FDA), permit number

411 FDA/DMDA11/703/202 and FDA/DMDA/11/704/202. We thank Tường Định and an
412 anonymous reviewer for the constructive comments and suggestions which greatly enhanced
413 the manuscript.

414

REFERENCES

416 Allen, K.E., Greenbaum, E., Welton, L.J., Bauer, A.M. (2019): High levels of hidden
417 phylogenetic structure within Central and West African *Trachylepis* skinks. *Salamandra*
418 **55**(4): 231–241.

419 Angel, F. (1943): Description d'un nouvel Anoure ovovivipare de la Haute Guinée français.
420 *Bull. Mus. Paris* **15**: 167–169.

421 Barbour, T., Loveridge, A. (1927): Some undescribed frogs and a new gecko from Liberia.
422 *Proc. New England Zool. Club* **10**: 13–18.

423 Barej, M.F., Schmitz, A., Günther, R., Loader, S.P., Mahlow, K., Rödel, M.-O. (2014): The
424 first endemic West African vertebrate family—a new anuran family highlighting the
425 uniqueness of the Upper Guinean biodiversity hotspot. *Front. Zool.* **11**: 1–11.

426 Blackburn, D.C., Kosuch, J., Schmitz, A., Burger, M., Wagner, P., Gonwouo, N.L., Hillers,
427 A., Rödel, M.-O. (2008): A new species of *Cardioglossa* (Anura: Arthroleptidae) from the
428 Upper Guinean forests of West Africa. *Copeia* **2008**(3): 603–612.

429 Boulenger, G.A. (1905): Descriptions of new West-African frogs of the genera *Petropedetes*
430 and *Bulua*. *Ann. Mag. Nat. Hist.* **15**: 281–283.

431 Bourgeron, P.S., Humphries, H.C., Jensen, M.E. (2001): Representativeness assessments. In:
432 A guidebook for integrated ecological assessment, pp. 292–306. Jensen, M.E., Bourgeron,
433 P.S., Eds, New York, Springer-Verlag.

434 Chabanaud, P. (1921): Contributions a l'étude de la faune herpetologique de l'Afrique
435 occidentale. II. Deuxième note. *Bull. Comité Études Hist. Sci. Afr. Occid. Fr.* **1921**: 445–472.

436 Channing, A., Rödel, M.-O. (2019): Field guide to the frogs & other amphibians of Africa.
437 Cape Town, Penguin Random House South Africa.

438 Chippaux, J.P. (2006): Les serpents d'Afrique occidentale et centrale. Paris, IRD éditions.

439 Cope, E.D. (1861): List of the Recent species of emydosaurian reptiles in the Museum of the
440 Academy of Natural Sciences. *Proc. Acad. Nat. Sci. Phila.* **12**: 549–550.

441 Cushing, W.M., Cotillon, S.E., Hutchinson, J.A., Pengra, B., Alfari, I., Botoni, E., Soulé, A.,
442 Herrmann, S.M., Tappan, G.G. (2016): Landscapes of West Africa: A window on a changing
443 world. *Garretson, United States Geological Survey.*

444 de Sousa, C., Fatoyinbo, L., Honzák, M., Wright, T.M., Murillo Sandoval, P.J., Whapoe, Z.E.,
445 Yonmah, J., Olatunji, E.T., Garteh, J., Stovall, A., Neigh, C.S., Portela, R., Gaddis, K.D.,
446 Larsen, T., Juhn, D. (2023): Two decades of land cover change and forest fragmentation in
447 Liberia: Consequences for the contribution of nature to people. *Conserv. Sci. Pract.* **5**(6):
448 e12933.

449 Deikumah, J.P., McAlpine, C.A., Maron, M. (2014): Biogeographical and taxonomic biases
450 in tropical forest fragmentation research. *Conserv. Biol.* **28**(6): 1522–1531.

451 Durso, A.M., Seigel, R.A. (2015): A snake in the hand is worth 10,000 in the bush. *J.*
452 *Herpetol.* **49**(4): 503–506.

453 Ernst, M., Rödel, M.-O., Blom, M.P. (2025): Towards a comprehensive view on evolutionary
454 refugia in West African rainforests. *Front. Biogeogr.* **18**: e139537.

455 Eskew, E.A., Todd, B.D. (2017): Too cold, too wet, too bright, or just right? Environmental
456 predictors of snake movement and activity. *Copeia* **105**(3): 584–591.

457 Falaschi, M. (2021): Phenology and temperature are the main drivers shaping the detection
458 probability of the common wall lizard. *Amphibia-Reptilia* **42**(3): 297–303.

459 Forskål, P. (1775): *Descriptiones animalium, avium, amphibiorum, piscium, insectorum,*
460 *vermium: quae in itinere orientali observavit.* Copenhagen, ex officina Mölleri.

461 Frost, D.R. (2025): *Amphibian Species of the World: an Online Reference.* Version 6.2.
462 American Museum of Natural History, New York.

463 <https://amphibiansoftheworld.amnh.org/index.php> doi.org/10.5531/db.vz.0001 [Accessed 15
464 October 2025].

465 Guibé, J., Lamotte, M. (1958): Une espèce nouvelle de batracien du Mont Nimba (Guinée
466 Française) appartenant au genre *Phrynobatrachus*: *Ph. maculiventris* n. sp. Bull. Mus. Natl.
467 Hist. Nat., 2e Sér. **30**: 255–257.

468 Hepner, S., Agonvonon, G.A., Ehbrecht, M., Iheaturu, C., Azihou, A.F., Ifejika Speranza, C.
469 (2025): Degradation and Fragmentation Effects on Structural Complexity in West-African
470 Forest Patches. *Biotropica* **57**(2): e70012.

471 Hillers, A., Rödel, M.-O. (2007): Rapid survey of amphibians and reptiles of North Lorma,
472 Gola and Grebo National Forests. In: A rapid biological assessment of North Lorma, Gola
473 and Grebo National Forests, Liberia, pp. 29–33. Hoke, P., Demey, R., Peal, A., Eds,
474 Arlington, Conservation International, Center for Applied Biodiversity Science.

475 Hoke, P., Demey, R., Peal, A. (2007): A rapid biological assessment of north Lorma, Gola and
476 Grebo national forests, Liberia. Arlington, Conservation International, Center for Applied
477 Biodiversity Science.

478 Ineich, I. (2003): Contribution à la connaissance de la biodiversité des régions afro-
479 montagnardes: les Reptiles du Mont Nimba. In: Le peuplement animal du Mont Nimba
480 (Guinée, Côte d'Ivoire, Liberia). Mém. Mus. nat. Hist. Nat. **190**: 597–637. Lamotte, M., Roy,
481 R., Eds, Paris, Muséum national d'Histoire naturelle.

482 IUCN (2024): The IUCN Red List of Threatened Species. Version 2024-2.

483 <https://www.iucnredlist.org> [Accessed 05 February 2025].

484 IUCN SSC Amphibian Specialist Group (2013): *Phrynobatrachus tokba*. The IUCN Red List
485 of Threatened Species 2013: e.T58145A18396312.

486 <https://dx.doi.org/10.2305/IUCN.UK.2013-2.RLTS.T58145A18396312.en>

487 IUCN SSC Amphibian Specialist Group (2014a): *Leptopelis macrotis*. The IUCN Red List of
488 Threatened Species 2014: e.T56263A16925323. <https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T56263A16925323.en>

489

490 IUCN SSC Amphibian Specialist Group (2014b): *Leptopelis occidentalis*. The IUCN Red
491 List of Threatened Species 2014: e.T56271A16925453.
492 <https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T56271A16925453.en>

493 Jensen, M.E., Bourgeron, P.S., Eds (2012): A guidebook for integrated ecological
494 assessments. New York, Springer Science & Business Media.

495 Jordaan, A., Heideman, N.J., Buschke, F.T. (2021): Topography-derived variables provide
496 insight into habitat occupancy of a cryptic snake, *Bitis atropos*. *Austral Ecol.* **46**(8): 1287–
497 1297.

498 Lamotte, M. (1998): Panorama d'une fauna oust fricaine d'amphibiens: le peuplement du mont
499 Nimba. *Bull. Soc. Herp. Fr.* **87–88**: 5–23.

500 Lewin, A., Feldman, A., Bauer, A.M., Belmaker, J., Broadley, D.G., Chirio, L., Itescu, Y.,
501 LeBreton, M., Maza, E., Meirte, D., Meiri, S. (2016): Patterns of species richness, endemism
502 and environmental gradients of African reptiles. *J. Biogeogr.* **43**(12): 2380–2390.

503 Loveridge, A. (1941): Report on the Smithsonian-Firestone expedition's collection of reptiles
504 and amphibians from Liberia. *Proc. U.S. Natl. Mus.* **91**: 113–140.

505 Luiselli, L., Diagne, T. (2014): *Kinixys erosa* (Schweigger 1812)—forest hinge-back tortoise,
506 serrated hinge-back tortoise, serrated hinged tortoise. *Chelonian Res. Monogr.* **5**: 1–13.

507 Luiselli, L., Dendi, D., Eniang, E.A., Fakae, B.B., Akani, G.C., Fa, J.E. (2019): State of
508 knowledge of research in the Guinean forests of West Africa region. *Acta Oecol.* **94**: 3–11.

509 Magurran, A.E. (2004): Measuring biological diversity. Oxford, John Wiley & Sons.

510 Mendes, D.M., de Freitas Leão, R., Toledo, L.F. (2015): Drift fences in traps: theoretical
511 evidence of effectiveness of the two most common arrays applied to terrestrial tetrapods. *Nat.*
512 *Conserv.* **13**(1): 60–66.

513 Mottram, P., van Huyssteen, R., Okoue, E.A., Petford, M., Gaugris, J. (2025): Mammal
514 diversity in southern Gabon: Grid-based vs. semi-random camera trapping. *Afr. J. Wildl. Res.*
515 **55**(1): 258–265.

516 Myers, N., Mittermeier, R.A., Mittermeier, C.G., Da Fonseca, G.A., Kent, J. (2000):
517 Biodiversity hotspots for conservation priorities. *Nature* **403**(6772): 853–858.

518 Nneji, L.M., Azevedo, J.A., Oyebanji, O.O., Ma, L., Elsen, P.R., Oladipo, S.O., Salako, G.,
519 Puschendorf, R., Pringle, R.M. (2023): Patterns of species richness and turnover in endemic
520 amphibians of the Guineo-Congolian rain forest. *Divers. Distrib.* **29**(8): 1035–1051.

521 Penner, J., Augustin, M., Rödel, M.-O. (2019): Modelling the spatial baseline for amphibian
522 conservation in West Africa. *Acta Oecol.* **94**: 31–40.

523 Penner, J., Rödel, M.-O., Luiselli, L., Trape, J.-F., Spawls, S., Malonza, P.K., Beraduccii, J.,
524 Chippaux, J.-P., LeBreton, M., Kusamba, C., Gonwouo, N.L. (2021): *Bitis nasicornis*. The
525 IUCN Red List of Threatened Species 2021: e.T13300910A13300919.
526 <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T13300910A13300919.en>

527 Penner, J., Wegmann, M., Hillers, A., Schmidt, M., Rödel, M.-O. (2011): A hotspot revisited—
528 a biogeographical analysis of West African amphibians. *Divers. Distrib.* **17**(6): 1077–1088.

529 Rödel, M.-O., Doumbia, J., Johnson, A.T., Hillers, A. (2009): A new small *Arthroleptis*
530 (Amphibia: Anura: Arthroleptidae) from the Liberian part of Mount Nimba, West Africa.
531 *Zootaxa* **2302**(1): 19–30.

532 Rödel, M.-O., Ernst, R. (2002): A new reproductive mode for the genus *Phrynobatrachus*:
533 *Phrynobatrachus alticola* has nonfeeding, nonhatching tadpoles. *J. Herpetol.* **36**(1): 121–125.

534 Rödel, M.-O., Glos, J. (2019): Herpetological surveys in two proposed protected areas in
535 Liberia, West Africa. *Zoosyst. Evol.* **95**(1): 15–35.

536 Rödel, M.-O., Adum, G.B., Aruna, E., Assemian, N.E., Barej, M.F., Bell, R.C., Burger, M.,
537 Demare, G., Doherty-Bone, T., Doumbia, J., Ernst, R., Gonwouo, N.L., Hillers, A.,
538 Hirschfeld, M., Jongsma, G.F.M., Kouamé, N.G., Kpan, T.F., Mohneke, M., Nago, S.G.A.,
539 Ofori-Boateng, C., Onadeko, A., Pauwels, O.S.G., Sandberger-Loua, L., Segniagbeto, G.H.,
540 Tchassem Fokoua, A.M., Tobi, E., Tohé, B., Zimkus, B.M., Penner, J. (2021): Diversity,
541 threats and conservation of western and central African amphibians (Senegal, The Gambia,
542 Guinea Bissau, Mali, Guinea, Sierra Leone, Liberia, Ivory Coast, Burkina Faso, Ghana, Togo,
543 Benin, Nigeria, Niger, Cameroon, Gabon, São Tome & Príncipe, Equatorial Guinea, Central
544 African Republic, Chad, Republic of the Congo, Democratic Republic of the Congo, northern
545 Angola). In: Status and threats of Afrotropical Amphibians – Sub-Saharan Africa,
546 Madagascar, Western Indian Ocean Islands. *Amphibian Biology*, Volume 11, Part 7, pp. 11–
547 101. Heatwole, H., Rödel, M.-O., Eds, Frankfurt am Main, Edition Chimaira.

548 Sandberger, L., Hillers, A., Doumbia, J., Loua, N.S., Brede, C., Rödel, M.-O. (2010):
549 Rediscovery of the Liberian Nimba toad, *Nimbaphrynoidea liberiensis* (Xavier, 1978)
550 (Amphibia: Anura: Bufonidae), and reassessment of its taxonomic status. *Zootaxa* **2355**: 56–
551 68.

552 Schiøtz, A. (1963): The amphibians of Nigeria. *Vidensk. Medd. Dansk Naturhist. Foren.*
553 Kjøbenh. **125**: 1–92.

554 Schiøtz, A. (1967): The treefrogs (Rhacophoridae) of West Africa. *Spolia Zool. Mus. Haun.*
555 **25**: 1–346.

556 Schiøtz, A. (1999): The treefrogs of Africa. Frankfurt am Main, Edition Chimaira.

557 Schweigger, A.F. (1812): *Prodromus Monographia Cheloniorum auctore Schweigger.*
558 Königsberg. *Arch. Naturwiss. Mathem.* **1**: 271–368, 406–458.

559 Tolley, K., Conradie, W., Pietersen, D., Weeber, J., Burger, M., Alexander, G.J., Eds (2023):
560 Conservation status of the reptiles of South Africa, Eswatini and Lesotho. Pretoria, South
561 African National Biodiversity Institute.

562 Tolley, K.A., Alexander, G.J., Branch, W.R., Bowles, P., Maritz, B. (2016): Conservation
563 status and threats for African reptiles. *Biol. Conserv.* **204**: 63–71.

564 Tortoise & Freshwater Turtle Specialist Group (1996): *Kinixys erosa*. The IUCN Red List of
565 Threatened Species 1996: e.T11002A3238083.

566 <https://dx.doi.org/10.2305/IUCN.UK.1996.RLTS.T11002A3238083.en>

567 Trape, J.F., Mané, Y. (2006): Le genre *Dasypeltis* Wagler (Serpentes: Colubridae) en Afrique
568 de l'Ouest: description de trois espèces et d'une sous-espèce nouvelles. *Bull. Soc. Herpetol.*
569 Fr. **119**: 27–56.

570 Trape, J.F., Trape, S., Chirio, L. (2012): Lézards, crocodiles et tortues d'Afrique occidentale
571 et du Sahara. Paris, IRD éditions.

572 Uetz, P. (2025): The Reptile Database. <http://www.reptile-database.org> [Accessed 23 May
573 2025].

574 Wüster, W., Chirio, L., Trape, J.F., Ineich, I., Jackson, K., Greenbaum, E., Barron, C.,
575 Kusamba, C., Nagy, Z.T., Storey, R., Hall, C. (2018): Integration of nuclear and
576 mitochondrial gene sequences and morphology reveals unexpected diversity in the forest
577 cobra (*Naja melanoleuca*) species complex in Central and West Africa (Serpentes: Elapidae).
578 *Zootaxa* **4455**(1): 68–98.

579 Xavier, F. (1978): Une espèce nouvelle de *Nectophrynoïdes* (Anoure, Bufonidae) des Monts
580 Nimba, *N. liberiensis*. 1.- Description de l'espèce. *Bull. Soc. Zool. France* **103**(4): 431–441.

581 **Table 1.** Amphibian species recorded during surveys in the Dugbe region, Liberia. IUCN
 582 status, number of observations, and West African endemism noted.
 583

TAXON	COMMON NAME	IUCN	OBSERVATIONS	WEST AFRICAN ENDEMIC
Class: Amphibia				
Order: Anura				
Family: Arthroleptidae				
<i>Arthroleptis poecilonotus</i> complex	Mottled Squeaker	—	90	—
<i>Astylosternus occidentalis</i>	Western Night Frog	LC	8	Yes
<i>Leptopelis macrotis</i>	Large-eared Tree Frog	NT	2	Yes
<i>Leptopelis occidentalis</i>	Western Tree Frog	NT	2	Yes
<i>Leptopelis spiritusnoctis</i>	Ghostly Tree Frog	LC	4	Yes
Family: Bufonidae				
<i>Sclerophrys maculata</i>	Northern Flat-backed Toad	LC	50	No
<i>Sclerophrys togoensis</i>	Togo Toad	LC	4	Yes
Family: Conrauidae				
<i>Conraua alleni</i>	Allen's Giant Frog	LC	4	Yes
Family: Dicroidiidae				
<i>Hoplobatrachus occipitalis</i>	African Tiger Frog	LC	10	No
Family: Hyperoliidae				
<i>Afrixalus dorsalis</i>	Striped Spiny Reed Frog	LC	20	No
<i>Afrixalus nigeriensis</i>	Nigerian Spiny Reed Frog	LC	24	Yes
<i>Hylambates boulengeri</i>	Boulenger's Wot-wot	LC	20	Yes
<i>Hyperolius chlorostictus</i>	Large Green Reed Frog	LC	24	Yes
<i>Hyperolius concolor</i>	Uniform Reed Frog	LC	21	Yes
<i>Hyperolius fusciventris</i>	Dark-bellied Reed Frog	LC	45	Yes
<i>Hyperolius guttulatus</i>	Dotted Reed Frog	LC	12	No
<i>Hyperolius picturatus</i>	Painted Reed Frog	LC	9	Yes
<i>Hyperolius soror</i>	Soror Reed Frog	LC	28	Yes
Family: Odontobatrachidae				
<i>Odontobatrachus natator</i>	Common Toothed Frog	LC	15	Yes
Family: Phrynobatrachidae				

<i>Phrynobatrachus allenii</i> complex	Allen's Puddle Frog	LC	45	Yes
<i>Phrynobatrachus fraterculus</i>	Brother's Puddle Frog	LC	5	Yes
<i>Phrynobatrachus guineensis</i>	Guinea Puddle Frog	LC	10	Yes
<i>Phrynobatrachus gutturosus</i>	Guttural Puddle Frog	LC	7	Yes
<i>Phrynobatrachus latifrons</i>	Savanna Puddle Frog	LC	14	Yes
<i>Phrynobatrachus liberiensis</i>	Liberian Puddle Frog	LC	43	Yes
<i>Phrynobatrachus phyllophilus</i>	Leaf-loving Puddle Frog	LC	9	Yes
<i>Phrynobatrachus plicatus</i>	Ridged Puddle Frog	LC	7	Yes
<i>Phrynobatrachus tokba</i>	Tokba Puddle Frog	LC	221	Yes
<i>Phrynobatrachus villiersi</i>	Villiers' Puddle Frog	LC	9	Yes
Family: Pipidae				
<i>Xenopus tropicalis</i>	Tropical Clawed Frog	LC	88	Yes
Family: Ptychadenidae				
<i>Ptychadena arnei</i>	Schiøtz's Grass Frog	DD	5	Yes
<i>Ptychadena cf. bibroni</i>	Bibron's Grass Frog	LC	44	No
<i>Ptychadena longirostris</i>	Snouted Grass Frog	LC	23	Yes
<i>Ptychadena pumilo</i>	Western Dwarf Grass Frog	LC	13	No
Family: Pyxicephalidae				
<i>Aubria subsigillata</i>	Brown Fishing Frog	LC	4	No
Family: Ranidae				
<i>Amniranana parva</i>	Lesser White-lipped Frog	—	7	Yes
Family: Rhacophoridae				
<i>Chiromantis rufescens</i>	Western Foam-nest Frog	LC	24	No
Order: Gymnophiona				
Family: Dermophiidae				
<i>Geotrypetes seraphini</i>	Seraphin's Caecilian	LC	12	No

585 **Table 2.** Reptile species recorded during surveys in the Dugbe region, Liberia. IUCN status,
 586 number of observations, and west African Endemism noted.

TAXON	COMMON NAME	IUCN	OBSERVATIONS	WEST AFRICAN ENDEMIC
Class: Reptilia				
Order: Testudines				
Family: Testudinidae				
<i>Kinixys erosa</i>	Forest Hinge-back Tortoise	DD	1	No
Order: Squamata				
Family: Agamidae				
<i>Agama africana</i>	West African Agama	LC	50	Yes
Family: Atractaspididae				
<i>Aparallactus modestus</i>	Western Forest Centipede-eater	LC	1	No
Family: Boidae				
<i>Calabaria reinhardtii</i>	Calabar Ground Python	LC	1	No
Family: Chamaeleonidae				
<i>Chamaeleo gracilis</i>	Slender Chameleon	LC	1	No
Family: Colubridae				
<i>Dipsadoboabrevirostris</i>	Short-headed Tree Snake	LC	1	Yes
<i>Dipsadoboaunderwoodi</i>	Underwood's Tree Snake	LC	1	No
<i>Hapsidophrys lineatus</i>	Black-lined Green Snake	LC	1	No
<i>Hapsidophrys smaragdina</i>	Emerald Snake	LC	1	No
<i>Thelotorniskirtlandii</i>	Forest Vine Snake	LC	2	No
<i>Rhamnophis aethiopissa</i>	Large-eyed Green Tree Snake	LC	1	No
Family: Elapidae				
<i>Naja guineensis</i>	Black Forest Cobra	NE	4	Yes
<i>Pseudohaje nigra</i>	Black Tree Cobra	LC	1	Yes
Family: Gekkonidae				
<i>Hemidactylus fasciatus</i>	Banded Leaf-toed Gecko	LC	3	No
<i>Hemidactylus muriceus</i>	Guinea Leaf-toed Gecko	LC	1	No
Family: Grayiidae				
<i>Grayia smithii</i>	Smith's African Water Snake	LC	2	No
Family: Lamprophiidae				
<i>Bothrophthalmus lineatus</i>	Red-black Striped Snake	LC	1	No

<i>Gonionotophis klingi</i>	Kling's File Snake	LC	1	Yes
<i>Hormonotus modestus</i>	Yellow Forest Snake	LC	1	No
<i>Mehelya poensis</i>	Equatorial File Snake	LC	1	No
Family: Natricidae				
<i>Afronatrix anoscopus</i>	African Brown Water Snake	LC	8	Yes
<i>Natriciteres variegata</i>	Variable Marsh Snake	LC	1	No
Family: Scincidae				
<i>Cophoscincopus durus</i>	Keeled Water Skink	LC	9	Yes
<i>Mochlus fernandi</i>	Fernand's Fire Skink	LC	7	No
<i>Trachylepis affinis</i>	Senegal Skink	LC	29	No
<i>Trachylepis maculilabris</i>	Speckle-lipped Skink	LC	7	No
<i>Trachylepis paucisquamis</i>	Tropical Skink	LC	4	Yes
Family: Typhlopidae				
<i>Afrotyphlops liberiensis</i>	Guinea Blind Snake	NE	1	Yes
Family: Varanidae				
<i>Varanus niloticus</i>	Nile Monitor	LC	6	No
Family: Viperidae				
<i>Atheris chlorechis</i>	Green Bush Viper	LC	3	Yes
<i>Bitis nasicornis</i>	Rhinoceros Viper	NE	1	No
<i>Bitis rhinoceros</i>	West African Gaboon Viper	LC	1	Yes
<i>Causus maculatus</i>	Spotted Night Adder	LC	3	No

587

588

589 **Table 3.** Diversity indices (Species Richness, Shannon Diversity, and Simpson Diversity)

590 generated from r/e sampling curves.

Assemblage	Diversity	Observed	Estimator	s.e.	LCL	UCL
Herpetofauna	Species richness	71.00	116.09	22.68	71.64	160.53
	Shannon diversity	28.34	29.78	1.35	27.13	32.43
	Simpson diversity	15.30	15.49	1.07	13.38	17.60
Amphibians	Species richness	38.00	38.00	1.33	38.00	40.61
	Shannon diversity	20.26	20.66	0.74	19.21	22.11
	Simpson diversity	11.91	12.05	0.82	10.44	13.65
Reptiles	Species richness	33.00	122.67	68.05	33.00	256.05
	Shannon diversity	12.69	16.97	3.25	10.61	23.34
	Simpson diversity	6.57	6.81	1.11	4.64	8.98

591

592

593 **Table 4:** Jaccard Similarity between species richness (SR) recorded in the Dugbe field survey
594 and IUCN predictive distributions for herpetofauna (total), amphibians, and reptiles. The
595 observed richness includes eight taxa not mapped or predicted by the IUCN.

Metric	Herpetofauna	Amphibians	Reptiles
Dugbe SR	71	38	33
IUCN SR	123	48	75
Species in Common	63	32	31
Jaccard Similarity Index	0.48	0.59	0.40

596

597

598 **Table 5.** Species Richness Comparison of Dugbe survey to the KBPPA and Grebo NF

599 surveys for all herpetofauna, amphibians, and reptiles.

Survey	Herps	Amphibians	Reptiles
Dugbe survey	71	38	33
Rödel & Glos 2019 (KBPPA)	48	34	14
Hillers & Rödel 2007 (Grebo NF)	34	29	5

600