



**Citation:** M. Tappi, G. Nardone, F.G. Santeramo (2022). On the relationships among durum wheat yields and weather conditions: evidence from Apulia region, Southern Italy. *Bio-based and Applied Economics* 11(2): 123-130. doi: 10.36253/bae-12160

**Received:** October 4, 2021

**Accepted:** April 27, 2022

**Published:** August 30, 2022

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Competing Interests:** The Author(s) declare(s) no conflict of interest.

**Editor:** Simone Cerroni.

**ORCID**

MT: 0000-0003-0682-5719

GN: 0000-0003-3816-0993

FGS: 0000-0002-9450-4618

Paper presented at the 10th AIEAA Conference

## On the relationships among durum wheat yields and weather conditions: evidence from Apulia region, Southern Italy

MARCO TAPPI\*, GIANLUCA NARDONE, FABIO GAETANO SANTERAMO

*University of Foggia (Italy)*

\* Corresponding author. E-mail: marco.tappi@unifg.it

**Abstract.** The weather index-based insurances may help farmers to cope with climate risks overcoming the most common issues of traditional insurances. However, the weather index-based insurances present the limit of the basis risk: a significant yield loss may occur although the weather index does not trigger the indemnification, or a compensation may be granted even if there has not been a yield loss. Our investigation, conducted on Apulia region (Southern Italy), aimed at deepening the knowledge on the linkages between durum wheat yields and weather events, i.e., the working principles of weather index-based insurances, occurring in susceptible phenological phases. We found several connections among weather and yields and highlight the need to collect more refined data to catch further relationships. We conclude opening a reflection on how the stakeholders may make use of publicly available data to design effective weather crop insurances.

**Keywords:** climate change, farming system, phenological phase, risk, weather insurance.

**JEL codes:** G22, Q14, Q18, Q54.

---

### INTRODUCTION

Farming activities are exposed and vulnerable to several risks, among which the weather risks are increasingly frequent and impactful due to climate change (Conradt et al., 2015). Among the several strategies available to reduce the weather impacts on farming systems, e.g., pest control, financial saving, agricultural and structural diversification (Vroege and Finger, 2020), the crop insurance programs can play an important role (Di Falco et al., 2014). In recent years, the attention for the weather index-based insurances (WIBIs) has been growing mainly because these tools may help to overcome some of the challenges associated with traditional indemnity-based insurances, e.g., asymmetric information, high transaction costs, moral hazard, and adverse selection (Norton et al., 2013; Dalhaus and Finger, 2016; Belissa et al., 2019; Ceballos et al., 2019). Differently from the traditional insurances, which provide pay-outs depending on actual yield losses, WIBIs indem-

nify the farmers when an index, computed on rainfall or temperature and highly correlated with farms performance (e.g., yields), is triggered (Conradt et al., 2015; Dalhaus and Finger, 2016). Therefore, farmers will be indemnified when the index exceeds a pre-determined threshold (Belissa et al., 2019). Moreover, WIBIs can be manipulated neither by the insurers or the insured because they are collected from historical and current dataset provided by recognized bodies (Belissa et al., 2020; Vroege et al., 2021). However, WIBIs present a limit, namely basis risk: a significant yield loss may occur even if the weather index does not trigger the payment (Conradt et al., 2015; Dalhaus et al., 2018) or a compensation may be granted even if there has not been a yield loss (Heimfarth and Musshoff, 2011). The contribution of our study is at least twofold: first, we provide empirical evidence on how yields and weather conditions are correlated, more specifically, we deepen the knowledge on the linkages between durum wheat yields and weather events occurring in susceptible phenological stages; second, we start a reflection on how stakeholders may make use of publicly available data to design an effective crop insurance scheme. We focused on the Apulia region (Southern Italy) which is the main national producer of durum wheat: almost a thousand of tons of production, i.e., accounting for 25% of the Italian durum wheat production, and about 344 thousand cultivated hectares, i.e., accounting for 28% of the Italian area utilized to grow durum wheat (ISMEA, 2020).

#### THE ITALIAN CROP INSURANCE SYSTEM

The Italy boasts a long tradition of public subsidies for agricultural risk management. The “Fondo di Solidarietà Nazionale” (FSN) was instituted in 1974 to finance both insurance policies and ex-post payments (Enjolras et al., 2012). Moreover, the EU Common Agricultural Policy allocated funds for agricultural insurances (art. 37 of EU Reg. 1305/2013) to cope economic losses due to adverse weather conditions, plant diseases, epizooties, and parasitic infestations (Santeramo et al., 2016; Rogna et al., 2021). Despite the public interventions, the participation level to insurance programs remains low (i.e., around 15 percent) mainly due to high costs of bureaucracy (i.e., complexity of procedures), delays in payments, lack of experience with crop insurance contracts or lack of high-quality information on existing insurance tools (Santeramo, 2019). The role of Defense Consortia, introduced both to facilitate the match of insurers and farmers in the subsidized crop insurance market and to reduce the asymmetric information, is not

negligible. It emerges a North-South territorial dualism that affects farmers participation: Defence Consortia are more effective in Northern Italy than in the Southern Italy and, also, the strong presence of producer organizations and cooperatives aggregates the crop insurance's demand in the Northern Italy (Santeramo et al., 2016). Moreover, farmers who trust more in the intermediaries assisting them are inclined to adopt insurance tools to cope the risk of production loss, while risk averse farmers tend to implement other risk management strategies as crop or financial diversification (Trestini et al., 2018). In Italy, only the 9.9 percent of Utilised Agricultural Area is covered by insurance contracts and 20.9 percent of production value is insured (ISMEA, 2021). According to a survey conducted by ISMEA in 2018 on low participation to the subsidized agricultural insurance systems, most Italian farmers renounce to subscribe insurance contracts due to economic reasons, highlighting the high costs of policies. The share of farmers who believe that their farms are not exposed to specific risks or who have had negative experiences when receiving compensation, losing trust on insurance market systems, is also not negligible. Indeed, Giampietri et al., 2020 found that the trust affects the decision-making process: under uncertainty, the trust may substitute the knowledge also overcoming the lack of experience, therefore, strong communication campaigns to improve farmers' participation are recommended. Moreover, focusing on the WIBIs, also subsidized by the Measure 17 of National Rural Development Program 2014-2020, a lack of knowledge emerged among big insured farmers, i.e., WIBIs were unknown to 93 percent of them (ISMEA, 2020). Furthermore, some farmers believe that index-based insurances are inadequate to manage the weather risks due to the distrust of the objectivity of the indexes and parameters used, also showing an aversion to any future subscriptions. Clearly, it is necessary to improve the appeal and communication of these innovative risk management tools, also considering that any intervention aimed at promoting farmer participation should improve the competition among insurance providers, also reducing at the same time the asymmetric information and opportunistic behaviour (Menapace et al., 2016; Rogna et al., 2021; Santeramo and Russo, 2021). In this complex scenario, we estimate the yield response equation to investigate the responsiveness of yield to climate, deepening the working principles of weather index-based insurance, through a case study on durum wheat crop in the Apulia region, also animating the debate on the use of publicly available data to the development of an effective and attractive tool to manage climatic risk in agriculture.

## DATA AND RESEARCH METHODOLOGY

An agronomic review on durum wheat allowed us to identify sensitive phenological stages of durum wheat in Apulia region and those critical weather events occurring in certain phenological stages that may cause significant production losses (Table 1).

Cold sensitivity is higher during the germination phase that occurs 10-15 days after sowing in which temperatures of few degrees centigrade below zero may cause considerable damages (Baldoni and Giardini, 2000; Angelini, 2007; Disciplinare di Produzione Integrata della Regione Puglia, 2021). Likewise, temperatures of few degrees centigrade below zero during the stem elongation phase may cause stems death and serious damages to the tissue of the internodes (Baldoni and Giardini, 2000; Angelini, 2007; Disciplinare di Produzione Integrata della Regione Puglia, 2021). Flowering stage occurs in late May and lasts about 10 days in which wheat crop is highly sensitive to cold stress that may cause death of flowers (Angelini, 2007; Baldoni and Giardini, 2000; Disciplinare di Produzione Integrata della Regione Puglia, 2021). Heat and drought stress during susceptible flowering and grain filling stages (i.e., after flowering, until the first decade of July) may cause considerable reductions in wheat yield and quality, leading the acceleration of leaf senescence process, reducing photosynthesis, causing oxidative damage, pollen sterility, also reducing physiological and metabolic imbalances, photosynthesis, grain numbers and weight (Angelini, 2007; Asseng et al., 2011; Li et al., 2013; Farooq et al., 2014; Rezaei et al., 2015; Zampieri et al., 2017; Makinen

et al., 2018). Heavy rainfall during the entire crop cycle may cause significant production losses due to the proliferation of pathogens, nutrient leaching, soil erosion, inhibition of oxygen uptake by roots (i.e., hypoxia or anoxia), waterlogging and lodging (Zampieri et al., 2017; Makinen et al., 2018).

Furthermore, we collected yearly total production (tons) and area harvested (hectares) data for durum wheat crop from the National Institute of Statistics (ISTAT), from 2006 to 2019, for each province of Apulia region, also calculating the respective yields (tons/hectare). Then, for the same time-period, we collected 10-days frequency weather data from six synoptic weather stations of the Institute for Environmental Protection and Research (ISPRA), one for each province of Apulia region: Bari (BA), Barletta-Andria-Trani (BT), Brindisi (BR), Foggia (FG), Lecce (LE), Taranto (TA). Weather data include 10-days average minimum temperature ( $^{\circ}\text{C}$ ), i.e., the average of daily minimum temperatures, 10 days average maximum temperature ( $^{\circ}\text{C}$ ), i.e., the average of daily maximum temperatures, and 10-days cumulative precipitation (mm), i.e., the average of daily precipitation.

Details on collected variables are shown in Table 2.

Our empirical approach is based on a panel data model that includes fixed effect (i.e., it is a major advantage of the panel rather than cross-sectional regression) both to control for unobservable variables such as seed varieties or soil quality that may vary across the space, i.e., provinces, and to catch the variation across the time within the Apulian provinces (Tack et al., 2015; Blanc and Schlenker, 2017; Kolstad and Moore, 2020).

**Table 1.** Phenological stages, weather events and critical limits of durum wheat in Apulia region.

Phenological stage	Weather event	Time interval	Critical limit	Reference
Sowing	Cold	From the first decade of November to the first decade of December	Temperature < 0 $^{\circ}\text{C}$	Baldoni and Giardini, 2000; Angelini, 2007; Disciplinare di produzione integrata della Regione Puglia, 2021
Germination	Cold	From the second decade of November to the second decade of December	Temperature < 0 $^{\circ}\text{C}$	
Stem elongation	Cold	From the second decade of March to the third decade of April	Temperature < 0 $^{\circ}\text{C}$	Baldoni and Giardini, 2000; Angelini, 2007 Angelini, 2007; Disciplinare di produzione integrata della Regione Puglia, 2021
Flowering	Cold	From the second decade of May to the first decade of June	Temperature < 0 $^{\circ}\text{C}$	
	Heat, drought		Temperature > 30-31 $^{\circ}\text{C}$	Angelini, 2007; Rezaei et al., 2015
Grain filling	Heat, drought	From the second decade of June to the first decade of July	Temperature > 34 $^{\circ}\text{C}$	Angelini, 2007; Asseng et al., 2011; Rezaei et al., 2015; Zampieri et al., 2017; Makinen et al., 2018
All phases	Excessive rainfall	From first decade of November to the first decade of July	Rainfall > 40 mm/day	Makinen et al., 2018

**Table 2.** Details on collected variables.

Variable (unit)	Frequency	Time-period	Province	Weather station - province (no. of obs, SR in km <sup>2</sup> )	Source
durum wheat yield (tons/hectares)	Yearly			-	ISTAT
average minimum temperature (°C)		2006-2019	Bari (BA) Bari (BA) Barletta-Andria-Trani (BAT)	Bari - BA (501, 5.138) Trani - BT (144, 1.543)	
average maximum temperature (°C)	10-days		Brindisi (BR) Foggia (FG) Lecce (LE) Taranto (TA)	Brindisi - BR (471, 1.839) Monte Sant'Angelo - FG (504, 7.008) Lecce - LE (471, 2.799) Marina di Ginosa - TA (471, 2.437)	ISPRA, UCEA, ARPA
cumulative precipitation (mm)					

Notes: missing data have been integrated including Research Unit for Climatology and Meteorology (UCEA) and Regional Agency for the Protection of the Environment (ARPA) datasets. Table includes no. of observations and spatial resolution (SR) of weather stations.

The relationship between durum wheat yields and weather events is synthesized as follows:

$$y_{it} = f(w_{it}) + \mu_i + \theta_t + \varepsilon_{it}$$

where  $y_{it}$  is the yield over the space ( $i$ ) and time ( $t$ ) as function ( $f$ ) of weather ( $w_{it}$ ), also including fixed effects over space ( $\mu_i$ ) and time ( $\theta_t$ ), error term and “controls” refers to other relevant exogenous variables ( $\varepsilon_{it}$ ) (Kolstad and Moore, 2020). More specifically, we conducted temporal and spatial autocorrelation identifying those contiguous provinces having a larger shared borders for a twofold check: (i) verify if the weather events occurring in a province may affect durum wheat yields in the contiguous province; (ii) control if the yields may be affect-

ed by weather events occurring at time  $t-1$ . Undoubtedly, both environmental and agronomic factors may justify the extreme variability of the durum wheat yield across the Apulian provinces: Foggia shows the highest average durum wheat yields while Lecce shows the lowest average yields, although it is characterized by lower yield variability than other provinces as Brindisi that, on the contrary, is more affected by environmental and agronomic factors, reason why it may benefit of crop insurance programs more than other provinces to cope yields fluctuations (Table 3).

## RESULTS

Our results clearly show that a relationship links weather conditions and production yields in the Apulia region. More specifically, precipitation seem to have a negative effect on durum wheat yields (Table 4).

However, controlling by spatial and temporal autocorrelation, the effects of temperatures have been caught. Minimum temperatures negatively affect durum wheat yields, while maximum temperatures positively affect the yields, both in a non-linear way. Indeed, we included the squares of weather variables to catch the nonlinearity, in other terms, the trade-off between weather and yields (Blanc and Schlenker, 2017). Our results clearly highlight that the weather affects the yields in a nonlinear way, therefore, variables have a statistically significant inverted-U shape relationship

**Table 3.** Durum wheat yields (tons/hectare) among Apulian provinces.

	Average	Minimum	Maximum	Standard deviation
Bari	0.234	0.170	0.306	0.045
BAT	0.224	0.200	0.260	0.020
Brindisi	0.285	0.180	0.420	0.071
Foggia	0.314	0.200	0.420	0.047
Lecce	0.189	0.160	0.220	0.018
Taranto	0.244	0.100	0.350	0.057

Notes: data include yearly durum wheat yield from 2006 to 2020. Source: ISTAT, 2020.

**Table 4.** Effects of weather variables on durum wheat yield.

Variables	Panel prov FE time trend	Panel temporal correlation prov FE time trend	Panel spatial correlation prov FE time trend	Panel temporal correlation prov FE time trend	Panel spatial correlation prov FE time trend
Temperature (min)	-0.00764 (0.10641)	-0.00124 (0.11715)	-0.46909*** (0.17058)	-0.45553** (0.18731)	
Temperature (min) sq.	0.00049 (0.00296)	-0.00023 (0.00320)	0.00892* (0.00490)	0.01384** (0.00544)	
Temperature (max)	0.22572 (0.14125)	0.28286* (0.15378)	0.61165** (0.25587)	0.66801** (0.27703)	
Temperature (max) sq.	-0.00523* (0.00278)	-0.00612** (0.00299)	-0.01530*** (0.00515)	-0.02022*** (0.00568)	
Precipitation	-0.01646** (0.00799)	-0.01625* (0.00844)	-0.03939** (0.01819)	-0.04670** (0.01954)	
Precipitation sq.	0.00008 (0.00006)	0.00007 (0.00006)	0.00019 (0.00017)	0.00024 (0.00018)	
Yield (lag)	-	0.10464*** (0.02153)	-	-0.09290*** (0.03579)	
Temperature (min) contig.	-	-	0.23065*** (0.06565)	0.18642*** (0.07019)	
Temperature (max) contig.	-	-	0.00822 (0.10765)	0.04557 (0.11545)	
Precipitation contig.	-	-	0.00537 (0.00704)	0.00771 (0.00837)	
Observations	1,837	1,638	914	833	
Number of id	6	6	4	4	

Notes: panel regression model was processed in STATA software. It includes provincial fixed effect, time trend, temporal (i.e., yield lag), and spatial (contiguous weather variables) autocorrelation.

Standard errors in parentheses.

\*\*\* Significant at the 1 percent level.

\*\* Significant at the 5 percent level.

\* Significant at the 10 percent level.

(Schlenker and Roberts, 2009; Lobell et al., 2011). Last but not least, minimum temperatures may affect the contiguous provinces. According to the scientific literature, any excess (or deficit) of temperature and precipitation (or their combinations) may cause severe yield losses on durum wheat (Baldoni and Giardini, 2000; Angelini, 2007; Asseng et al., 2011; Li et al., 2013; Farooq et al., 2014; Rezaei et al., 2015; Zampieri et al., 2017; Makinen et al., 2018). Furthermore, we estimated the model for each phenological phase of durum wheat to capture the potential heterogeneity in the effect of weather variables, also controlling by spatial and temporal autocorrelation. Our results show that the relationship between weather variables and yields is valid only for some weather variables in certain phenological phases. More specifically, the maximum temperatures and precipitation positively affect durum wheat yield

in a nonlinear way when occur in the germination and grain filling stages, respectively (Table 5).

Moreover, minimum temperatures may affect the contiguous provinces. Clearly, ten-days data we have collected does not highlight the dynamics between weather events occurring in certain phenological stages and durum wheat yields mainly because the impacts of daily weather are not captured. Moreover, most variables are not statistically significant: this limit opens a reflection on data disaggregation level and on the need to collect more spatially and temporally refined data, also laying the foundations for the development of an effective index that reflects the responsiveness of the yields to climatic conditions to be implemented in the WIBIs. The evidence resulting from our econometric model on phenological stages is also in contrast with the literature: germination stage is highly sensitive to cold stress (Bal-

**Table 5.** Effects of weather variables on yield by phase.

Variables	sowing	germination	stem elongation	flowering	grain filling
Yield (lag)	-0.11883 (0.20660)	0.05952 (0.20523)	0.17798* (0.09219)	-0.04474 (0.18593)	0.09403 (0.14041)
Temperature (min)	0.95845 (2.53724)	-0.00051 (1.74362)	0.50020 (1.26379)	-1.32087 (4.06620)	-0.65587 (3.83238)
Temperature (min) sq.	-0.01783 (0.11363)	0.01530 (0.08655)	-0.01201 (0.05223)	0.03550 (0.10882)	0.02171 (0.08353)
Temperature (max)	3.15220 (12.35641)	23.00804** (10.88917)	-2.73726 (2.21349)	7.62398 (8.51643)	-1.65011 (6.74553)
Temperature (max) sq.	-0.15964 (0.35336)	-0.76330** (0.33477)	0.06023 (0.05582)	-0.15868 (0.15987)	0.01396 (0.11320)
Precipitation	0.04601 (0.12015)	-0.07450 (0.11228)	-0.03735 (0.07473)	-0.43463 (0.42173)	0.42332* (0.24351)
Precipitation sq.	-0.00034 (0.00088)	0.00054 (0.00084)	0.00049 (0.00101)	0.01188 (0.01680)	-0.00826* (0.00463)
Temperature (min) contig.	1.05294** (0.41397)	0.86957** (0.35021)	0.62187*** (0.17188)	0.52210 (0.35845)	0.55304** (0.23765)
Temperature (max) contig.	0.38942 (1.25128)	0.17524 (1.33537)	-0.06474 (0.34861)	0.22627 (0.52741)	0.00512 (0.37530)
Precipitation contig.	-0.05370 (0.05168)	0.01278 (0.04199)	-0.01394 (0.03275)	-0.10017 (0.11446)	-0.05635 (0.04998)
Observations	42	44	125	43	67
Number of id	4	4	4	4	4

Notes: panel regression model was processed in STATA software. It includes provincial fixed effect, time trend, temporal (i.e., yield lag), and spatial (contiguous weather variables) autocorrelation.

Notes: standard errors in parentheses

\*\*\* Significant at the 1 percent level.

\*\* Significant at the 5 percent level.

\* Significant at the 10 percent level.

doni and Giardini, 2000, Angelini, 2007; *Disciplinare di Produzione Integrata della Regione Puglia*, 2021), while there are not evidences on heat stress during this stage. However, our study may help the debate suggesting precise directions for the future research.

## CONCLUSIONS

Participating in index-based crop insurance schemes is a key challenge to improve the resilience of farming systems and adopting effective subsidies to enhance participation in the schemes is a pressing goal for policymakers. In this complex scenario, we investigated how temperatures and precipitation are correlated with yields data to reflect on potential designs for the index-based insurance schemes. While not novel (e.g., Chen et al., 2014), we found that weather changes affect durum wheat yields in a nonlinear way and some weather events occurring in certain phenological phases may

have an impact on the yields. Our results are important to show that even with aggregated data the evidence is striking. However, focusing on phenological stages, our findings are in contrast with the literature highlighting the complexity of the phenomenon and the need to rely on more temporally and spatially disaggregated data. Although we provided clear evidence on the weather-yield relationship, it is impossible to design a WIBI using 10-days weather data. Therefore, our contribution may help the debate suggesting precise directions for the future research: first, a major effort should be devoted to the collection of weekly or daily weather observations, also identifying empirical damage thresholds that can be verified at farm-level, as well as the collection of production area or municipal data; a promising approach could be the Growing Degree Days tool so as to calibrate the more precisely the growing stages in a view to a better explanation of weather risks on crop performances (Conradt et al., 2015; Dalhaus et al., 2018; Lollato et al., 2020); last but not least, the design of the index-

based insurance schemes needs of further investigation because establishing a triggering index is a major challenge for the *stakeholders* involved in the implementation of the insurance schemes. The debate on crop insurance schemes is still vivid, and it will be so also in the next decade due to the central role that the risk management (old and novel) tools will have in the new CAP (Meuwissen et al., 2018; Severini et al., 2019; Cordier and Santeramo, 2020).

## REFERENCES

- Angelini, R. (2007). *Coltura & cultura. Il grano*. ART SpA - Bologna.
- Asseng, S., Foster, I.A.N., and Turner, N.C. (2011). The impact of temperature variability on wheat yields. *Global Change Biology*, 17(2), 997-1012.
- Auffhammer, M., Hsiang, S. M., Schlenker, W., & Sobel, A. (2013). Using weather data and climate model output in economic analyses of climate change. *Review of Environmental Economics and Policy*, 7(2), 181-198.
- Baldoni, R. and Giardini, L. (2000). *Coltivazioni erbacee. Cereali e proteaginose*. In Toderi, G., and D'Antuono L.F., *Frumento (Triticum sp.pl.)*. Patron Editore.
- Belissa, T., Bulte, E., Cecchi, F., Gangopadhyay, S., and Lensink, R. (2019). Liquidity constraints, informal institutions, and the adoption of weather insurance: A randomized controlled Trial in Ethiopia. *Journal of Development Economics*, 140, 269-278.
- Belissa, T., Lensink, R., and Winkel, A. (2020). Effects of Index Insurance on Demand and Supply of Credit: Evidence from Ethiopia. *American Journal of Agricultural Economics*, 102(5), 1511-1531.
- Blanc, E., & Schlenker, W. (2017). The use of panel models in assessments of climate impacts on agriculture. *Review of Environmental Economics and Policy*, 11(2), 258-279.
- Ceballos, F., Kramer, B., and Robles, M. (2019). The feasibility of picture-based insurance (PBI): Smartphone pictures for affordable crop insurance. *Development Engineering*, 4, 100042.
- Chen, C.C., McCarl, B.A., and Schimmelpfennig, D.E. (2004). Yield variability as influenced by climate: A statistical investigation. *Climatic Change*, 66(1), 239-261.
- Conradt, S., Finger, R., and Spörri, M. (2015). Flexible weather index-based insurance design. *Climate Risk Management*, 10, 106-117.
- Conradt, S., Finger, R., and Bokusheva, R. (2015). Tailored to the extremes: Quantile regression for index-based insurance contract design. *Agricultural economics*, 46(4), 537-547.
- Cordier, J. and Santeramo, F. (2020). Mutual funds and the Income Stabilisation Tool in the EU: Retrospect and Prospects. *EuroChoices*, 19(1), 53-58.
- Dalhaus, T. and Finger, R. (2016). Can gridded precipitation data and phenological observations reduce basis risk of weather index-based insurance? *Weather, Climate, and Society*, 8(4), 409-419.
- Dalhaus, T., Musshoff, O., and Finger, R. (2018). Phenology information contributes to reduce temporal basis risk in agricultural weather index insurance. *Scientific reports*, 8(1), 1-10.
- Disciplinare di Produzione Integrata della Regione Puglia (2021). [http://burp.regione.puglia.it/documents/10192/56088259/DET\\_67\\_2\\_3\\_2021.pdf/bb22795d-6335-4498-bc3a-7a13bf8d140d](http://burp.regione.puglia.it/documents/10192/56088259/DET_67_2_3_2021.pdf/bb22795d-6335-4498-bc3a-7a13bf8d140d). Accessed 31 August 2021
- Di Falco, S.D., Adinolfi, F., Bozzola, M., and Capitanio, F. (2014). Crop insurance as a strategy for adapting to climate change. *Journal of Agricultural Economics*, 65(2), 485-504.
- Enjolras, G., Capitanio, F., and Adinolfi, F. (2012). The demand for crop insurance: Combined approaches for France and Italy. *Agricultural economics review*, 13(389-2016-23488), 5-22.
- Farooq, M., Hussain, M., and Siddique, K. H. (2014). Drought stress in wheat during flowering and grain-filling periods. *Critical reviews in plant sciences*, 33(4), 331-349.
- Giampietri, E., Yu, X., & Trestini, S. (2020). The role of trust and perceived barriers on farmer's intention to adopt risk management tools. *Bio-based and Applied Economics Journal*, 9(1050-2021-213), 1-24.
- Kolstad, C. D., & Moore, F. C. (2020). Estimating the economic impacts of climate change using weather observations. *Review of Environmental Economics and Policy*, 14(1), 1-24.
- Li, Y.F., Wu, Y., Hernandez-Espinosa, N., and Peña, R. J. (2013). Heat and drought stress on durum wheat: Responses of genotypes, yield, and quality parameters. *Journal of Cereal Science*, 57(3), 398-404.
- Lobell, D. B., Bänziger, M., Magorokosho, C., & Vivek, B. (2011). Nonlinear heat effects on African maize as evidenced by historical yield trials. *Nature climate change*, 1(1), 42-45.
- Lollato, R.P., Bavia, G.P., Perin, V., Knapp, M., Santos, E.A., Patrignani, A., and DeWolf, E.D. (2020). Climate-risk assessment for winter wheat using long-term weather data. *Agronomy Journal*, 112(3), 2132-2151.
- Mäkinen, H., Kaseva, J., Trnka, M., Balek, J., Kersebaum, K. C., Nendel, C., Gobin, A., Olesen, J.E., Bindi,

- M., Ferrise, R., Moriondo, M., Rodriguez, A., Ruiz-Ramos, M., Takác, J., Bezák, P., Ventrella, D., Ruget, F., Capellades, G., and Kahiluoto, H. (2018). Sensitivity of European wheat to extreme weather. *Field Crops Research*, 222, 209-217.
- Menapace, L., Colson, G., and Raffaelli, R. (2016). A comparison of hypothetical risk attitude elicitation instruments for explaining farmer crop insurance purchases. *European Review of Agricultural Economics*, 43(1), 113-135.
- Meuwissen, M. P., de Mey, Y., and van Asseldonk, M. (2018). Prospects for agricultural insurance in Europe. *Agricultural Finance Review*.
- Norton, M.T., Turvey, C., and Osgood, D. (2013). Quantifying spatial basis risk for weather index insurance. *The Journal of Risk Finance*.
- Rezaei, E.E., Webber, H., Gaiser, T., Naab, J., and Ewert, F. (2015). Heat stress in cereals: mechanisms and modelling. *European Journal of Agronomy*, 64, 98-113.
- Rogna, M., Schamel, G., and Weissensteiner, A. (2021). The apple producers' choice between hail insurance and anti-hail nets. *Agricultural Finance Review*.
- Santeramo, F. G., Goodwin, B. K., Adinolfi, F., and Capitanio, F. (2016). Farmer participation, entry and exit decisions in the Italian crop insurance programme. *Journal of Agricultural Economics*, 67(3), 639-657.
- Santeramo, F.G. and Ford Ramsey, A. (2017). Crop Insurance in the EU: Lessons and Caution from the US. *EuroChoices*, 16(3), 34-39.
- Santeramo, F.G. (2018). Imperfect information and participation in insurance markets: evidence from Italy. *Agricultural Finance Review*, 78(2), 183-194.
- Santeramo, F. G. (2019). I learn, you learn, we gain experience in crop insurance markets. *Applied Economic Perspectives and Policy*, 41(2), 284-304.
- Santeramo, F.G. and Russo, I. (2021). Aspetti comportamentali della partecipazione ai programmi di assicurazione agricola agevolata nell'Italia meridionale. *Italian Review of Agricultural Economics*, 76(2), 73-90.
- Schlenker, W., & Roberts, M. J. (2009). Nonlinear temperature effects indicate severe damages to US crop yields under climate change. *Proceedings of the National Academy of sciences*, 106(37), 15594-15598.
- Severini, S., Biagini, L., and Finger, R. (2019). Modeling agricultural risk management policies—The implementation of the Income Stabilization Tool in Italy. *Journal of Policy Modeling*, 41(1), 140-155.
- Tack, J., Barkley, A., & Nalley, L. L. (2015). Effect of warming temperatures on US wheat yields. *Proceedings of the National Academy of Sciences*, 112(22), 6931-6936.
- Trestini, S., Giampietri, E., & Smiglak-Krajewska, M. (2018). Farmer behaviour towards the agricultural risk management tools provided by the CAP: a comparison between Italy and Poland (No. 2038-2018-2993).
- Vroege, W. and Finger, R. (2020). Insuring Weather Risks in European Agriculture. *EuroChoices*, 19(2), 54-62.
- Vroege, W., Bucheli, J., Dalhaus, T., Hirschi, M., and Finger, R. (2021). Insuring crops from space: the potential of satellite-retrieved soil moisture to reduce farmers' drought risk exposure. *European Review of Agricultural Economics*, 48(2), 266-314.
- Woodard, J. D. and Garcia, P. (2008). Weather derivatives, spatial aggregation, and systemic risk: implications for reinsurance hedging. *Journal of Agricultural and Resource Economics*, 34-51.
- Zampieri, M., Ceglar, A., Dentener, F., and Toreti, A. (2017). Wheat yield loss attributable to heat waves, drought and water excess at the global, national and subnational scales. *Environmental Research Letters*, 12(6), 064008.