



Citation: Mitka J., Novikov A., Rottensteiner W.K. (2021) The taxonomic circumscription of *Aconitum* subgenus *Aconitum* (Ranunculaceae) in Europe. *Webbia. Journal of Plant Taxonomy and Geography* 76(1): 11-45. doi: 10.36253/jopt-10006

Received: November 10, 2020

Accepted: December 12, 2020

Published: April 27, 2021

Copyright: © 2021 Mitka J., Novikov A., Rottensteiner W.K. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/webbia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Competing Interests: The Author(s) declare(s) no conflict of interest.

Editor: Riccardo M. Baldini

ORCID

JM: <https://orcid.org/0000-0001-8472-1742>

AN: <https://orcid.org/0000-0002-0112-5070>

The taxonomic circumscription of *Aconitum* subgenus *Aconitum* (Ranunculaceae) in Europe

JÓZEF MITKA^{1,*}, ANDRIY NOVIKOV², WALTER K. ROTTENSTEINER³

¹ Jagiellonian University in Kraków, Institute of Botany, Botanical Garden, Kopernika 27, 31-501 Kraków, Poland

² State Natural History Museum NAS of Ukraine, Teatralna str. 18, Lviv 79008, Ukraine

³ Klosterwiesgasse 12, A-8010 Graz, Austria

*Corresponding author. E-mail: j.mitka@uj.edu.pl

Abstract. In this article, we present a revised taxonomic circumscription of *Aconitum* subg. *Aconitum* (Ranunculaceae) in Europe. In total, the subgenus contains some 250 species with the major center of diversity in Eastern Asia. Altogether 94 taxa (species and infraspecific taxa, including hybrids) occur in Europe. Among them, 22 are native species, and 28 are nothospecies (including hybrid formulae). The research is based on former (since Linnaeus) and recent species diagnoses integrating herbarium and field studies carried out in the Alps, Carpathians, Balkans, Spanish Sierra Nevada, Sudetes, and Corsica. The subgenus includes three sections in Europe: the diploid sect. *Cammarum*, the tetraploid sect. *Aconitum*, and the monospecific, allopolyploid sect. *Angustifolium*. Additionally, a triploid, hybridogenous nothosection *Acomarum* (sect. *Aconitum* × sect. *Cammarum*) is presented. For each species, type citation, a concise morphological description, including infraspecific variation and hybridization, geographical distribution, and iconography sources are given. Also, a key to the determination of all taxa is presented. The proposed system scrutinizes former and recent species concepts and gives a base for further studies on the genus' phylogeny and biotechnology.

Keywords: *Aconitum*, geographical distribution, Europe, hybrids, Linnaean taxonomy, nomenclature, species concept.

INTRODUCTION

The genus *Aconitum* L. (monkshood) comprises ca. 300–400 species distributed in temperate regions of the northern hemisphere, with a center of diversity in the eastern Himalaya, southwestern China, and Japan (Kadota 1987; Li and Kadota 2001; Luo et al. 2005). From a total number of ca. 250 species belonging to the subgenus *Aconitum*, 22 species may be found in Europe, with eight species occurring exclusively in the Carpathian and Balkans Mts (Boroń et al. 2020). The subgenus includes both the “temperate” forest (diploid) and “cold” high-mountain (tetraploid) species, which are differentiated both morphologically (Starmühler 1998; Mitka 2003; Novikoff

and Mitka 2011) and cytogenetically (Ilnicki and Mitka 2009, 2011).

Genera *Aconitum* L., *Delphinium* L., *Consolida* (DC.) S.F Gray, *Aconitella* Spach, and *Staphisagria* J. Hill form the monophyletic tribe *Delphinieae* Schröd., subtribe *Delphininae* Benth. & Hook. f. (Tamura 1993; Keener et al. 1999; Turland and Barrie 2001; Jabbour and Renner 2011a, b). Zygomorphic flowers and the presence of diterpene alkaloids turned out to be the synapomorphies to this taxonomic group (Johansson 1995; Jabbour et al. 2009, 2014). Genus *Aconitum* consists of the following monophyletic subgenera: subg. *Aconitum*, subg. *Fletcherum* (Tamura) Y.Hong & Q.E.Yang, subg. *Galeata* (Rapaics) Y.Hong & Q.E.Yang, and subg. *Lycocotonum* (DC.) Peterm. (Kita et al. 2000; Utelli and Ito 2000; Luo et al. 2005; Jabbour and Renner 2011b; Hong et al. 2017). The monophyly and taxonomic rank of subg. *Anthora* (Rapaics) Peterm. is unclear and deserves further studies (Novikoff and Mitka 2015). The endemic of the Qinghai-Tibetan Plateau *A. gymnanthrum* Maximowicz has been recently transferred from the monotypic *Aconitum* subg. *Gymnaconitum* (Stapf) Rapaics to an independent genus *Gymnaconitum* (Stapf) Wei Wang & Z.D.Chen based on molecular data (Wang et al. 2013).

Aconitum subg. *Aconitum* is known for its remarkable morphological plasticity and notorious hybridization, leading to difficulties in the circumscription of taxa (Kadota 1981; Tutin et al. 1993; Sutkowska et al. 2013, 2017a, b). To clarify the sectional treatment of the subgenus, we used both cytogenetic and morphological criteria (Starmühler 1996b, 2000; Ilnicki and Mitka 2009, 2011; Joachimiak et al. 1999; Mitka and Starmühler 2000; Novikoff and Mitka 2011).

The subgenus *Aconitum* in Europe has basic (monoploid, i.e., in a single complete set of chromosomes) number $x = 8$ and ploidy levels $2x$, $3x$, $4x$, and $6x$. It consists of the diploid sect. *Cammarum* DC. ($2n (2x) = 16$), the triploid nothosection *Acomarum* Starm. ($2n (3x) = 24$), the tetraploid sect. *Aconitum* ($2n (4x) = 32$), and the monospecific hexaploid sect. *Angustifolium* (Seitz) Rottensteiner, represented by the allopolyploid *A. angustifolium* Rchb. ($2n (6x) = 48$) (Seitz 1969; Zieliński 1982a, b; Simon et al. 2001; Ilnicki and Mitka 2009, 2011).

Section *Aconitum* includes high-mountain species that inhabit open sites in the subalpine and alpine zones. Section *Cammarum* is represented by forest species, which usually are distributed in montane (up to ca. 1150–1200 m above sea level), foreland and lowland/upland areas (Mitka 2000, 2002; Novikoff and Mitka 2011). Intersectional hybrids (*Aconitum* sect. *Aconitum* × *Aconitum* sect. *Cammarum*) are circumscribed within

nothosect. *Acomarum* Starm. (Wacław ska-Ćwiertnia and Mitka 2016; Starmühler 2001).

We also include to the list *A. carmichaelii* from the section *Euchylodea* with $2n (4x) = 32$, $2n (6x) = 48$ or, often, $2n (8x) = 64$ (Yang 1996; Li and Kadota 2001). In Europe, *A. carmichaelii* is used as an ornamental plant, while in China (native area) it is mostly cultivated for medicinal purposes (Yu et al. 2016; Zhao et al. 2017).

In the present article, we propose an annotated Linnaean taxonomic system of *Aconitum* subg. *Aconitum* in Europe. It is based on the classical works by Reichenbach (1819, 1821, 1827, 1840), Gáyer (1906, 1909, 1911, 1912), Götz (1967), Seitz (1969), Skalický (1982), and Flora Europaea (Tutin et al. 1993), supplemented by our recent systematic studies carried out in the Alps, Carpathians, Sudetes, and Balkans.

We adopt the following taxonomic treatment of *Aconitum* subgenus *Aconitum* in Europe (see also Appendix):

Aconitum L. subg. *Aconitum*

1. Sect. *Aconitum*

- 1A. Subsect. *Aconitum* [syn. sect. *Napellus* Wolf (DC.) subsect. *Napellus* (Wolf) Rapaics]
 - 1a. Ser. *Aconitum*
 - 1b. Ser. *Castellana* Rottensteiner, **ser. nov.**
 - 1c. Ser. *Taurica* Mucher ex Starm.
 - 1d. Nothoser. *Acorica* Mucher ex Starm. (ser. *Aconitum* × ser. *Taurica*)

1B. Subsect. *Burnatii* Rottensteiner

2. Sect. *Cammarum* DC.

- 2A. Subsect. *Cammarum* (DC.) Rapaics
 - 2a. Ser. *Variegata* Steinberg ex Starm.
 - 2b. Ser. *Toxica* (Rchb.) Mucher
 - 2c. Nothoser. *Toxigata* Starm. (ser. *Toxica* × ser. *Variegata*)

3. Nothosect. *Acomarum* Starm. (sect. *Aconitum* × sect. *Cammarum*)

- 4. Sect. *Angustifolium* (Seitz) Rottensteiner, **stat. nov.**
- 5. Sect. *Euchylodea* Rchb.
 - 5a. Ser. *Japonica* (Nakai) Kadota

MATERIAL AND METHODS

We have evaluated all recognized taxa from the *Aconitum* subg. *Aconitum* distributed in Europe based on our experience of many years of field investigations, morphological and biogeographical studies, as well as analysis of herbarium material. For each taxon, we provided comprehensive data on its distribution and morphological characteristics. We reviewed all available

sources for taxon synonymy and type citations. Basing on morphological data, a binary key for the identification of the representatives of the subgenus *Aconitum* has been developed.

The chromosome numbers were investigated in the Carpathians and Sudetes Mts (Joachimiak et al. 1999; Ilnicki and Mitka 2009, 2011; Mitka et al. 2007) or taken from the on-line DCDB database (<http://hdl.handle.net/2445/95875>; Bosch et al. 2016).

RESULTS

Key to the taxa of Aconitum subgenus Aconitum in Europe

1. Root napiform; stem ± stiff upright, rarely ± flexuous (*A. sect. Burnatii*); caudine leaves usually without reticulate venation; seeds smooth to transverse wrinkled or with transverse lamellae on one side, the uppermost segment of the leaf lobe with two, rarely with 4–6 teeth 2
- 1.* Root globose; stem upright to bent to overhanging or spreading-climbing, flexuous; caudine leaves with reticulate venation; seeds always with transverse lamellae on one side, the uppermost segment of the leaf lobe, located in its middle, with 4–12 teeth *sect. Cammarum* 35
2. Carpels sterile *nothosect. Acomarum* (*A. sect. Aconitum* × *A. sect. Cammarum*) 3
- 2.* Carpels fertile 14
3. Hood outside glabrous (or with just some single hairs) 4
- 3.* Hood outside hairy 11
4. Cauline leaves net-veined *A. superbum* × *A. variegatum*
- 4.* Cauline leaves not net-veined 5
5. Pedicel glabrous or with just some single hairs at the top; filaments glabrous or just sparsely pilose 6
- 5.* Pedicel hairy at least above the bracteoles; filaments ± densely pilose 7
6. Carpels mostly completely glabrous *A. ×acutum* (*A. tauricum* × *A. variegatum*)
- 6.* Carpels pilose on the suture *A. ×berdaui* nothosubsp. *berdaui* (*A. firmum* subsp. *firmum* × *A. variegatum* subsp. *variegatum*)
7. Pedicels with glandular hairs 8
- 7.* Pedicels without glandular hairs 9
8. Pedicel pilose with strict glandular hairs *A. ×mielichhoferi* (*A. degeneri* × *A. tauricum*)
- 8.* Pedicel pilose with curved and s-formed glandular hairs *A. pilipes* × *A. tauricum*
9. Pedicels above the bracteoles with curved, crisped and strict hairs ... *A. ×cammarum* (*A. napellus* × *A. variegatum*)
 - a. Tepals with monotonic deep blue color *fo. cammarum*
 - a.* Tepals white and blue strips *fo. bicolor*
- 9.* Pedicels above the bracteoles only with curved and crisped hairs 10
10. Bracteoles triangular to linear, 2–5 mm long *A. ×exaltatum* (*A. plicatum* × *A. variegatum* subsp. *variegatum*)
- 10.* Bracteoles lanceolate to spatulate, 5–7 mm long *A. ×schneebergense* (*A. napellus* × *A. variegatum*)
11. Cauline leaves net-veined *A. superbum* × *A. vitosanum*
- 11.* Cauline leaves not net-veined 12
12. Tepals outside with only eglandular hairs *A. ×berdaui* nothosubsp. *walasii* (*A. firmum* subsp. *moravicum* × *A. variegatum* subsp. *variegatum*)
- 12.* Tepals outside with eglandular and glandular hairs 13
13. Tepals outside with curved eglandular hairs and s-formed glandular hairs *A. napellus* × *A. pilipes*
- 13.* Tepals outside with curved eglandular hairs and strict glandular hairs. *A. ×acuminatum* (*A. degeneri* × *A. napellus*)
14. Cauline leaves with uppermost leaf segments only 1–3 (–4) mm broad; tepals light blue to whitish; hood ± triangular; seeds with transverse lamellae *sect. Angustifolium* ... *A. angustifolium*
 - a. Pedicels and tepals outside glabrous *fo. angustifolium*
 - a.* Pedicels and tepals outside ± pubescent .. *fo. carniolicum*
- 14.* Cauline leaves mostly with broader leaf segments; tepals mostly blue to violet; hood ± hemispherical; seeds smooth to transverse wrinkled *sect. Aconitum* ... 15
15. Stem at least in the inflorescence region flexuous (zigzag); inflorescence axis, pedicels and tepals outside glandular pilose *subsect. Burnatii* ... 16
- 15.* Stem stiff, upright; inflorescence axis, pedicels and tepals outside with a different pubescence *subsect. Aconitum* ... 19
16. Hood about as broad as high *A. maninense*
- 16.* Hood distinctly broader than high 17
17. Stem in the lower third glabrous; caudine leaves almost glabrous; nectary spur globose *A. pentheri*

- 17.* Stem in the lower third pubescent or pilose; caudine leaves densely pilose; nectary spur hooky..... 18
18. Stem in the lower third pubescent; caudine leaves with narrow segments; pedicels straight; filaments glabrous or pilose..... *A. burnatii*
- 18.* Stem in the lower third pilose; caudine leaves with broad segments; pedicels bent; filaments always pilose *A. nevadense*
19. Nectary spur elongated and bent backward..... ser. *Castellana* ... *A. castellanum*
- 19.* Nectary spur not elongated and not bent backward 20
20. Nectary spur distinctly globose; tepals outside glabrous to densely pubescent..... ser. *Aconitum*...21
- 20.* Nectary acephalous; tepals outside glabrous or only sparsely pubescent..... ser. *Taurica*...32
21. Nectary claw erect or only slightly curved at the top..... *A. corsicum*
- 21.* Nectary claw distinctly curved..... 22
22. Cauline leaves net-veined..... 23
- 22.* Cauline leaves not net-veined 30
23. Tepals outside densely curved and crisped eglandular pubescent, plant above 2 m high..... *A. superbum*
- 23.* Tepals outside with a different pubescence or glabrous.. 24
24. Tepals outside densely curved and crisped eglandular pubescent and glandular pilose, bracteoles divided..... *A. firmum* subsp. *skerisorae*
- 24.* Tepals outside glabrous or hairy 25
25. Tepals outside and pedicels glabrous..... *A. firmum*
- a. Bracteoles all similar, undivided, lanceolate....subsp. *fissurae*
- a.* Bracteoles of lower flowers bigger and divided b
- b. Bracteoles of lower flowers in ± deeply divided subsp. *firmum*
- b.* Bracteoles of lower flowers only bigger or just little divided.....
....nothosubsp. *fussianum* (subsp. *firmum* × subsp. *fissurae*)
- 25.* Tepals outside and pedicels eglandular pubescent hairy
- a. Pedicels eglandular pubescent
..... *A. firmum* subsp. *moravicum*
- a.* Pedicels below the bracteoles glabrous
..... *A. firmum* subsp. *moravicum* × *A. variegatum* subsp. *variegatum*
- 25.** Tepals outside glabrous or hairy, pedicels glandular or glandular/eglandular pubescent 26
26. Pedicels glandular pilose only above bracteoles.....
. *A. ×mariae* nothosubsp. *mariae* (*A. firmum* subsp. *firmum* × *A. maninense*)
- 26.* Pedicel with a different pubescence..... 27
27. Pedicels curved and crisped eglandular and glandular pubescent, mainly above the bracteoles, tepals outside glabrous *A. ×nanum* (*A. bucovinense* × *A. firmum*)
- 27.* Pedicels with a different pubescence 28
28. Pedicels sparsely glandular pubescent only above the bracteoles..... *A. ×czarnohorense* (*A. firmum* × *A. ×nanum*)
- 28.* Pedicels with a different pubescence 29
29. Pedicels and tepals outside curved and crisped eglandular pubescent and glandular pilose *A. ×mariae* nothosubsp. *paxii* (*A. firmum* subsp. *moravicum* × *A. maninense*)
- 29.* Pedicels only sparsely curved and crisped eglandular pubescent and glandular pilose, mainly just above the bracteoles, tepals outside glabrous..... *A. ×zapalowiczii* (*A. firmum* × *A. ×mariae* nothosubsp. *paxii*)
30. Pedicels densely curved and crisped eglandular and glandular pubescent *A. bucovinense*
- a. Carpels glabrous or rarely pilose on the ventral side fo. *bucovinense*
- a.* Carpels pubescent fo. *orthotricha*
- 30.* Pedicels eglandular hairy 31
31. Cauline leaves with linear to narrow lanceolate segments, the uppermost 2–4 mm broad, longly acuminate; flowers mauve to mauvish-blue; hood 18–20 mm high from base... *A. anglicum*
- 31.* Cauline leaves with mostly broader segments, which are shortly acuminate; flowers dark blue to violet; hood 13–18 (–20) mm high from the base..... *A. napellus*
- a. Carpels densely pubescent; often with bulbils in the lower leaves subsp. *lobelii*
- a.* Carpels glabrous or just sparsely pubescent on the backside; never with bulbils b
- b. Carpels sparsely pubescent on the backside c
- b.* Carpels glabrous d
- c. Bracteoles 2–3 (–4) mm long nothosubsp. *polatschekii* (subsp. *formosum* × subsp. *lobelii*)
- c.* Bracteoles (3–) 5–8 (–20) mm long ...nothosubsp. *seitzii* (subsp. *lobelii* × subsp. *napellus*)

- d. Filaments of stamens toothed subsp. *lusitanicum*
- d.* Filaments of stamens not toothed..... e
- e. Bracteoles (3–) 5–8 (–20) mm long; carpels 3 (–4).....
..... subsp. *napellus*
- e.* Bracteoles shorter; carpels 2–3 f
- f. Bracteoles 1–2 (–3) mm long; carpels 2–3
..... subsp. *formosum*
- f.* Bracteoles (2–) 3 (–4) mm long; carpels (2–) 3 notho-
..... subsp. *hinterhuberi* (subsp. *formosum* × subsp. *napellus*)
- 32. Tepals outside always glabrous ser. *Taurica*
 - A. Stem rigid, bracteoles linear to lanceolate 2–5 (–7) mm,
carpels 3; tepals outsides always glabrous .. *A. tauricum*
 - a. Pedicels with strict and crisped, glandular or eglandu-
lar hairs above the bracteoles; carpels often sparsely
pubescent on the suture subsp. *latemarensense*
 - a.* Pedicels with only few hairs on the top or glabrous;
carpels mostly glabrous b
 - b. Pedicels with solitary hairs at the top; carpels mostly
glabrous nothosubsp. *hayekianum* (subsp.
latemarensense × subsp. *tauricum*)
 - b.* Pedicels glabrous; carpels glabrous....subsp. *tauricum*
 - I. Inflorescence with many, ± long side racemes
..... var. *eustachyum*
 - I.* Inflorescence without or only with few short
side racemes var. *tauricum*
 - i. Nectaries and filaments glabrous....fo. *tauricum*
 - i.* Nectaries and filaments ± densely pilose
..... fo. *taurericum*
 - A.* Stem weekly ramified, bracteoles triangular to lin-
ear, 1–2 (–3) mm long; carpels 2 (–3) *A. clusianum*
 - 32.* Tepals outside sparsely hairy.....33
 - 33. Bracteoles triangular to linear, 1–2 (–3) mm long; carpels
2 (–3) *A. plicatum*
 - a. Pedicels eglandular pubescent, carpels glabrous.....
..... subsp. *plicatum*
 - a.* Pedicels eglandular and/or glandular pilose; carpels pubes-
cent..... subsp. *sudeticum*
 - 33.* Bracteoles linear, lanceolate or spatulate, 3–5 (–8) mm
long; carpels 2–3.....34
 - 34. Bracteoles situated always shortly below the flower at the
top of the pedicel; carpels (2–) 3.....
..... *A. ×teppneri* (*A. napellus* × *A. tauricum*)
 - a. Carpels glabrous.....b
 - a.* Carpels sparsely pubescent on the backside c
 - b. Pedicels sparsely curved and crisped eglandular pubes-
centnothosubsp. *teppneri* (*A.*
napellus subsp. *napellus* × *A. tauricum* subsp. *tauricum*)
 - b.*Pedicels glandular and eglandular curved and crisped
pubescent and pilose..... nothosubsp. *kerneri* (*A.*
napellus subsp. *napellus* × *A. tauricum* subsp. *latemarensense*)
 - c. Pedicels sparsely curved and crisped eglandular pubes-
centnothosubsp. *goetzii* (*A.*
napellus subsp. *lobelii* × *A. tauricum* subsp. *tauricum*)
 - c.* Pedicels glandular and eglandular curved and crisped
pubescent and pilose..... nothosubsp. *haderlappii* (*A.*
napellus subsp. *lobelii* × *A. tauricum* subsp. *latemarensense*)
 - 34.* Bracteoles situated distant from the flower, below to or in
the middle of the pedicel; carpels 2–3.....
..... *A. ×bavaricum* (*A. napellus* × *A. plicatum*)
 - a. Carpels glabrous.....nothosubsp. *bavaricum* (*A. napellus*
..... subsp. *napellus* × *A. plicatum*)
 - a.* Carpels sparsely pubescent on the backside
.....nothosubsp. *lusenense* (*A. napellus* subsp. *lobelii* × *A.*
plicatum)
 - 35. Pedicels and tepals outside glabrous or eglandular pubes-
centser. *Variegata*...36
 - 35.* Pedicels and tepals outside glandular pilose or very rarely
glabrous (*A. degenii* subsp. *rhaeticum*, *A. toxicum* subsp.
bucegiense).....39
 - 36. Pedicels eglandular curved pubescent; carpels glabrous or
pubescent on the backside
 - a. bracteoles at the middle of pedicel or below
.....sect. *Euchylodea*...*A. carmichaelli*
 - a.* bracteoles in the upper part below flower.....37
 - 36.*Pedicels mostly glabrous; tepals outside always glabrous;
carpels glabrous or pilose on the suture.....38
 - 37. Pedicels and tepals outside eglandular curved pubescent;
carpels mostly pubescent on the backside *A. vitosanum*
 - 37.*Pedicels sparsely eglandular curved pubescent; tepals out-
side glabrous; carpels mostly glabrous
..... *A. ×aquinonare* (*A. variegatum* × *A. vitosanum*)
 - 38. Hood not much higher than broad; claws of the nectaries
strongly curved; spurs of the nectaries mostly reaching the
top of the hood; carpels 3, glabrous *A. vivantii*
 - 38.* Hood distinctly higher than broad; claws of the nectaries
upright; spurs of the nectaries not reaching the top of the
hood; carpels 3–5, glabrous or pilose on the suture.....
..... *A. variegatum*

- a. Carpels 3, glabrous..... subsp. *nasutum*
 a.* Carpels 3–5, pilose at least on the suture..... b
 b. Carpels 3, only sparsely pilose on the top of the suture .. nothosubsp. *podobnikianum* (subsp. *nasutum* × subsp. *variegatum*)
 b.* Carpels 3–5, densely pilose on the suture subsp. *variegatum*
 c. Bracteoles 2–3 (–5) mm long, linear to narrow lanceolate; carpels always 3..... var. *carniolicum*
 c.* Bracteoles (4) 5–8 (–25) mm long, spatulate, ovate or leaflike; carpels (3–) 5..... d
 d. Leaf sheaths underside pubescent; pedicels sparsely pubescent below the bracteoles; carpels pilose on the suture and often also on the backside var. *stiriacum*
 d.* Leaf sheaths underside glabrous; pedicels glabrous; carpels only pilose on the suture var. *variegatum*
 39. Tepals outside densely glandular pilose or very rarely glabrous (*A. degenerii* subsp. *rhaeticum*, *A. toxicum* subsp. *bucegiense*)... ser. *Toxicaria*... 40
 39.* Tepals outside totally or almost glabrous nothoser. *Toxigata*... 46
 40. Hood distinctly higher than broad; nectary claw upright; nectary spur not reaching the top of the hood 41
 40.* Hood about as high as broad or broader than high; nectary claw curved; nectary spur always reaching the top of the hood 42
 41. Pedicels and tepals outside with curved and s-formed glandular hairs..... *A. pilipes*
 41.* Pedicels and tepals outside with straight glandular hairs.... *A. lasiocarpum*
 a. The whole pedicel glandular pilose....subsp. *lasiocarpum*
 a.* Pedicel only above the bracteoles glandular pilose, below the bracteoles glabrous or curved or crisped eglandular pubescent subsp. *kotulae*
 42. Bracteoles broad ovate, with reticulate venation....*A. toxicum*
 a. Tepals outside glabrous or almost glabrous..... b
 a.* Tepals outside curved or straight glandular pilose c
 b. Pedicels glabrous subsp. *bucegiense*
 b.* Pedicels just sparsely straight glandular pilose, mainly above the bracteoles..... nothosubsp. *ungarianum* (subsp. *bucegiense* × subsp. *toxicum*)
 c. Pedicels and tepals outside only with straight glandular hairs subsp. *toxicum*
 c.* Pedicels and tepals outside also or only with curved glandular hairs..... d
 d. Pedicels and tepals outside with curved and straight glandular hairs..... nothosubsp. *nyaradyanum* (subsp. *crispulum* × subsp. *toxicum*)
 d.* Pedicels and tepals outside only with curved glandular hairs subsp. *crispulum*
 42.* Bracteoles linear to narrow ovate 43
 43. Pedicels and tepals outside with straight, curved and s-formed glandular hairs
 *A. × pilosiusculum* (*A. degenerii* × *A. pilipes*)
 43.* Pedicels and tepals outside only with straight glandular hairs 44
 44. Bracteoles narrow ovate or spatulate, with branching veins *A. × dragulescuanum* (*A. degenerii* × *A. toxicum*)
 a. Pedicels, bracteoles and tepals outside only with straight glandular hairs..... nothosubsp. *dragulescuanum* (*A. degenerii* subsp. *degenerii* × *A. toxicum* subsp. *toxicum*)
 a.* Pedicels, bracteoles and tepals outside with curved and straight glandular hairs..... nothosubsp. *grintescuanum* (*A. degenerii* subsp. *degenerii* × *A. toxicum* subsp. *crispulum*)
 44.* Bracteoles linear to lanceolate, without branching veins..... 45
 45. Nectary claw distinctly curved; nectary spur capitate or slightly recurved *A. degenerii*
 a. Tepals outside glabrous b
 a.* Tepals outside glandular pilose..... c
 b. Pedicels glabrous subsp. *rhaeticum*
 b.* Pedicels sparsely glandular pilose, mainly above the bracteoles..... nothosubsp. *lippertianum* (subsp. *paniculatum* × subsp. *rhaeticum*)
 c. Bracteoles linear, situated about in the middle of the pedicel..... subsp. *paniculatum*
 I. Nectaries and filaments glabrous var. *laxiflorum*
 I.* Nectaries glabrous or pilose; filaments ± densely pubescent var. *turrachense*
 c.* Bracteoles lanceolate to spatulate, situated at the top of the pedicel e
 e. Carpels 5, densely pilose subsp. *valesiacum*
 e.* Carpels 3–5, mostly glabrous f
 f. Pedicels below the bracteoles almost glabrous; carpels 3–5, mostly glabrous nothosubsp. *gandogerii* (subsp. *paniculatum* × subsp. *valesiacum*)

- f.* Pedicels below the bracteoles sparsely to densely glandular pilose; carpels 3, glabrous subsp. *degenii*
- II. Pedicels below the bracteoles only sparsely glandular pilose var. *intermedium*
- II.* Pedicels below the bracteoles densely glandular pilose.. var. *degenii*
- i. Hood ± hemispheric fo. *degenii*
 - i.* Hood cymbiform, spur of nectaries capitate..... fo. *craciunelense*
- 45.* Nectary claw almost upright and only very slightly curved; nectary spur elongated and half or almost totally recurved..... *A. ×gayeri* (*A. degenii* × *A. lasiocarpum*)
46. Hood distinctly higher than broad; nectary claw upright; nectary spur not reaching the top of the hood 47
- 46.* Hood about as high as broad or slightly higher; nectary claw curved; nectary spur always reaching the top of the hood 48
47. Pedicels sparsely crisped and s-formed glandular pilose, mainly above the bracteoles; carpels (3–) 5 mostly ± densely pilose *A. ×austriacum* (*A. pilipes* × *A. variegatum*)
- 47.* Pedicels sparsely straight glandular pilose, mainly above the bracteoles; carpels 3–5, pilose only on the suture *A. ×pawlowskii* (*A. lasiocarpum* × *A. variegatum*)
48. Bracteoles narrow, without reticulate veination 49
- 48.* Bracteoles broad spatulate to narrow ovate, with reticulate venation..... 50
49. Tepals outside sparsely glandular pilose; carpels 3–5..... *A. ×hebegynum* (*A. degenii* × *A. variegatum*)
- 49.* Tepals outside glandular pilose and eglandular pubescent; carpels 3..... *A. ×tuscheticum* (*A. degenii* × *A. vitosanum*)
50. Carpels glabrous or pilose on the suture..... *A. ×bartokianum* (*A. toxicum* × *A. variegatum*)
- a. Pedicels and tepals outside curved glandular pilose; carpels glabrous nothosubsp. *rapaicsianum* (*A. toxicum* subsp. *crispulum* × *A. variegatum* subsp. *nasutum*)
 - a.* Pedicels and tepals outside straight glandular pubescent b
 - b. Carpels glabrous..... nothosubsp. *bartokianum* (*A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *nasutum*)
 - b.* Carpels pilose on the suture..... nothosubsp. *sooanum* (*A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *variegatum*)
- 50.* Carpels entirely pubescent..... *A. lasiocarpum* × *A. toxicum*

TAXONOMIC TREATMENT

Aconitum L. sect. *Aconitum* subsect. *Aconitum* ser. *Aconitum* in Europe is represented by six species, four nothospecies, and one hybrid formula; ser. *Castellana* Rottensteiner – by single species; ser. *Taurica* Mucher ex Starm. – by two species; nothoser. *Acorica* Starm. (ser. *Aconitum* × ser. *Taurica*) – by one species and two nothospecies (Mitka et al. 2017).

Aconitum sect. *Aconitum* subsect. *Burnatii* Rottensteiner consists of four endemic species distributed in the Sierra Nevada, Maritime Alps, French Massif Central, Balkans, and Western Carpathians (Rottensteiner 2018).

Aconitum sect. *Cammarum* DC. ser. *Variegata* Steinberg ex Starm. in Europe comprises three species and one nothospecies; ser. *Toxica* (Rchb.) Mucher is represented by four species, three nothospecies, and one hybrid formula; nser. *Toxigata* Starm. (ser. *Toxica* × ser. *Variegata*) – by five nothospecies.

Aconitum sect. *Angustifolium* Rottensteiner consists of one species (Seitz 1969).

Aconitum nothosect. *Acomarum* Starm. (Aconitum sect. *Aconitum* × Aconitum sect. *Cammarum*) consists of seven nothospecies and four hybrid formulae (Starmüller 2001; Wacławska-Ćwiertnia and Mitka 2016).

1. – Sect. *Aconitum*

1A. – Subsect. *Aconitum*

Description

Hood falciform, hemispherical, i.e., above rostrum wide-convex, or rounded-conic, 1–1.5 times higher than wide; claws of the nectaries bent and reaching the top of the hood, spurs of the nectaries acephalous, capitate or slightly bent, reaching the top of hoods; carpels 2–3, inflorescence axis, pedicels and tepals outside eglandular, glandular/eglandular pubescent/pilose or glabrous; leaves divided into 3 or 5 (–7) lobes and dissected to the base, the uppermost leaf segment linear to ovate, sometimes with 1–2 teeth; nectary spur capitate; seeds without membranous lamellae or rugulose, black to black-brown, with one or three longitudinal wings, then one wing more developed than the other two; root fusiform.

1a. – Ser. *Aconitum*

Type species: *A. napellus* L.

Diagnostic characters: nectary spur distinctly globose; tepals outside glabrous to densely pubescent.

Aconitum anglicum Stapf, Curtis's Bot. Mag. 151: tab. 9088. 1926

Iconotype: Stapf, Curtis's Bot. Mag. 151: tab. 9088, 1926.

Synonyms: *A. napellus* auct., *A. napellus* L. subsp. *napellus* (sensu Seitz 1969)

Distribution: Southwest and West England and East Wales.

Diagnostic characters: tubers up to 9 cm, by 3 cm at the top; stem with fine curled hairs; leaves 5- to 3-partite, deeply laciniate with lobes linear to narrowly linear-lanceolate and acuminate, the uppermost 2–4 mm wide, almost glabrous; pedicels (1–) 2 (–3) cm long, quite erect; flowers mauve to mauvish-blue, minutely downy, curved eglandular pubescent; hood 18–20 mm high from the base; lower tepals strongly deflexed; lateral tepals 12–17 mm; claws of the nectaries curved almost horizontal, with a capitate spur; filaments hairy; carpels 3, glabrous.

Iconography: Stapf (1926); Ross-Craig (1948); Clapham et al. (1957).

Literature: Seitz (1969); Starmühler (1998).

Hybrids: unknown.

Aconitum bucovinense Zapał., Rozpr. Wydz. Mat.-Przyr. Akad. Umiej., Dział B. Nauki Biol. 48: 89–90. 1908a, *pro hybr.* [reprinted in Zapałowicz 1908b].

Type citation: "In montanis et subalpinis Bucovinae australis: Jakobeny, Pojana niegri prope Dorna Kandreny (*Rehman*)".

Typus: lectotype or, perhaps, holotype: KRAM 132396! (Wacławska-Ćwiertnia and Mitka 2016).

Synonyms: *A. callibotrys* Rchb. subsp. *bucovinense* (Zapał.) Grinč., *A. callibotrys* Rchb. subsp. *rigidum* (Rchb.) Grinč., *A. firmum* Rchb. subsp. *bucovinense* (Zapał.) Aschers. & Graebn.

Distribution: Eastern Carpathians, Southern Carpathians.

Diagnostic characters: the uppermost leaf segment 3–5 mm wide; hood and pedicels densely curved and crisped eglandular pubescent and often glandular pilose; claws of the nectaries curved, glabrous or pilose; bracteoles linear to obovate-lanceolate; filaments and stamens pilose; carpels glabrous or, rarely, pilose on the dorsal side (fo. *bucovinense*) or pubescent (fo. *orthotricha*).

Iconography: Mitka (2003: 179, fig. 12A–C); Novikoff and Mitka (2011: 49, fig. 5F,G).

Literature: Starmühler (1998); Starmühler (2000); Mitka (2003); Starmühler (2010); Wacławska-Ćwiertnia and Mitka (2016).

Lower taxa:

a) fo. ***bucovinense***

Diagnostic characters: carpels glabrous or, rarely, pilose on the dorsal side.

b) fo. ***orthotricha*** Gáyer, Magyar Bot. Lap. 9: 168. 1909

Type citation: did not provided by the author.

Typus: not designated.

Synonyms: none.

Diagnostic characters: carpels pubescent.

Hybrids:

a) ***Aconitum ×nanum*** (Baumg.) Simonk., Enum. Fl. Transsilv.: 64. 1887

Type citation: "In mtb. Csibesz et Arszuluj".

Typus: lectotype, (Romania) in m. Arszuluj (1812) – CL-Baumgarten 4745!

Synonyms: *A. napellus* L. var. *nanum* Baumg., *A. tauricum* Wulfen subsp. *nanum* (Baumg.) Gáyer, *A. tauricum* Wulfen subsp. *nanum* (Baumg.) Grinč.

Hybrid origin: *A. bucovinense* × *A. firmum*.

Diagnostic characters: hoods glabrous, pedicels eglandular and glandular pubescent.

b) ***Aconitum ×czarnohorense*** (Zapał.) Mitka, The genus *Aconitum* in Poland and adjacent countries: 77. 2003

Type citation: "W Górzach Pokucko-Marmaroskich, szczególnie na Czarnej Horze i w Alpach Rodneńskich w kraju kosodrzewu często, najwyżej tam 2010–2030 m; liczne okazy z Czarnej Hory i z Pietrosu w Alpach Rodneńskich".

Typus: lectotype, Góry Pokucko-Marmaroskie, Pietrosz, 8.08.1880, leg. et det. H. Zapałowicz, KRAM 246853! (Wacławska-Ćwiertnia and Mitka 2016).

Synonyms: *A. napellus* L. em Skalický var. *czarnohorense* Zapał. fo. *czarnohorense*, *A. napellus* L. var. *czarnohorense* Zapał. fo. *amoenum* Zapał., *A. napellus* L. var. *czarnohorense* Zapał. fo. *glabratum* Zapał., *A. napellus* L. var. *czarnohorense* Zapał. fo. *hooverianum* Zapał., *A. napellus* L. var. *czarnohorense* Zapał. fo. *nanum* Zapał. non Baumg., *A. napellus* L. em Skalický var. *czarnohorense* Zapał. fo. *rodnense* Zapał., *A. napellus* L. var. *czarnohorense* Zapał. fo. *tenuiseptum* Zapał., *A. napellus* L. em Skalický var. *czarnohorense* Zapał. fo. *turkulense* Zapał.

Hybrid origin: *A. bucovinense* × *A. ×nanum*.

Diagnostic characters: hoods glabrous, pedicels below bracteoles glandular pilose or glabrous.

Iconography: (Mitka 2003: 179, fig. 12E); Novikoff and Mitka 2011: 49. fig. 5K).

Aconitum corsicum Gáyer, Magyar Bot. Lap. 8: 181. 1909

Type citation: "Corsica: M. Coscione (Kralik – W; Reverchon – D)".

Typus: paratype, WU 0034394.

Synonyms: *A. lobelianum* Rouy-Fouc., *A. napellus* L. var. *compac-*

tum Rapaics., *A. napellus* L. subsp. *corsicum* (Gáyer) W. Seitz, *A. pyramidale* Mill. var. *corsicum* (Gáyer) P. Fourn.
Distribution: Corsica.

Diagnostic characters: hood and pedicels densely pubescent, inflorescence rigid, in lower part ramified; claws of the nectaries erected, at theirs top slightly bent, spur of the nectaries capitate or acephalous, reaching the top of the hoods; filaments pilose; bracteoles linear to linear-lanceolate 3–5 mm; carpels glabrous; the uppermost leaf segment 2–4 mm wide.

Iconography: Jeanmonod and Gamisans (2007).

Literature: Gáyer (1909); Starmühler (1998).

Hybrids: unknown.

Aconitum firmum Rchb., Uebers. Aconitum: 20. 1819

Type citation: “in Carpathorum tractu Townson!, Genersich!, Kitaibel!, Wahlenberg!, Sadler! in Transylvania Baumgarten! in Carinth. Alp. Fladnitz. v. Vest!”

Typus: iconotype, “Rchb., Mon. Acon.: 85, Tab. XIV, fig. 1, 1821” (Skalicky 1990).

Synonyms: *A. flerovii* Steinb. in Komarov, *A. palmatifidum* Rchb., p.p., *A. romanicum* Wołoszczak, *A. tatrae* Borb., *A. koelleanum* Rchb. var. *firmum* (Rchb.) Rchb., *A. napellus* L. var. *firmum* (Rchb.) Pawł., *A. napellus* L. var. *babiogorense* Zapáč. fo. *babiogorense*, *A. napellus* L. var. *babiogorense* Zapáč. fo. *subfissum* Zapáč., *A. napellus* L. var. *carpaticum* Zapáč. fo. *carpaticum*, *A. napellus* L. var. *subtatrense* Zapáč. fo. *subtatrense*, *A. napellus* L. var. *subtatrense* Zapáč. fo. *abnorme* Zapáč., *A. napellus* L. var. *subtatrense* Zapáč. fo. *latisectum* Zapáč., *A. napellus* L. var. *tatrense* Zapáč., *A. skerisorae* auct., non Gáyer, *A. tauricum* auct. Fl. Carpathorum, non Wulfen.

Distribution: Carpathian Mts, Apuseni Mts, Balkans, Central Russian Highlands.

Diagnostic characters: bracteoles (2.5–) 3–17 mm long, at least those of the lowest flowers in terminal raceme divided or at least toothed; pedicels and tepals outside glabrous (subsp. *firmum*, subsp. *fissurae*, and nothosubsp. *fussianum*), curved and crisped eglandular pubescent and glandular pilose (subsp. *skerisorae*), or eglandular pubescent (subsp. *moravicum*); bracteoles 2–5 (–7) mm long, wether toothed nor divided; stamens glabrous (subsp. *firmum*) or villose (subsp. *fissurae* and nothosubsp. *fussianum*).

Iconography: Bauhin (1671: 183); Reichenbach (1821: t. xiv, fig. 1); Mitka (2003: 178); Novikoff and Mitka (2011: 49).

Literature: Skalicky (1990); Starmühler (1998); Starmühler (2000); Starmühler and Mitka (2001); Mitka (2003); Starmühler (2010); Novikoff and Mitka (2011); Mitka et al. (2017).

Lower taxa:

a) subsp. *firmum*

Diagnostic characters: pedicels and tepals outside glabrous; bracteoles ± deeply divided; stamens glabrous.

b) subsp. *fissurae* Nyár., Enum. Pl. Cheia Turzii: 132. 1939

Type citation: “[Romania] Transsilvania, Cheia Turzii, Repezieul caprelor; Nr. 435214; 06.09.1936; leg. E.I. Nyárády (CL)” (Starmühler and Mitka 2001).

Typus: lectotype, CL 435214.

Synonyms: *A. romanicum* Wołoszczak, *A. flerovii* Steinberg in Komarov.

Diagnostic characters: pedicels and tepals outside glabrous; bracteoles entire or slightly toothed; stamens villose.

c) subsp. *skerisorae* (Gáyer) Starm., Siebenbürg. Arch. 36: 18. 2000

Type citation: “Plantae Hungariae exsiccatae, Comit. Kolozs, in lapidosis fl. Melegszamos ad pag. Melegszamos, Nr. 60705; 17.07.1904 leg. Z. Zsák (BP); *A. skerisorae* Gáy., rev. Gáyer 1910” (Starmühler 2000).

Typus: neotype, BP 60705.

Synonyms: *A. skerisorae* Gáyer, *A. tatrae* Borbás var. *skerisorae* Soó, *A. napellus* Rchb. subsp. *skerisorae* Seitz non Gáyer (Soó 1972).

Diagnostic characters: pedicels and tepals outside eglandular pubescent and glandular pilose; bracteoles from lanceolate to divided; stamens villose or rarely glabrous.

d) subsp. *moravicum* Skalický, Preslia 54 (2): 115 1982

Type citation: “[Czech Republic] Beskydy, u cesty na západ, svahu Smrku, 1025 m, Nr. 322; 27.07.1943; leg. R. Kurka (PCR)” (Starmühler and Mitka 2001).

Typus: holotype, PCR 322.

Synonyms: *A. napellus* L. subsp. *firmum* (Rchb.) Gáyer var. *carpaticum* Maloch.

Diagnostic characters: pedicels and tepals outside eglandular pubescent; bracteoles at least those of the lowest flowers divided or at least toothed, rarely undivided; stamens mostly pilose.

e) nothosubsp. *fussianum* Starm. (*A. firmum* subsp. *firmum* × *A. firmum* subsp. *fissurae*) – see hybrids of *A. firmum*.

Hybrids:

a) *Aconitum firmum* nothosubsp. *fussianum* Starm., Siebenbürg. Arch. 36: 17. 2000

Type citation: "Transilvania (Siebenbürgen, Erdély), Muntii Apuseni (Siebenbürgisches Westgebirge, Nyugati-havasok), Muntii Vladeasa (Vlegyásza-hegyseg), W Huedin (Bánffyhunyad), im Valea Draganului (Nagysobos) beim Dorf Lunca, 735 m". Typus: holotype, GZU 000225133.

Synonyms: none.

Hybrid origin: *A. firmum* subsp. *firmum* × *A. firmum* subsp. *fissurae*.

Diagnostic characters: tepals outside glabrous; bracteoles toothed or lobed; stamens villose.

b) *Aconitum ×nanum* (Baumg.) Simonk. (*A. bucovinense* × *A. firmum*) – see *A. bucovinense*.

c) *Aconitum ×czarnohorense* (Zapał.) Mitka (*A. firmum* × *A. ×nanum*) – see *A. bucovinense*.

d) *Aconitum ×mariae* Rottensteiner, Mitka & Novikov, nothospec. (hybr.) nov.

Typus: [Slovakia] Hungaria, comitatus Szepes, Montes Bélaenses, in valle Drechslerhäuschen sub monte Stirnberg, alt. cca. 1400–1500 m, solo calc.; 01.09.1907; leg. E.G. Nyárády (holotype, SIB).

Diagnosis

A. ×mariae differs from *A. firmum* by glandular pilose pedicels and differs from *A. maninense* by glabrous tepals.

Description

Inflorescence axis glabrous, pedicels glandular pubescent and pilose, often only above the bracteoles, bracteoles linear, lanceolate, spathulate or even divided, situated in the upper half of the pedicel, tepals outside glabrous, filaments of the stamens glabrous or sparsely pilose, carpels glabrous or sparsely pilose on the backside.

Hybrid origin: *A. firmum* s.str. × *A. maninense*.

Eponymy

This new *Aconitum* hybrid is named after a distinguished botanist, our friend and colleague from the Institute of Botany of the Jagiellonian University in Kraków, Professor Maria Zająć (1955–2018).

e) *Aconitum ×mariae* Rottensteiner, Mitka & Novikov nothosubsp. *paxii* (Starm.) Mitka, comb. nov.

Basionym: *A. firmum* Rchb. nothosubsp. *paxii* Starm. in Starmühler & Mitka, Thaiszia 10 (2): 127, 2001.

Type citation: "Polen, Galizien, Nord-Karpaten, Hohe Tatra (Tatry Wysokie), S-Ufer vom Großen Fischsee (Morskie Oko), N 49°11.64', E 20°04.12', 1395 m alt.; Hochstaudenflur; 13.08.1997; leg. U. et W. Starmühler" (Starmühler and Mitka 2001).

leg. U. et W. Starmühler" (Starmühler and Mitka 2001).

Typus: holotype, GZU 000232750.

Synonyms: *A. palmatifidum* Rchb. fo. *piliferum* Gáyer.

Hybrid origin: *A. firmum* subsp. *moravicum* × *A. maninense*.

Diagnostic characters: pedicels curved and crisped eglandular pubescent and glandular pilose.

Iconography: Mitka (2003: 178, fig. 11E).

f) *Aconitum ×zapalowiczii* (Starm.) Mitka, Studia Universitatis Babeş-Bolyai, Biologia, 62: 166. 2017

Basionym: *A. firmum* Rchb. nothosubsp. *zapalowiczii* Starm. in Starmühler & Mitka, Thaiszia 10 (2): 128, 2001.

Type citation: "Polen, Galizien, Nord-Karpaten, Hohe Tatra (Tatry Wysokie), S-Ufer vom Großen Fischsee (Morskie Oko), N 49°11.64', E 20°04.12', 1395 m alt.; Hochstaudenflur; 13.08.1997; leg. U. et W. Starmühler" (Starmühler and Mitka 2001).

Typus: holotype, GZU 000232751.

Synonyms: none.

Hybrid origin: *A. firmum* × *A. ×mariae* nothosubsp. *paxii*.

Diagnostic characters: pedicels only sparsely curved and crisped eglandular pubescent and glandular pilose, mainly just above the bracteoles.

Iconography: Mitka (2003: 178, fig. 11F).

Aconitum napellus L., Sp. Pl. ed. 1, 1: 352. 1753

Type citation: "in Helvetia, Bavaria, Gallia".

Typus: lectotype in Herb. Clifford: 214, A. 3, sheet A, BM 000628795 (Seitz 1969).

Synonyms: *A. autumnale* Clus. ex Rchb., *A. bernhardianum* Rchb., p.p., *A. compactum* Rchb., *A. eminens* Koch. ex Rchb., *A. laxum* Rchb., *A. linnaeanum* Gáyer, *A. lobelianum* Rchb., *A. meyeri* Rchb., *A. neomontanum* Wulfen, *A. neubergense* Clus. ex Rchb., *A. pyramidale* Mill. ex Rchb., *A. strictum* Bernh., *A. vulgare* DC., *Delphinium napellus* (L.) Baill.

Distribution: Alps, Pyrenees, Iberian Peninsula, Bohemian Massif.

Diagnostic characters: inflorescence axis, pedicels and hoods outside pubescent; filaments not toothed; nectaries bent horizontal, spur of nectaries capitate; inflorescence rigid or loose/ramified; bracteoles (3–) 5–8 (–20) mm long and carpels (2–) 3, densely pubescent (subsp. *lobelii*) or carpels sparsely pubescent on the backside only (nothosubsp. *seitzii*); carpels glabrous, bracteoles linear to lanceolate, rarely spathulate (4–) 5–8 (–15) mm (subsp. *napellus*), 1–2 (–3) mm long (subsp. *formosum*); or nectaries hardly bent; filaments ± toothed, carpels 2–3 (subsp. *lusitanicum*); bracteoles (2–) 3 (–4) mm long and carpels (2–) 3 glabrous (nothosubsp. *hinterhuberi*) or carpels sparsely pubescent on the backside (nothosubsp. *polatschekii*).

Iconography: Matt., New Kreuterb.: 472, fig. D, 1563; Clus. Rar. Pl. Hist. 2: 96 *pro Aconit. Lycoc. vi. Napellus vulgar.*, 1601; Jacq. Fl. Austr. v. 4, t. 381, 1776.

Literature: Cappelletti and Poldini (1984); Molero and Blanché (1986); Mucher (1991a); Starmühler (1999); Starmühler (2001); Starmühler (2002).

Lower taxa:

a) subsp. *napellus*

Diagnostic characters: carpels 3 (–4), glabrous; bracteoles linear to lanceolate, rarely spatulate (4–) 5–8 (–15) mm; filaments not toothed.

b) subsp. *lobelii* Mucher, Phyton (Horn) 31(1): 130. 1991

Type citation: "Ostalpen, Steiermark, Salzkammergut, Totes Gebirge, Loser, Panoramastraße, Hochstaudenflur, 1020 m, 9.8.1989, leg. W. Mucher, GZU" (Mucher 1991c).

Typus: holotype, GZU 000274111. Isotypes, GJO, GZU 000274112 – GZU 000274117, W, WU.

Synonyms: *A. lobelianum* Host, p.p., non Rchb., *A. lobelianum* (Rchb.) Host. in Gáyer, p.p.

Diagnostic characters: carpels (2–) 3, densely pubescent; bracteoles linear to lanceolate, (3–) 5–8 (–20) mm; filaments not toothed, pilose.

c) subsp. *formosum* (Rchb.) Gáyer in Hegi, Fl. Mitteleur. 3: 498. 1912

Type citation: "Salzburger Alpen (Untersberg, Schafberg)" (Gáyer 1912: 498)

Typus: iconotype, "Reichenbach 1825, Ill. Spec. Acon. Gen., icon. 65" (Starmühler 1997b).

Synonyms: *A. formosum* Rchb., *A. napellus* L. [var.] *formosum* (Rchb.) Koch, *A. napellus* [var.] β *hemisphaericum* fo. *formosum* (Rchb.) Beck.

Iconography: Gáyer (1912: 499, fig. 655 a–c).

Diagnostic characters: carpels 2–3, glabrous; bracteoles linear to lanceolate, only 1–2 (–3) mm long; filaments not toothed.

d) subsp. *lusitanicum* Rouy, Le Naturaliste 6: 405. 1884

Type citation: "Prov. de Tras-os-montes: environ de Bracançà, à San Martin ho d'Augueira. – Sept. 1854. – leg. E. Schmitz" (Rouy 1884).

Typus: not designated.

Synonyms: none.

Diagnostic characters: carpels 2–3, glabrous; filaments toothed.

e) nothosubsp. *hinterhuberi* (*A. napellus* subsp. *formo-*

sum \times *A. napellus* subsp. *napellus*) – see hybrids of *A. napellus*.

f) nothosubsp. *polatschekii* (*A. napellus* subsp. *formosum* \times *A. napellus* subsp. *lobelii*) – see hybrids of *A. napellus*.

Hybrids:

a) *Aconitum napellus* L. nothosubsp. *hinterhuberi* Starm., Fritschiana 18: 12. 1999

Type citation: "*Aconitum brauneanum*, Untersberg, 1825, leg. Hinterhuber (WU-Keck). Dieser Beleg wurde bereits von Seitz 1966 als "*A. napellus* ssp. *hians* gegen ssp. *neomontanum*" revidiert" (Starmühler 1999).

Typus: holotype, WU-Keck.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *formosum* \times *A. napellus* subsp. *napellus*.

Diagnostic characters: carpels (2–) 3, glabrous; bracteoles (2–) 3 (–4) mm long; filaments of stamens not toothed.

b) *Aconitum napellus* L. nothosubsp. *polatschekii* Mucher ex Starm., Fritschiana 18: 13. 1999

Type citation: "Ostalpen, Steiermark, Mariazellerland, Südufer des Hubertussees an der niederösterreichischen Grenze, feuchte Wiese, 840 m; 04.08. 1989; leg. W. Mucher" (Starmühler 1999).

Typus: holotype, GZU 000274119. Isotype, W.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *formosum* \times *A. napellus* subsp. *lobelii*.

Diagnostic characters: carpels (2–) 3 sparsely pubescent on the backside; bracteoles (2–) 3 (–4) mm long.

c) *Aconitum napellus* L. nothosubsp. *seitzii* Mucher ex Starm., Fritschiana 18: 14. 1999

Type citation: "Ostalpen, Steiermark, Mariazellerland, in der Walster, Bachufer beim Kaiser Franz Josef Denkmal 810 m; 04.08.1989; leg. W. Mucher" (Starmühler 1999).

Typus: holotype, GZU 000274118. Isotype, W.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *lobelii* \times *A. napellus* subsp. *napellus*.

Diagnostic characters: carpels (2–) 3 sparsely pubescent on the backside; bracteoles (3–) 5–8 (–20) mm long.

Aconitum superbum Fritsch, Verh. K.K. Zool.-Bot. Ges. Wien 45: 370. 1895

Type citation: "Mala Vrata unweit vom Berge Koprivnica zwis-

chen Bugojno und Kupres/Bosnien (Sostarič, Aug. 1895)".
 Typus: holotype, WU 2540.
 Synonyms: *A. adriaticum* Gáyer, *A. sostaricianum* Fritsch, *A. napellus* L. subsp. *superbum* (Fritsch) W. Seitz
 Distribution: Western and Central Balkans.

Diagnostic characters: plant 200–300 cm high; inflorescence axis, pedicels and hoods outside eglandular pubescent; leaves on the lower side with distinct reticulate venation, the uppermost leaf segment ovate; filaments pilose; carpels 2 (–3) glabrous.

Iconography: Seitz (1969: 16).
 Literature: Fritsch (1895); Gáyer (1909).
 Hybrids:

- a) *A. superbum* × *A. variegatum* – more studies required.
- b) *A. superbum* × *A. vitosanum* – more studies required.

1b. – Ser. *Castellana* Rottensteiner, series nova

Type species: *A. castellanum* (Molero & Blanché)
 Rottensteiner.

Diagnosis

Spurs of nectaries are elongated and backward bent.

Description

Monotypical series with only one species, separated from the other series of subsection *Aconitum* by the unique character of the elongated and backward bent spur of the nectaries.

Aconitum castellanum (Molero & Blanché) Rottensteiner, stat. nov.

Basionym: *A. napellus* L. subsp. *castellanum* Molero and Blanché, Anal. Jardin Bot. Madrid 41: 213. 1984.
 Type citation: "Laguna del Marquesado, 30S XK1349, loco humido umbraque – in populeto – ad 1400 m, J. Molero et A. Rovira die 20-VIII-1983 legerunt".
 Typus: holotype, BCF 71554.
 Distribution: Iberian Peninsula.

Diagnostic characters: stem and leaves glabrous or glabrescent; inflorescence ramified, curved eglandular pubescent; hood hemispheric-falciform; nectaries hardly curved, spur of the nectaries elongated and bent backward; filaments not dentate, pilose; carpels 3–5, glabrous.

Iconography: Molero and Blanché (1986: 238).
 Literature: Molero and Blanché (1984), Molero and Puig (1990).

Hybrids: unknown.

1c. – Ser. *Taurica* Mucher ex Starm.

Type species: *A. tauricum* Wulfen

Diagnostic characters: inflorescence simple or shortly ramosed; bracteoles triangle to linear, glabrous or at margins ciliate; hood and pedicels glabrous; nectary spur acephalous or hardly capitate.

Aconitum clusianum Rchb., Mon. Aconit.: 91. 1821

Type citation: "Bohemian Sudetes: Aupagrund".
 Typus: iconotype: Reichenbach, Mon. Aconit. Tab. III, 1. 1821
 Distribution: Sudetes, Bohemian Massif.

Diagnostic characters: plant 30–150 cm high, inflorescence weakly ramified, hood falciform or rounded-conic, pedicels glabrous; carpels 2 (–3) glabrous; spur of nectaries acephalous or capitate; bracteoles triangle to linear 1–2 (–3) mm long; filaments pilose; the uppermost leaf segment ovate or lanceolate-ovate 3–5 mm wide.

Iconography: Reichenbach (1821: tab. XIII, fig. 1); Mitka (2003, 177: fig. 10D).
 Literature: Mitka (2003).
 Hybrids: unknown.

Aconitum tauricum Wulfen in Koelle, Spicil. Observ. Acon.: 15. 1788

Type citation: "Frequentissimum in apricis Taureri Malnizesis, Radstadienis, etc."
 Typus: iconotype, "Typus gesehen von Gáyer, Magyar Bot. Lap. P. 146 (1909): (W) – ist vernichtet; Abbildung bei Reichenbach, illustr. Aconit. T. 63 (1827)" (Seitz 1969).
 Synonyms: *A. dolomiticum* A.Kern.; *A. hayekianum* Gáyer; *A. koelleanum* Rchb., p.p.; *A. napellus* L. subsp. *koelleanum* (Rchb.) Mucher; *A. parviflorum* Host.; *A. taurericum* Rchb.; *A. trichocarpum* Rchb., *A. napellus* L. subsp. *tauricum* (Wulfen) Gáyer in Hegi, *A. napellus* L. var. *tauricum* (Wulfen) Sér., *A. napellus* fo. *koelleanum* (Rchb.) Gürke in Richter-Gürke.
 Distribution: Alps.

Diagnostic characters: plant 60–80 cm high; inflorescence rigid; hoods falciform or hemispherical; inflorescence axis and tepals outside glabrous, pedicels glabrous (subsp. *tauricum*), with solitary hairs at the top (nothosubsp. *hayekianum*) or pilose above bracteoles (subsp. *latemarensis*); inflorescence with a lot of long side racemes (var. *eustachyum*) or without or only a few short side racemes (var. *tauricum*), filaments glabrous (fo. *tauricum*) or pilose (fo. *taurericum*); spur of nectaries acephalous or hardly capitate; the uppermost leaf segments 1–3 mm wide.

Iconography: Clusii Histor. Rarior. Plant p. XCV, n. IV, 1601; Reichenbach (1827: t. 63).

Literature: Mucher (1993a); Starmüller (2001).

Lower taxa:

a) subsp. *tauricum* var. *tauricum* fo. *tauricum*

Diagnostic characters: inflorescence compact, without or with only a few short side racemes; pedicels, carpels, nectaries and filaments glabrous.

b) subsp. *tauricum* var. *tauricum* fo. *taurericum* (Rchb.) Gáyer, Magyar Bot. Lap. 8: 145. 1909

Type citation: none.

Typus: not designated, probably iconotype is Rchb., Monogr. Acon.: 87, tab. 12, figs. 2, 3, 1820.

Synonyms: *A. taurericum* Rchb., *A. tauricum* Wulfen in Koelle subsp. *tauericum* (Rchb.) Grinč.

Diagnostic characters: inflorescence compact, without or with only a few short side racemes; pedicels, carpels, nectaries and filaments pilose.

c) subsp. *tauricum* var. *eustachyum* (Rchb.) Starm., Hladnikia 6: 42. 1996

Type citation: none.

Typus: not designated, probably iconotype is Monogr. Aconit.: 96, tab. 15, fig. 3, 1820.

Synonyms: *A. eustachyum* Rchb., *A. napellus* L. subsp. *eustachyum* (Rchb.) Gáyer in Hegi, *A. napellus* L. [var.] γ *eustachyum* (Rchb.) Fiori.

Diagnostic characters: inflorescence ramified; pedicels with solitary hairs or glabrous; carpels mostly glabrous.

d) subsp. *latemarense* (Degen & Gáyer) Starm. in W.Maurer, Fl. Steiermark 1: 84. 1996

Type citation: "Tirol: in lapidosis subalpinis ad pedem montis Latemar prope Karersee, alt. ca.1600–1800 m, 30.VIII.1906, Degen".

Typus: not designated.

Synonyms: *A. latemarense* Degen & Gáyer, *A. tauricum* Wulfen in Koelle var. *latemarense* (Degen & Gáyer) Mucher.

Diagnostic characters: pedicels with mixed glandular and eglandular pubescence; carpels often sparsely pubescent on the suture.

e) nothosubsp. *hayekianum* (*A. tauricum* subsp. *latemarense* × *A. tauricum* subsp. *tauricum*) – see hybrids of *A. tauricum*.

Hybrids:

a) *Aconitum ×acutum* Rchb. (*A. tauricum* × *A. variegatum*) – see nsect. *Acomarum*.

b) *Aconitum ×mielichhoferi* Rchb. (*A. degenii* × *A. tauricum*) – see nsect. *Acomarum*.

c) *Aconitum tauricum* Wulfen in Koelle nothosubsp. *hayekianum* (Gáyer) Grinč. in Savulescu, Flora Reipublicii Populare Române, 2: 475. 1953

Type citation: "Sannotaler Alpen" (Gáyer 1911)

Type specimens: not designated.

Synonyms: *A. dolomiticum* Hayek, p.p., *A. ranunculifolium* Rchb. var. *dolomiticum* Evers, p.p.

Hybrid origin: *A. tauricum* subsp. *latemarense* × *A. tauricum* subsp. *tauricum*.

Diagnostic characters: pedicels with solitary hairs at the top; carpels mostly glabrous.

d) *Aconitum pilipes* × *A. tauricum* – see nothosect. *Acomarum*.

1d. – Nothoser. *Acorica* Mucher ex Starm. (*Aconitum* ser. *Aconitum* × *Aconitum* ser. *Taurica*)

Type species: *A. plicatum* Rchb., designated here.

Diagnostic characters: spur of nectaries acephalous or hardly capitate, hoods pilose.

Aconitum plicatum Köhler ex Rchb., Uebers. Aconitum: 29. 1819

Type citation: "In Sudetis Köehler!".

Typus: iconotype in Reichenbach, Icon. Fl. Germ. Helv., Vol. 4: Icon. T. 98, Nr. 4708d. 1840 (Starmüller 1997a).

Synonyms: *A. amoenum* Rchb., *A. bernhardianum* Rchb., p.p., *A. callibotryon* Rchb., *A. hians* Rchb., *A. koehleri* Rchb., *A. laetum* Rchb., *A. multifidum* Koch ex Rchb., *A. rigidum* Rchb.

Distribution: Sudetes, Bohemian Massif, Bavarian Alps.

Diagnostic characters: plant 30–150 cm high, inflorescence weakly ramified, hood falciform or rounded-conic, carpels glabrous, pedicels eglandular pubescent (subsp. *plicatum*) or glandular/eglandular pilose (subsp. *sudeticum*); carpels 2 (–3) glabrous; spur of nectaries acephalous or capitate; bracteoles triangle to linear 1–2 (–3) mm long; filaments pilose; uppermost leaf segment ovate or lanceolate-ovate 3–5 mm wide.

Iconography: Mattioli (1563: 472, fig. D); Clusius (1601: 96, pro *Aconit. Lycoc. vi. Napellus vulgar.*); Jacquin (1776: t. 381); Reichenbach (1840: 48, fig. 4708 d); Mitka (2003: 177).

Literature: Cappelletti and Poldini (1984); Molero and Blanché

(1986); Mucher (1991a); Starmühler (1999); Starmühler (2001); Starmühler (2002); Mitka (2003).

Lower taxa:

a) subsp. *plicatum*

Diagnostic characters: pedicels eglandular pubescent; carpels glabrous.

b) subsp. *sudeticum* Mitka, The genus *Aconitum* in Poland and adjacent countries: 76. 2003.

Type citation: "Sudety Wschodnie, Masyw Śnieżnika, nad potokiem Czarna, 960 m, pod Dzikim Stokiem, 10.8.1988, leg. Z. Szeląg" (Mitka 2003).

Typus: holotype, KRAM. Isotypes, KRAM. Paratypus, KRA. Synonyms: none.

Diagnostic characters: pedicels eglandular and/or glandular pilose; carpels pubescent.

Hybrids:

a) *Aconitum ×bavaricum* Starm., Ber. Bayer. Bot. Ges. 71: 105. 2001, nothosubsp. *bavaricum*

Type citation: "Aconita rarissima selecta (2001), Deutschland, Bayern, Bayerischer Wald, NNE Bodenmais, Großer Arber, E-Hang unter dem Gipfel, über der Bodenmais-Mulde, N 49°06.66', E 13°08.08', 1435 m alt.; 31.08.2000; leg. W. Starmühler (GZU)" (Starmühler 2001).

Typus: holotype, GZU 000232791. Isotype, M, Herb. STARMÜHLER.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *napellus* × *A. plicatum*.

Diagnostic characters: bracteoles situated distant from the flower, carpels glabrous.

b) *Aconitum ×bavaricum* Starm. nothosubsp. *lusenense* Starm., Ber. Bayer. Bot. Ges. 71: 105. 2001

Type citation: "Ranunculaceae subtrib. Delphiniinae exsiccatae, Fasc. 5 (2001), Deutschland, Bayern, Bayerischer Wald, NE Grafenau, Lusen, am Weg vom Teufelsjoch zur Böhmküche, 1045 m alt.; Quellufer; 29.08.2000; leg. W. Starmühler (GZU)" (Starmühler 2001).

Typus: holotype, GZU 000232817. Isotype, GJO, JACA, KRA, LE, LG, LI, M, MEL, NY, PE, Herb. STARMÜHLER, TBI, TNS, W, Z.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *lobelii* × *A. plicatum*.

Diagnostic characters: carpels sparsely pubescent on the backside.

c) *Aconitum ×teppneri* Mucher ex Starm., Ber. Bayer. Bot. Ges. 71: 107. 2001, nothosubsp. *teppneri*

Type citation: "Ostalpen, Steiermark, Hochschwab, Schönleiten, Latschengebüsch, 1680 m, 22.8.1989, leg. W. Mucher (GZU)" (Starmühler 2001).

Typus: holotype, GZU 000274190. Isotypes, GZU 000274195, GZU 000274196, W, WU.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *napellus* × *A. tauricum* subsp. *tauricum*.

Diagnostic characters: pedicels sparsely curved and crisped eglandular pubescent, carpels glabrous.

d) *Aconitum ×teppneri* Mucher ex Starm. nothosubsp. *goetzii* Mucher ex Starm., Ber. Bayer. Bot. Ges. 71: 108. 2001

Type citation: "Österreich, Steiermark, Totes Gebirge, Pühringer Hütte – Salzofen, 21.8.1934, leg. W. Möschl (GZU)" (Starmühler 2001).

Typus: holotype, GZU 000274189.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *lobelii* × *A. tauricum* subsp. *tauricum*.

Diagnostic characters: pedicels sparsely curved and crisped eglandular pubescent; carpels sparsely pubescent on the backside.

e) *Aconitum ×teppneri* Mucher ex Starm. nothosubsp. *haderlappii* Starm., Fritschiana 30: 14. 2002

Type citation: "Slowenien, Kranjska (Krain), Kamniške Alpe (Steiner Alpen), N Kamnik (Stein), bei der Cojzova koča (Zois-Hütte) auf dem Kokrsko sedlo (Kanker-Sattel), 1790 m alt.; Hochstaudenflur; 20.08.1991; leg. W. Mucher (GZU)" (Starmühler 2002).

Typus: holotype, GZU 000232745.

Synonyms: none.

Hybrid origin: *A. napellus* subsp. *lobelii* × *A. tauricum* subsp. *latemarense*.

Diagnostic characters: pedicels glandular and eglandular curved and crisped pubescent and pilose, carpels sparsely pubescent on the backside.

f) *Aconitum ×teppneri* Mucher ex Starm. nothosubsp. *kernerii* Starm., Fritschiana 30: 15. 2002

Type citation: "Aconitum dolomiticum A. Kerner ex Hayek, Flora stiriaca exsiccata Nr. 845, Stiria inferior [Slovenija, Štajerska], in uliginosis prope refugium Kocbekhütte in monte Molčka planina in Alpibus Lithopolitanis (Sanntaler-sive Steineralpen), solo calcareo, 1760–1800 m s.m.; 08.1905 et 1907; leg. A. Hayek et L. Deranc (GZU)" (Starmühler 2002).

Typus: holotype, GZU. Isotypes, GZU 000274195, GZU 000274196, LJU, WU.

Synonyms: *A. dolomiticum* Hayek, p.p., *A. ranunculifolium* Rchb. var. *dolomiticum* Evers, p.p.

Hybrid origin: *A. napellus* subsp. *napellus* × *A. tauricum* subsp. *latemarensis*.

Diagnostic characters: pedicels glandular and eglandular curved and crisped pubescent and pilose, carpels glabrous.

1B. – Subsect. *Burnatii* Rottensteiner

Type species: *A. burnatii* Gáyer.

Description

Differs from subsection *Aconitum* (series *Aconitum*) by the flexuous zigzag stem (at least in the region of the inflorescence) and by glandular pilose inflorescence.

Aconitum burnatii Gáyer, Magyar Bot. Lap. 8: 141. 1909

Type citation: “Alpes Maritimae. Italia, au Col de la Madonna d. Fenestre (Bourgeau pl. alp. mar. n. 13 – W)”.

Typus: not designated.

Synonyms: *A. divergens* Pančić subsp. *burnatii* (Gáyer) W. Seitz, *A. delphinense* Gáyer, *A. napellus* L. subsp. *burnatii* (Gáyer) J.-M. Tison.

Distribution: Maritime Alps, France (Massif Central).

Diagnostic characters: stem in the lower third pubescent; leaves densely pubescent; the uppermost leaf segment (2–) 3–4 mm; pedicels straight; bracteoles linear 2–3 mm long; hood broader than high, falciform; spur of nectaries elongate capitate, filaments sparsely pilose or glabrous; carpels glabrous or pubescent.

Iconography: Molero and Blanché (1986: 238).

Literature: Seitz (1969); Seitz (1970); Rottensteiner (2018).

Hybrids: unknown.

Aconitum maninense (Skalický) Mitka, Studia Universitatis Babeş-Bolyai, Biologia, 62, Sp. Iss. 2017

Basionym: *A. firmum* Rchb. subsp. *moravicum* Skalický var. *maninense* Skalický, Preslia 57: 136, 1985.

Type citation: “Proles localis (natio) angustiarum Maninska úžina et Kostelecká úžina inter Považská Bystrica et Súľov.”

Typus: holotype, PRC 725!

Distribution: Western Carpathians.

Diagnostic characters: stem in the lower third glabrous; leaves sparsely pubescent, the uppermost leaf segment 4–6 mm wide; pedicels straight to bent; bracteoles linear to obovate-lanceolate and then divided; hood as broad

as high, rounded-conic or hemispherical; spur of nectaries hooked; filaments glabrous or sparsely pilose; carpels glabrous or pubescent.

Iconography: Mitka (2003: 178).

Literature: Starmühler and Mitka (2001); Mitka (2003); Mitka et al. (2017).

Hybrids:

a) *Aconitum ×mariae* Rottensteiner, Mitka & Novikov (*nothospec. nov.*) nothosubsp. *mariae* (*A. firmum* subsp. *firmum* × *A. maninense*) – see *A. firmum*.

b) *Aconitum ×mariae* Rottensteiner, Mitka & Novikov nothosubsp. *paxii* (Starm.) Mitka **comb. nov.** (*A. firmum* subsp. *moravicum* × *A. maninense*) – see *A. firmum*.

Aconitum nevadense Uechtr. ex Gáyer, Magyar Bot. Lap. 8: 180. 1909

Type citation: “Sierra Nevada. (Willkomm n. 321 – W), Aleazaba (M. Jimenes – K)”.

Typus: not designated.

Distribution: Sierra Nevada, Pyrenees.

Synonym: *A. occidentale* Timb. Lagr. f. ex Gáyer

Diagnostic characters: stem in the lower third pilose; leaves densely pubescent, uppermost leaf segment (2–) 3–4 mm wide; pedicels bent; bracteoles triangle or linear 1–3 (–4) mm long; hood broader than high, falciform to hemispherical; spur of nectaries capitate; filaments pilose; carpels glabrous or pilose.

Iconography: Molero and Blanché (1986: 236).

Literature: Gáyer (1909); Rottensteiner (2018).

Hybrids: unknown.

Aconitum pentheri Hayek, Ann. Nat. Hofmus. 31: 68. 1917

Type citation: “In monte Žlep prope Ipek, 1800 m.s.m., leg. A. Penther”.

Typus: iconotype in Hayek, Ann. Nat. Hofmus. 31: Taf. 1, 1917.

Synonym: *A. divergens* Pančić nom. illeg. p.p., *A. burnatii* Gáyer subsp. *pentheri* (Hayek) Jalas.

Distribution: Balkans.

Diagnostic characters: stem in the lower third glabrous; leaves ± glabrous, the uppermost leaf segment 3–5 mm wide; pedicels bent; bracteoles linear-lanceolate 1–3 mm long; hood broader than high, falciform; spur of nectaries globose; filaments pilose; carpels glabrous.

Iconography: Hayek (1917).

Literature: Seitz (1969).
Hybrids: unknown.

2. – Sect. *Cammarum* DC.

2A. – Subsect. *Cammarum* (DC.) Rapaics

Description

Leaves divided into 3 or 5 lobes and dissected to the base, uppermost leaf segment ovate or linear with more than two teeth; hoods conical, i.e., above rostrum distinctly elongated, 1.6–2.5 times higher than wider, rarely hemispherical, i.e., above rostrum wide-convex; claws of the nectaries erect or slightly bent, mostly not reaching the top of the hood; spurs of nectaries backward bent or semi-spiral coiled, do not reach the top of hoods; carpels 3 or 5; seeds with membranous 4–6 lamellae and with one conspicuously winged edge, goldish to light brown; root globose.

2a. – Ser. *Variegata* Steinberg ex Starm.

Type species: *A. variegatum* L.

Diagnostic characters: indumentum glabrous or eglandular pubescent.

Aconitum variegatum L., Sp. Pl. ed. 1: 532. 1753

Type citation: "In Italiae, Bohemiae montibus".

Typus: lectotype, Herb. Linn. No. 695.8, LINN (Götz 1967).

Synonyms: *A. cammarum* Jacq., *A. bernhardianum* Wallr., *A. dominii* Sill., *A. gracile* Rchb., p.p., *A. judenbergense* Rchb. (Gáyer), *A. macranthum* Rchb., *A. rostratum* Bernh.

Distribution: Montane and submontane areas of Europe (Alps, Balkans, Sudetes, Transilvania), descending to the lowlands. In Central Europe, it attains the eastern limit of the geographical distribution in the West Carpathians.

Diagnostic characters: inflorescence axis and tepals outside glabrous; hoods conical; filaments glabrous or sparsely pilose; stalks of nectaries erect, glabrous or sparsely pilose; carpels glabrous (subsp. *nasutum*), pilose on the suture (subsp. *variegatum*) or only sparsely pilose on the top of the suture (nothosubsp. *podobnikianum*), pedicel glabrous and bracteoles spathulate to incised (var. *variegatum*) or bracteoles filiform to narrow lanceolate (var. *carniolicum*), or pedicel at least under the bracteoles sparsely pubescent (var. *stiriacum*).

Iconography: Clusius (1601: 98, *pro Aconit. Lycoc. × Thora Italica*); Reichenbach (1827: t. 34; 1840: t. 82 fig. 4682); Novikoff and Mitka (2011: 44).

Literature: Cappelletti and Poldini (1984); Mucher (1991b, 1992); Starmühler (1997a); Mitka and Starmühler (2000); Mitka (2003); Novikoff and Mitka (2011).

Lower taxa:

a) subsp. *variegatum* var. *variegatum*

Diagnostic characters: bracteoles (4) 5–8 (25) mm long, spatulate, ovate or leaflike; carpels 3–5, densely pilose only along the suture; pedicels glabrous.

b) subsp. *variegatum* var. *carniolicum* Starm., Fritschiana 10: 1. 1997

Type citation: "Slowenien, Krain, ENE Laibach (Ljubljana), etwa 3 km NW Sagor (Zagorje), im Tal vom Kotredesch-Bach (Kotredeščica) E der Ruine Gallenberg (Gamberk), 460 m alt., GF 9855/2; Gebüsch; 31.08.1996; leg. A. Podobnik; det. W. Starmühler (LJU)" (Starmühler 1997a).

Typus: holotype, LJU. Isotypes, CL, GJO, GZU, IBF, JACA, KL, KRA, LE, LI, LG, LJM, M, MEL, NY, OSC, PE, Herb. PODOBNIK, SIB, Herb. STARMÜHLER, TBI, TNS, TK, W, WU, Z.

Synonyms: none.

Diagnostic characters: bracteoles short, only 2–3 (5) mm long, linear to narrow lanceolate; carpels always only 3, densely pilose along the suture.

c) subsp. *variegatum* var. *stiriacum* Mucher, Verh. Zool.-Bot. Ges. Österreich 129: 144. 1992

Type citation: "Österreich, Steiermark, Koralpe, Seekar, Hochstauden-flur, 1915 m, 11.9.1991, leg. W. Mucher, GZU" (Mucher 1992).

Typus: holotype, GZU 000206408. Isotypes, GZU 000206409 – GZU 000206420, KL, KRA, W, WU, Herb. MUCHER.

Synonyms: none.

Diagnostic characters: bracteoles (4) 5–8 (25) mm long, spatulate, ovate or leaflike; pedicels sparsely pubescent below the bracteoles; carpels (3–) 5, pilose on the suture and often also on the backside.

Iconography: Mucher (1992: 145, fig. 1).

d) subsp. *nasutum* (Fischer ex Rchb. em. Rupr.) Götz, Feddes Repert. 76 (1–2): 36. 1967

Type citation: "Saxonia metalifera" (Reichenbach 1823: tab. IX, fig. 2).

Typus: not designated, probably iconotype is Rchb. Illustr. Spec. Aconit., tab. IX, fig. 2, and tab. X, 1823.

Synonyms: *A. nasutum* Fischer ex Rchb., p.p.

Diagnostic characters: carpels 3, glabrous.

e) nothosubsp. *podobnikianum* – see hybrids of *A. variegatum*.

Hybrids:

a) *Aconitum × austriacum* Mucher. (*A. pilipes* × *A. variegatum*) – see nothoser. *Toxigata*.

b) *Aconitum × aquilonare* A. Kerner ex Gáyer, Magyar Bot. Lapok 10: 201. 1911

Type citation: "Croatia – Fužine. Borbás – HUW" (Gáyer, 1911).
Typus: lectotype, WU-Kerner 0034380.

Synonym: *A. variegatum* L. fo. *aquilonare* (Gáyer) M. Niketić.
Hybrid origin: *A. variegatum* × *A. vitosanum*.

Diagnostic characters: pedicels sparsely eglandular curved pubescent; tepals outside glabrous; carpels mostly glabrous.

c) *Aconitum × hebegynum* DC. (*A. degenii* × *A. variegatum*) – see nothoser. *Toxigata*.

d) *Aconitum × pawlowskii* Mitka & Starm. (*A. lasiocarpum* × *A. variegatum*) – see nothoser. *Toxigata*.

e) *Aconitum variegatum* L. nothosubsp. *podobnikianum* Mucher, Carinthia II 183/103 (2): 522. 1993

Type citation: "Aconitum variegatum L. subsp. *variegatum*, Slovenija, Nanos, ad orientem a refugio alpino Vojkova koca dicto, ad margines silvarum, 1240 m s.m., 0250/1, Nr. 117683(1), 4. 10. 1987, leg. A. Podobnik, LJU. Isotypes: GZU, LJU" (Mucher 1993).
Typus: holotype, LJU 117683(1). Isotypes, GZU 000274191, LJU.

Synonyms: none.

Hybrid origin: *A. variegatum* subsp. *nasutum* × *A. variegatum* subsp. *variegatum*.

Diagnostic characters: pedicels glabrous or sparsely pubescent only below bracteoles; stamens glabrous; carpels 3, only sparsely pilose on the top of the suture.

Aconitum vitosanum Gáyer, Magyar Bot. Lap. 8: 203. 1909

Type citation: "Bulgaria: Vitoša, inter saxa i. monte medio, Keck et Pichler Pl. Bulg. 1890".

Typus: holotype, W (?). Isotypes, WU-Halácsy-Europ. 0034442, WU 0034443, WU 0034444.

Synonyms: *A. variegatum* L. var. *diffusum* (Rchb.) Penev & Simeon. (*nom. nud.*).

Distribution: Balkans, Northern Italy.

Diagnostic characters: Inflorescence axis and pedicels pubescent; petals outside curved eglandular pubescent;

hoods hemispherical or rounded-conic; filaments glabrous; stalks of nectaries glabrous; carpels glabrous or pubescent on the backside.

Iconography: Penev and Simeonovski (1970: 63).

Literature: Starmühler (1996c).

Hybrids: *A. × aquilonare* A. Kern. ex Gáyer (*A. variegatum* × *A. vitosanum*) – see *A. variegatum*.

***Aconitum vivantii* Rottensteiner, nom. nov.**

Basionym: *A. variegatum* subsp. *pyrenaicum* Vivant & Delay, Bull. Soc. Bot. France, Lettres Bot. 127: 501, 1981. There is a new name necessary in the rank of a species because the name *A. pyrenaicum* is already occupied by a taxon from the subgenus *Lycocodium*.

Type citation: "haute vallée d'Aspe (Pyr.-Atl.) en amont du hameau des Forges d'Abel. ... vers 1700 m d'altitude au-dessus de la hetaïre-sapinére d'Esplungére ..." .

Typus: not designated.

Synonyms: none.

Distribution: Pyrenees: Navarra, Guipúzcoa (Spain).

Diagnostic characters: inflorescence axis and tepals outside glabrous; pedicels almost always glabrous or very rarely sparsely pubescent just below flowers; hoods hemispherical or rounded-conic; filaments glabrous; nectaries strongly bent, mostly reaching the top of the hood, glabrous; carpels 3, glabrous.

Iconography: Molero and Blanche (1986: 240).

Hybrids: most probably with *A. napellus*.

2b. – Ser. *Toxica* (Rchb.) Mucher

Type species: *A. toxicum* Rchb.

Diagnostic characters: indumentum glandular pilose.

***Aconitum degenii* Gáyer, Magyar Bot. Lap. 5: 123. 1906**

Type citation: "Comit. Beszterce-Naszód: in lapidosis, umbrosis ad cacumen montis calcarei Craciunel prope Rodnam; in lapidosis graminosis montis calc. Galarin; secus rivulos montanos ad Rodna-Borberek; Comit. Csik: in montibus calcareis Ócsém et Nagy-Hagymás; Comit. Máramaros: Oberhalb der Luhyer Klause Hoverla im Fichtenwalde; Comit. Torda-Aranyos: in valle Csepilor sub alpe Nagy-Bihar; Comit. Kolozs: ad margines silvarum prope Rogozsely sub tractu Vlegyászae".

Typus: lectotype, CL-Soó!

Synonyms: *A. hebegynum* auct. Fl. Carpathorum non DC., *A. molle* Rchb., *A. paniculatum* Lam., p.p. (*nom. inval.*).

Distribution: Alps, Carpathians, Balkans.

Diagnostic characters: hood rounded-conic; carpels (3–5 pubescent (subsp. *valesiacum*) or glabrous, pedicels

and petals glandular pilose, bracteoles in the middle of pedicels (subsp. *paniculatum*), filaments glabrous (var. *laxiflorum*) or pubescent (var. *turrachense*), pedicels and petals glabrous (subsp. *rhaeticum*); bracteoles in the upper part of the pedicels or just below flower (subsp. *degenii*), spur of nectaries semi-spiral coiled (fo. *degenii*) or capitulate (fo. *craciunelense*); pedicels below bracteoles and hood ± glabrous (var. *intermedium*).

Iconography: Mucher (1993a: 55); Novikoff and Mitka (2011: 46). Literature: Mucher (1993a); Starmühler (1996d, 1996e); Ilnicki and Mitka (2011); Novikoff and Mitka (2011); Ziman et al. (2015).

Lower taxa:

a) subsp. *degenii* var. *degenii* fo. *degenii*

Diagnostic characters: bracteoles lanceolate to spathulate, in the upper part of pedicel; pedicels below the bracteoles densely glandular pilose; tepals outside glandular pilose; hood ± hemispheric; carpels 3, glabrous.

b) subsp. *degenii* var. *degenii* fo. *craciunelense* Gáyer, Magyar Bot. Lap. 5: 126. 1906

Type citation: "Comit. Beszterce-Naszód: ad cacumen montis Craciunel. (Degen, H.D.)".

Type specimens: not designated.

Synonyms: none.

Diagnostic characters: bracteoles lanceolate to spathulate, in the upper part of pedicel; pedicels below the bracteoles densely glandular pilose; tepals outside glandular pilose; hood cymbiform; carpels 3, glabrous.

c) subsp. *degenii* var. *intermedium* (Zapał.) Mitka, Caryologia 64(4): 449. 2011

Type citation: "Płaj między Mołodą a Jajcem nad Łomnicą, Tomatek w Polenicy Popowiczowskiej, Jabłonica na Przesmyku Tatarskim (*Wołoszczak*), Jawornik koło Mikuliczyna (*Rehman*), Bystrzec pod Czarną Horą (*Ślęzakiński*)" (Zapałowicz 1908b: 221). Typus: lectotype, KRAM 132196 (Mitka 2003).

Synonyms: *A. hebegynum* auct. Fl. Carpathorum, non DC., *A. paniculatum* Lam. var. *prutense* Zapał., *A. paniculatum* fo. *latiusculum* Zapał.

Diagnostic characters: bracteoles lanceolate to spathulate, in the upper part of pedicel; pedicels below the bracteoles only sparsely glandular pilose; tepals outside glandular pilose; carpels 3, glabrous.

d) subsp. *paniculatum* (Arcang.) Mucher var. *turrachense* (Mucher) Mucher, Phyton (Horn) 33 (1): 63. 1993

Type citation: "Ostalpen, Steiermark, Gurktaler Alpen, 2 km südlich von Predlitz, Gebüschsaum, 1020 m, 26.8.1989, leg. W. Mucher (GZU)" (Mucher 1993a).

Typus: holotype, GZU 000274182. Isotypes, GJO, GZU 000274183, GZU 000274185 – GZU 000274188, W. Paratype, GZU 000274184.

Synonyms: *A. paniculatum* Lam. subsp. *paniculatum* var. *turrachense* Mucher.

Diagnostic characters: bracteoles linear, in the middle of the pedicel; tepals outside glandular pilose; nectaries glabrous or pilose; filaments ± densely pubescent.

Iconography: Mucher (1990).

e) subsp. *paniculatum* (Arcang.) Mucher var. *laxiflorum* (Rchb.) Mucher, Phyton (Horn) 33 (1): 60. 1993

Typus: lectotype, see Mucher (1993a) for details.

Synonyms: *A. cernuum* Rchb. non Wulfen [var.] *α laxiflorum* Rchb., *A. paniculatum* subsp. *paniculatum* var. *paniculatum* (nom. inval.).

Diagnostic characters: bracteoles linear, in the middle of the pedicel; tepals outside glandular pilose; nectaries and filaments glabrous.

Iconography: Mucher (1993a).

f) subsp. *rhaeticum* Starm., Fritschiana 7: 2. 1996

Type citation: "Schweiz, Graubünden, Oberhalbstein S Tiefencastel, Abhang zum rechten Ufer des Adont N Präsanz (Personz), 1360–1375 m alt., GF 9423/2; *Alnus incana*–*Prunus padus*-Wald; 14.08.1995; leg./det. W. Starmühler (GZU)" (Starmühler 1996d).

Typus: holotype, GZU 000212973. Isotypes: LE, M, NY, PE, TI, W, Z, Herb. STARMÜHLER.

Synonyms: none.

Diagnostic characters: tepals outside, pedicels and nectaries glabrous.

Iconography: Starmühler (1996d).

g) subsp. *valesiacum* (Gáyer) Mucher, Phyton (Horn) 33(1): 64. 1993

Type citation: "Mauvoisin, vallee de Bagne [Suisse, Valais] (BP, Götz 1967)" (Mucher 1993a).

Typus: holotype, BP.

Synonyms: *A. valesiacum* Gáyer, *A. variegatum* L. subsp. *valesiacum* (Gáyer) Greuter & Burdet.

Diagnostic characters: bracteoles lanceolate to spathulate, situated at the pedicel's top; pedicels above the bracteoles densely glandular pilose, below the bracteoles – sparsely pilose or glabrous; tepals outside glandular pilose; carpels 5, densely pilose; nectaries and filaments glabrous.

Iconography: Götz (1967), Mucher (1993a).

- h) nothosubsp. ***gandogeri*** Mucher (*A. degenii* subsp. *paniculatum* × *A. degenii* subsp. *valesiacum*) – see hybrids of *A. degenii*.
- i) nothosubsp. ***lippertianum*** Starm. (*A. degenii* subsp. *paniculatum* × *A. degenii* subsp. *rhaeticum*) – see hybrids of *A. degenii*.

Hybrids:

- a) ***Aconitum degenii*** Gáyer nothosubsp. ***gandogeri***
Mucher, Phyton (Horn) 33 (1): 65, 1993

Type citation: “Suisse, Vaud, Pays d’Enhaut, Vallee de Gerignoz, LaMontagnette, alt. 1445 m, cs. 579/143; 17.8.1952; leg. A. Maillefer, LAU (as *A. paniculatum* Lam.)” (Mucher 1993).

Typus: holotype, LAU.

Synonyms: none.

Hybrid origin: *A. degenii* subsp. *paniculatum* × *A. degenii* subsp. *valesiacum*.

Diagnostic characters: the filaments of the hybrid are in most cases pilose, although both parents have glabrous filaments; pedicels below the bracteoles almost glabrous; carpels mostly glabrous; tepals outside glandular pilose.

- b) ***Aconitum degenii*** Gáyer nothosubsp. ***lippertianum***
Starm., Fritschiana 7: 11. 1996

Type citation: “Schweiz, Kt. Graubünden, Oberhalbstein S Tiefencastel, Abhang zum rechten Ufer des Adont N Präsanz (Personz), 1325 m alt., GF 9423/2; *Alnus incana*-Wald mit *Corylus avellana*, *Sorbus aucuparia*, *Berberis vulgaris* etc.; 14.08.1995; leg./det. W. Starmühler (GZU)” (Starmühler 1996e).

Typus: holotype, GZU 000212947. Isotypes, LE, M, PE, Herb. STARMÜHLER, Z.

Synonyms: none.

Hybrid origin: *A. degenii* subsp. *paniculatum* × *A. degenii* subsp. *rhaeticum*.

Diagnostic characters: pedicels sparsely glandular pilose above the bracteoles; tepals glabrous.

- c) ***Aconitum ×dragulescuanum*** Mucher (*A. degenii* subsp. *degenii* × *A. toxicum* subsp. *toxicum*) – see *A. toxicum*.

- d) ***Aconitum ×gayeri*** Starm., Stapfia 45: 41. 1996

Type citation: “Tanssilvania, Rodna – Borbereki [= Valea Vinului], Nr. 135419; 28.8.1897; L. Walz (CL)” (Starmühler, 1996b).

Typus: holotype, CL 135419! Isotype, Detto, Nr. 135418 (CL).

Synonyms: none.

Hybrid origin: *A. degenii* × *A. lasiocarpum*.

Diagnostic characters: hood rounded-conic or conic, petals outside pilose, carpels pilose, spur of nectaries reaching the top of the hood, pedicels above bracteoles densely glandular pilose, below bracteoles sparsely pilose, bracteoles linear to lanceolate.

- e) ***Aconitum ×hebegynum*** DC. (*A. degenii* × *A. variegatum*) – see nothoser. *Toxigata*

- f) ***Aconitum ×mielichhoferi*** Rchb. (*A. degenii* × *A. tauricum*) – see nsect. *Acomarum*

- g) ***Aconitum ×pilosiusculum*** (Ser.) Gáyer, Magyar Bot. Lapok 10: 200. 1911

Type citation: “Neotype: Patura-ge de Lavaraz, Alpes de Bex, 26.9.1879, Jaccard (ZT)” (Starmühler 1996a).

Typus: neotype, ZT.

Synonyms: *A. variegatum* L. var. *pilosiusculum* Ser., *A. rostratum* Bernh. ex DC. var. *pilosiusculum* Ser.

Hybrid origin: *A. degenii* × *A. pilipes*.

Diagnostic characters: hood rounded-conic, bracteoles lanceolate to spatulate, tepals outside pilose, carpels pilose, pedicels glandular pilose and eglandular pubescent.

- Aconitum lasiocarpum*** Rchb., Uebers. Aconitum: 55. 1819

Type citation: “in Hungar Comit. Marmarosch. *Sadler!*”.

Typus: neotype, “An schattigen Stellen in der Voralpenregion auf dem Gebirge Craciunel bei Alt Rodna, 1.8.1882, leg. *Porcius*” – WU-Kerner” (Mitka 2003).

Synonyms: *A. cammarum* L. var. *beskidene* Zapał., *A. dasycarpum* Schur.

Distribution: Western and Eastern Carpathians, Podilya (Ukraine).

Diagnostic characters: inflorescence axis and tepals outside glandular pilose, hood conical; pedicels glandular pilose (subsp. *lasiocarpum*) or below bracteoles glabrous or sparsely eglandular pilose and curved and crisped eglandular pubescent (subsp. *kotulae*); bracteoles linear or lanceolate, rarely spatulate, (2.5–) 3–6 (–12) mm; carpels glandular pilose.

Iconography: Reichenbach (1827: t. 9); Mitka (2003: 175); Novikoff and Mitka (2011: 44).

Literature: Gáyer (1911); Starmühler (1993, 1996a); Joachimiak et al. (1999); Mitka and Starmühler (2000); Mitka (2003).

Lower taxa:

a) subsp. *lasiocarpum*

Diagnostic characters: pedicels entirely glandular pilose.

b) subsp. *kotulae* (Pawl.) Starm. & Mitka, Acta Soc. Bot. Polon. 69 (2): 150. 2000

Type citation: "Tatry Wysokie [High Tatras]: dol. Koprowa – skraj lasu. ok. 1300 m: 19.8.1925; leg. B. Pawłowski (KRAM-Pawłowski 320228 – Neotype)" (Mitka and Starmühler 2000).

Typus: neotype, KRAM-Pawłowski 320228. Paratype, KRAM 132168.

Synonyms: *A. variegatum* L. subsp. *kotulae* Pawł., *A. variegatum* fo. *kotulae* (Pawl.) Skalický, *A. paniculatum* Lam. var. *podolicum* Zapał. fo. *podolicum*, *A. paniculatum* Lam. var. *podolicum* Zapał. fo. *latilobum* Zapał., *A. podolicum* (Zapał.) Voroshylov, *A. hebegeynum* auct., p.p., *A. beskidense* (Zapał.) Gáyer, *A. cammarum* Jacq. var. *koscieliskanum* Zapał., p.p., *A. cammarum* Jacq. var. *beskidense* Zapał., *A. gracile* Rchb. subsp. *grosserratum* (Gáyer) Grinč. fo. *beskidense* Zapał.

Diagnostic characters: pedicels below bracteoles glabrous or sparsely eglandular pilose and curved and crisped eglandular pubescent.

Hybrids:

a) *Aconitum ×gayeri* Starm. (*A. degenii* × *A. lasiocarpum*) – see *A. degenii*.

b) *Aconitum lasiocarpum* × *A. toxicum* – hood round-ed-conic, bracteoles linear-ovate with reticulate venation, pedicels glandular pilose, carpels pilose.

c) *Aconitum ×pawlowskii* Mitka & Starm. (*A. lasiocarpum* × *A. variegatum*) – see nothoser. *Toxigata*.

***Aconitum pilipes* (Rchb.) Gáyer, Magyar Bot. Lap. 10: 200. 1911**

Basionym: *A. cammarum* L. var. *pilipes* Rchb., Fl. Germ. Exurs. 738, 1832.

Type citation: "Salvesental bei Imst".

Typus: neotype, WU-Kerner.

Distribution: Western and Eastern Alps.

Diagnostic characters: hood conical, outside glandular pilose; pedicels curved and s-formed glandular pilose and eglandular pubescent; bracteoles spatulate 3–6 mm, situated just below the flower; filaments pilose; carpels 5, entirely pubescent.

Iconography: Starmühler (1996b: 171).

Literature: Starmühler (1996b, 2001).

Hybrids:

a) *Aconitum ×austriacum* Mucher, Carinthia II 183/103(2): 524. 1993

Type citation: "Österreich, Salzburg, Hohe Tauern, Habachtal, Weg von der Moar-Alm zum Noitro-Steig, Hochstaudenflur, 1540 m, 30. 8. 1991, leg. W. Mucher, GZU" (Mucher 1993b).

Typus: holotype, GZU 000206194. Isotypes: GJO, GZU 000206195, KL, KRA,W, WU 0034381, Herb. MUCHER.

Synonyms: none.

Hybrid origin: *A. pilipes* × *A. variegatum*.

Diagnostic characters: carpels sparsely pubescent or pubescent at the suture.

b) *Aconitum ×pilosiusculum* (Ser.) Gáyer (*A. degenii* × *A. pilipes*) – see *A. degenii*.

c) *Aconitum napellus* × *A. pilipes* – see nothosect. *Aconitum*

d) *Aconitum pilipes* × *A. tauricum* – see nothosect. *Aconitum*

***Aconitum toxicum* Rchb., Uebers. Aconitum: 43. 1819**

Type citation: "Hab. in Transylvaniae alpibus. Baumgarten!".

Typus: lectotype, "Transsylv., in graminosis alpestris, [I]827, leg. Baumgarten (as *A. neomontanum*, rev. E. Götz 1966 as *A. toxicum* Rchb.)", W (Mucher 1993).

Synonyms: *A. bosniacum* Beck, *A. diabolicum* Gáyer, *A. neomontanum* auct. non Koelle, *A. schurii* Beck., *A. schurii* fo. *retyezatense* Gáyer, *A. toxicum* Rchb. *bosniacum* (Beck) Hayek, *A. toxicum* Rchb. fo. *diabolicum* (Gáyer) Grint., *A. toxicum* [var.] β *schurii* Beck.

Distribution: Transilvania, Eastern Carpathians (Ceahlau Mts), Southern Carpathians, Balkans.

Diagnostic characters: pedicels glandular pilose (subsp. *toxicum*), glandular/eglandular pubescent (subsp. *crispulum* and nothosubsp. *nyaradyanum*), sparsely straight glandular pilose (nothosubsp. *ungarianum*), or glabrous (subsp. *bucegiense*); hood outside glandular pilose (subsp. *toxicum*), glandular/eglandular pubescent (subsp. *crispulum* and nothosubsp. *nyaradyanum*), or glabrous (subsp. *bucegiense* and nothosubsp. *ungarianum*); filaments in the upper part, pilose; hood rounded-conic or conical; carpels glabrous; bracteoles ovate, sometimes divided, with reticulate venation.

Iconography: Reichenbach (1827: t. 37).

Literature: Mucher (1993a, 1993b); Mitka (2000); Starmühler (2000).

Lower taxa:

a) subsp. ***toxicum***

Diagnostic characters: pedicels and tepals outside with straight glandular trichomes.

b) subsp. ***crispulum*** (Nyár.) Mucher, Phyton (Horn) 33(1): 70. 1993

Type citation: "Transsilvania, distr. Brasov, ad rivum Gr. Weidenbach sub mtibus Bucegi prope pagum Rasnov, alt. cca. 720 m s.m., 12.8.1929, leg. E.I. Nyárády (SIB-Nyárády)" (Mucher 1993a).

Typus: lectotype, SIB-Nyárády. Paratype, SIB-Nyárády.

Synonyms: *A. toxicum* Rchb. var. *crispulum* Nyárády in Borza.

Diagnostic characters: pedicels and tepals outside with curved glandular trichomes.

Iconography: Mucher (1993a).

c) subsp. ***bucegiense*** (Nyár.) Mucher, Phyton (Horn) 33(1): 68. 1993

Type citation: "Transsilvania, distr. Brasov, ad pedem montium Bucegi, prope Valea Topliei, versus pagum Rasnov, in querketis, una cum typo, alt. cca 700 m s.m., Nr. 601109, 22.9.1940, leg. P. Cretzoiu, det. E.I. Nyárády (CL)" (Mucher 1993a).

Typus: holotype, CL. Isotypes, CL, SIB.

Synonyms: *A. toxicum* Rchb. var. *bucegiense* Nyár.

Diagnostic characters: pedicels glabrous.

Iconography: Mucher (1993a).

d) nothosubsp. ***nyaradyanum*** Mucher – see hybrids of *A. toxicum*.

e) nothosubsp. ***ungarianum*** Starm. – see hybrids of *A. toxicum*.

Hybrids:

a) ***Aconitum ×dragulescuanum*** Mucher, Phyton (Horn) 33(1): 72. 1993 nothosubsp. ***dragulescuanum***

Type citation: "Transsilvania, in praeruptis et graminosis rupium calc. Mt. Hagimasul Mare, supra rivum Nagyág, alt. 1500–1740 m s.m., distr. Ciuc; 19.8.1948; J.E. Nyárády (SIB-Nyárády)" (Mucher 1993).

Typus: holotype, SIB-Nyárády.

Synonyms: none.

Hybrid origin: *A. degenii* subsp. *degenii* × *A. toxicum* subsp. *toxicum*.

Diagnostic characters: pedicels, bracteoles and tepals outside with straight glandular trichomes only.

b) ***Aconitum ×dragulescuanum*** Mucher nothosubsp. ***rintescuanum*** Starm., Siebenbürg. Arch. 36: 16. 2000

Type citation: "Rumänien, Siebenbürgen (Transsilvania, Erdély), Süd-Karpaten (Carpatii Meridionali, Déli Kárpátok), Retezat-Gebirge (Muntii Retezatului, Retezat-havasok) S Hotzing (Hateg, Hátszeg), Nationalpark Retezat, am Weg von der Gura Zlata-Hütte zum Retezat, am Bach Dobrun, 1470 m"

Typus: holotype, GZU 000232748.

Synonyms: none.

Hybrid origin: *A. degenii* subsp. *degenii* × *A. toxicum* subsp. *crispulum*.

Diagnostic characters: pedicels, bracteoles and tepals outside with curved and straight glandular trichomes.

c) ***Aconitum lasiocarpum*** × ***A. toxicum*** – bracteoles spathulate or linear-ovate with reticulate venation, hood slightly conical or round-conic, carpels pilose.

d) ***Aconitum toxicum*** Rchb. nothosubsp. ***nyaradyanum*** Mucher, Phyton (Horn) 33(1): 71. 1993

Type citation: "Transsilvania, distr. Brasov, ad rivum Gr. Weidenbach sub mtibus Bucegi prope pagum Rasnov, alt. cca. 720 m s.m., Nr. 195107; 12.8.1929; leg. E.I. Nyárády (CL)" (Mucher 1993a).

Typus: holotype, CL 195107.

Synonyms: none.

Hybrid origin: *A. toxicum* subsp. *crispulum* × *A. toxicum* subsp. *toxicum*.

Diagnostic characters: pedicels and tepals outside glandular/eglandular pubescent.

e) ***Aconitum toxicum*** Rchb. nothosubsp. ***ungarianum*** Starm., Siebenbürg. Arch. 36: 21. 2000

Type citation: "Siebenbürgen (Transsilvania, Erdély), Burzenländer Senke (Depresiunea Birsei, Barcasági-medence) SW Kronstadt (Brasov, Brassó), S Rosenau (Risnov, Barcarozsnyó), am rechten Ufer vom Grossen Weidenbach (Ghimbasel), 720 m."

Typus: holotype, GZU 000232749.

Synonyms: none.

Hybrid origin: *A. toxicum* subsp. *bucegiense* × *A. toxicum* subsp. *toxicum*.

Diagnostic characters: tepals outside glabrous or almost glabrous; pedicels sparsely glandular pilose, mainly above the bracteoles.

2c. – Nothoser. *Toxigata* Starm. (ser. *Variegata* × ser. *Toxica*)

Type species: *A. ×hebegynum* DC., designated here.

Diagnostic characters: indumentum glandular and eglandular pubescent.

Aconitum ×austriacum Mucher, Carinthia II 183/103(2): 524. 1993

Type citation: “Österreich, Salzburg, Hohe Tauern, Habachtal, Weg von der Moar-Alm zum Noitroi-Steig, Hochstaudenflur, 1540 m, 30. 8. 1991, leg. W. Mucher” (Mucher, 1993b).

Typus: holotype, GZU 000206194. Isotypes, GZU 000206195, WU 0034381, GJO, KL, KRA, W, Herb. MUCHER.

Synonyms: none.

Hybrid origin: *A. pilipes* × *A. variegatum*.

Diagnostic characters: Pedicels sparsely crisped to s-formed glandular pilose, mainly above the bracteoles; tepals outside glabrous; hood higher than broad; nectaries claw upright; nectaries spur curved backward, not reaching the top of the hood; carpels (3) 5, mostly +/- densely pilose.

Iconography: Mucher (1993b).

Literature: Starmühler (1996a).

Aconitum ×bartokianum Starm., Siebenbürg. Arch. 36: 10. 2000

Type citation: “Aconita rarissima selecta (1999), Rumänien, Transilvania (Siebenburgen, Erdély), Süd Karpaten, Munții Retezatului [...] 14.08.1992; leg. W. Mucher et U. Starmühler (GZU)”.

Typus: holotype, GZU 000225135. Isotypes, CL, GJO, Herb. STARMÜHLER, TNS.

Synonyms: none.

Hybrid origin: *A. toxicum* × *A. variegatum*.

Distribution: Transilvania, Southern Carpathians.

Diagnostic characters: hood conical; bracteoles spathulate or linear-ovate with reticulate venation; nectaries reaching the top of the hood; filaments glabrous or sparsely pilose. Pedicel and tepals outside straight glandular pilose; carpels glabrous – nothosubsp. *bartokianum*. Pedicel and tepals outside curved glandular pilose; carpels glabrous – nothosubsp. *rapaicsianum*. Pedicel and tepals outside straight glandular pilose; carpels pilose on the suture – nothosubsp. *sooanum*.

Iconography: Starmühler (2000: 11).

Literature: Starmühler (2000).

Lower taxa/hybrids:

a) nothosubsp. *bartokianum*

Hybrid origin: *A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *nasutum*.

Diagnostic characters: pedicel and tepals outside straight glandular pilose; carpels glabrous.

b) nothosubsp. *rapaicsianum* Starm., Siebenbürg. Arch. 36: 14. 2000

Type citation: “Aconita rarissima selecta (1999), Rumänien, Transilvania (Siebenburgen, Erdély), Süd-Karpaten, Munții Retezatului (Retezat-Gebirge, Retyezát-havasok), Nationalpark Retezat, S-Hang des Retezat, 1980 m alt.; Hochstaudenflur mit *Achillea distans*, *Adenostyles alliariae*, *Angelica sylvestris*, *Carduus personata*, *Deschampsia cespitosa*, *Rumex alpinus*, *Seriecio ovatus*, *Senecio cacaliaster* etc.; 14.08.1992; leg. W. Mucher et U. Starmühler (GZU)” (Starmühler 2000).

Typus: holotype, GZU 000225136. Isotypes, CL, GJO, Herb. STARMÜHLER, TNS.

Synonyms: none.

Hybrid origin: *A. toxicum* subsp. *crispulum* × *A. variegatum* subsp. *nasutum*.

Diagnostic characters: pedicel and tepals outside curved glandular pilose; carpels glabrous.

Iconography: Starmühler (2000: 11).

c) nothosubsp. *sooanum* Starm., Siebenbürg. Arch. 36: 15. 2000

Type citation: “Hungaria media, Alpes Biharienses, ad balnea “Biharfüred”; 13.07.1914; leg. L.T. Haisz (CL-Soó)” (Starmühler 2000).

Typus: holotype, CL-Soó.

Synonyms: none.

Hybrid origin: *A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *variegatum*.

Diagnostic characters: pedicel and tepals outside straight glandular pilose; carpels pilose along the suture.

Iconography: Starmühler (2000: 11).

Aconitum ×hebegynum DC., Regn. Veg. Syst. Nat.: 376. 1817, pro spec.

Type citation: “Hab. in dumetis Alpium Vallesiae, Austriae (Jacq.) et in Carpathorum convallibus (Wahlenb.)”.

Typus: neotype, “(Schweiz) cultivé, provenant d'un pied vivant récolté à la Gemmi – G-DC” (Starmühler 1997b).

Synonyms: none.

Hybrid origin: *A. degenii* × *A. variegatum*.

Distribution: Alps, Balkans.

Diagnostic characters: Hood about as high as broad or slightly higher; nectary claw curved; nectary spur backward bent always reaching the top of the hood; bracteoles linear-lanceolate, rarely spathulate 4–6 (–8) mm long, mostly glandular pilose. Pedicels and tepals outside sparsely glandular pilose; carpels 3–5 glabrous or sparsely glandular pubescent.

Iconography: Mucher (1993b: 519–527).

Literature: Gáyer (1906).

Aconitum ×pawłowskii Mitka & Starm., Acta Soc. Bot. Pol. 69: 151. 2000

Type citation: “(Tatra Mts.) Tatry, pod Raptawickimi Turniami (dol. Kościeliska) od półn., ok. 1300 m, piargi wapienne, 3.9.1932, leg. B. Pawłowski”.

Typus: holotype, KRAM-Pawłowski 320221.

Synonyms: none.

Hybrid origin: *A. lasiocarpum* × *A. variegatum*.

Distribution: West Carpathians.

Diagnostic characters: Hood conical; nectary claw curved; nectary spur backward bent not reaching the top of the hood; bracteoles linear-lanceolate, lanceolate, glabrous or ciliate 2.5–6 (–12) mm long. Pedicels above bracteoles straight glandular pilose, below bracteoles sparsely straight glandular pilose or appressed curved pubescent; carpels 3–5 on the suture pilose.

Iconography: Mitka and Starmühler (2000: 151); Mitka (2003: 173).

Literature: Mitka and Starmühler (2000); Mitka (2003).

Aconitum ×tuscheticum (N.Busch) N.Busch, Opred. Rast. Kryma i Kavkaza 30. 1919

Basionym: *A. caucasicum* N.Busch subsp. *pubiceps* (Rupr.) N.Busch var. *tuscheticum* N.Busch, Tr. Jur’jev. Bot. Sada 1: 3, 1900.

Type citation: “Dzhvari-Vozeli. 13.VII.[18]76. Radde” (LE).

Typus: holotype, LE (?).

Synonyms: *A. caucasicum* N.Busch subsp. *pubiceps* (Rupr.) N. Busch var. *tuscheticum* N.Busch, *A. nasutum* Rchb. subsp. *tuscheticum* (N.Busch) A.N.Luferov, *A. pubiceps* Rupr. var. *tuscheticum* (N.Busch) Grossh.

Hybrid origin: *A. degenii* × *A. vitosanum*.

Distribution: Caucasus, Slovenia.

Diagnostic characters: plant 50–100 cm high; inflorescence axis, pedicels and tepals outside glandular and eglandular pubescent.

Iconography: none.

Literature: Luferov (2000); Starmühler (1996c).

3. – Nothosect. *Acomarum* Starm. (= nothosect. *Acopellus* Starm.)

Type species: *A. ×cammarum* L. em. Fries, designated here.

Diagnosis

Carpels totally or partially sterile.

Description

Leaves divided into 3 or 5 lobes and dissected to the base; hoods hemispherical or conical, 1.2–2.0 (–2.5) times higher than wider; claws of the nectaries erect or slightly bent, reaching or not the top of the hood; spurs of nectaries backward bent or semi-spiral coiled; carpels 3 or 5; seeds with membranous 4–6 lamellae and with one conspicuously winged edge; inflorescence ramosc or rarely dense ramified; root elongated.

Aconitum ×acuminatum Rchb., Uebers. Aconitum 46. 1819

Type citation: “Hab. in monte Baldo Heineken! Hb. Mertens”.

Typus: not designated.

Synonym: *A. lobelianum* Host. in Gáyer fo. *brachytrichum* Gáyer.

Hybrid origin: *A. degenii* × *A. napellus*.

Distribution: Alps.

Diagnostic characters: hood glandular pilose and eglandular pubescent, pedicels above bracteoles glandular pilose and below eglandular pubescent, carpels 3, glabrous or overall pubescent.

Iconography: Reichenbach (1827: t. 68).

Literature: Starmühler (2001).

Aconitum ×acutum Rchb., Uebers. Acon. 21. 1819

Type citation: “Hab. in valle Kalstahl Tyrol. Sieber! in Alp. Seethal supra Judenburg v. Vest!”.

Typus: not designated.

Synonym: *A. ×zahlbruckneri* Gáyer.

Hybrid origin: *A. tauricum* × *A. variegatum*.

Distribution: Eastern and Southeastern Alps.

Diagnostic characters: hood glabrous, hemispherical; pedicels glabrous or eglandular pubescent; filaments glabrous or sparsely pilose, stalks of nectaries glabrous or pubescent; carpels mostly glabrous.

Iconography: Reichenbach (1821: Tab. 12, 2 pro species).

Literature: Starmühler (2001).

Aconitum ×berdaui Zapał., Rozpr. Wydz. Mat.-Przyr. Akad. Umiej., Dział B. Nauki Biol. 48: 88. 1908a, pro hybr. [reprinted in Zapałowicz, 1908b].

Type citation: "In regione subalpina Tatorum: in valle Kościeliska et altero loco non indicato (*Berdaui*), Podspady ad Jaworzynka".

Typus: lectotype, "Dol. Kościeliska, ?08.1855, leg. *Berdaui*, det. Zapałowicz (16.02.1908), KRAM 133482!" (Waclawska-Ćwiertnia and Mitka 2016).

Synonyms: *A. lengyelii* Gáyer, *A. odotandrum* Wissjul.

Hybrid origin: *A. firmum* × *A. variegatum*.

Distribution: Europe: Sudetes, Transilvania and the Balkans.

Diagnostic characters: inflorescence axis, pedicels and tepals outside pilose (nothosubsp. *walasii*) or glabrous (nothosubsp. *berdaui*); hoods conical; filaments glabrous or sparsely pilose, spur of the nectaries semi-spiral coiled; carpels pilose on the suture.

Iconography: Mitka (2003: 173); Novikoff and Mitka (2011: 44).

Literature: Wissjulina (1939); Starmühler and Mitka (2001); Mitka (2003), Novikoff and Mitka (2011), Waclawska-Ćwiertnia and Mitka (2016).

Lower taxa/hybrids:

a) nothosubsp. ***berdaui***

Hybrid origin: *A. firmum* subsp. *firmum* × *A. variegatum* subsp. *variegatum*.

Diagnostic characters: inflorescence axis, pedicels and tepals outside glabrous.

Iconography: (Mitka 2003: 173, fig. 6D).

b) nothosubsp. ***walasii*** (Mitka) Mitka, The genus *Aconitum* in Poland and adjacent countries: 56. 2003

Type citation: "Poland, Western Carpathians, Pogórze Spisko-Gubałowskie Foothills), Podhale, Wzn. Gubałowskie, Roztoki, żwirowiska nad Czarnym Dunajcem, 22.8.1957, leg. E. Pancer" (Mitka 2003).

Typus: holotype, KRA 003852.

Synonyms: *A. xlengyelii* Gáyer nothosubsp. *walasii* Mitka.

Hybrid origin: *A. firmum* subsp. *moravicum* × *A. variegatum* subsp. *variegatum*.

Diagnostic characters: inflorescence axis, pedicels and tepals outside pilose.

Iconography: Mitka (2003: 176, fig. 9E).

Aconitum ×cammarum L. em. Fries, Sp. Pl. 1 (ed. 2): 751. 1762 em. Nov. Flor. Suec. (ed. 2): 171. 1828

Type citation: "Austria, Styria".

Typus: not designated. Conflict of original material has been revealed (The Linnaean Plant Name Typification Project, 2006).

Synonyms: *A. intermedium* DC., *A. stoerkianum* Rchb.

Hybrid origin: *A. sect. A.* (*A. napellus* ?) × *A. variegatum*.

Distribution: Europe (garden plant, sometimes naturalized).

Diagnostic characters: hood glabrous; tepals deep blue (fo. *cammarum*) or with white strips (fo. *bicolor*); pedicels above bracteoles sparsely curved pubescent, below bracteoles glabrescent; carpels glabrous; filaments pilose.

Iconography: Reichenbach (1827: t. 71, *A. stoerkianum*); Mitka (2003: 173, fig. 6F), Novikoff and Mitka (2011: 44).

Literature: Mitka (2003).

Lower taxa:

a) var. ***cammarum*** fo. ***cammarum***

Diagnostic characters: tepals with monotonic, usually deep blue, color.

b) var. ***cammarum*** fo. ***bicolor*** (Schult) DC., Syst. Nat. [Candolle] 1: 372. 1817

Type citation: none.

Typus: not designated.

Synonyms: none.

Diagnostic characters: tepals variegate, with white and blue stripes.

Aconitum ×exaltatum Bernh. ex Rchb., Ill. Spec. Acon.: Tab. LXXII. 1827

Type citation: "In montium Sudetorum Aupagrund, *Funk!*".

Type specimens: not designated, probably BM 000613639.

Synonyms: *A. exaltatum* var. *hamatum* Rchb. [= *A. hamatum* (Rchb.) Gáyer], *A. speciosum* Otto ex Rchb.

Hybrid origin: *A. plicatum* × *A. variegatum*.

Distribution: Sudetes, Bohemian Massif, Bavarian Alps.

Diagnostic characters: inflorescence axis and hoods outside glabrous; hoods rounded-conic or slightly elongated; pedicels glabrous or glandular curved and crisped pubescent, stamens pilose, spur of nectaries backward bent or capitate; carpels mostly entirely sterile.

Iconography: Mitka (2003: 177, fig. 10F).

Literature: Mitka (2003).

Aconitum ×mielichhoferi Rchb., Uebers. Acon.: 29. 1819

Type citation: "Hab. in Alp. Carinth. Salisb. (in quodam certi harum regionum loco) *Mielichhoferi*".

Typus: not designated.

Synonym: *A. ×ronnigeri* Gáyer.

Hybrid origin: *A. degenerii* × *A. tauricum*.

Distribution: Eastern and Southeastern Alps.

Diagnostic characters: pedicels at least above the bracteoles densely glandular pilose; bracteoles filiform to spathulate; tepals outside sparsely to densely glandular pilose; hood about as broad as high; nectaries always pilose, their stalks distinctly bent, their spur capitate, always reaching the top of the hood; carpels 3, glabrous to densely hairy.

Literature: Starmüller (2001).

Aconitum ×schneebergense Gáyer, Magyar Bot. Lapok 8: 189. 1909

Type citation: "Hab. in Schneeberg Austriae Inf. (*Portenschlag* - W, duo exempla: unum cum nota: *A. neomontanum*, Schneeberg, alterum: Austria Inf.)".

Typus: not designated.

Synonyms: *A. ×algovicense* Gáyer, *A. laxum* Gáyer, *A. ×virgatum* Rchb. ex Gáyer.

Hybrid origin: *A. napellus* × *A. variegatum*.

Distribution: Alps.

Diagnostic characters: pedicels above the bracteoles ± densely crimped eglandular pubescent, below the bracteoles ± glabrescent; tepals outside glabrous; hood about as high as broad; nectaries mostly glabrous, hardly curved, with a capitate to hooked spur, always reaching the top of the hood; filaments always hairy; carpels 3–4, glabrous or pilose on the suture.

Literature: Starmüller (2001).

Hybrid formulae:

Aconitum napellus* × *A. pilipes – hood rounded, pedicels above bracteoles glandular pubescent and below bracteoles eglandular pubescent; spur of nectaries backward bent or capitate, carpels glabrous or entirely pubescent, sterile.

Aconitum pilipes* × *A. tauricum – hood rounded; pedicels glandular pubescent, spur of nectaries capitates; carpels glabrous or slightly pubescent.

Aconitum superbum* × *A. variegatum – more studies are required.

Aconitum superbum* × *A. vitosanum – more studies are required.

4. – Sect. *Angustifolium* (Seitz) Rottensteiner, stat. nov.

Basionym: *A. subsect. Angustifolium* Seitz, Feddes Repert. 80 (1): 60. 1969

Type species: *A. angustifolium* Bernh. ex Rchb.

Diagnosis characters: seeds tricorn, pyramidal, membranous 6–8 lamellae, root napiform.

Description: stem up to 60 cm, glabrous; leaves with 5–7 lobes, narrowly dissected; leaf segments linear to lanceolate, up to 4 (–6) mm wide; inflorescence compact, pyramidal or slightly ramified; pedicels firm, with tri-lobed or entire linear bracteoles; flowers violet; hood from crescent-shape to hemispherical, glabrous or sparsely pubescent; nectaries hardly bent with ± capitate spurs, always reaching the top of the hood; filaments glabrous; carpels 2–3, glabrous or pilose on the dorsal or ventral side.

Aconitum angustifolium Bernh. ex Rchb., Mon. Aconit.: 95, t. 15, fig. 2. 1821

Type citation: "In Styriae valle Wochinensi: *Bernhardi!* (ibid.) circa lacum, in altitude circiter 2000 pendum: *de Vest!* In Sibiria: *Pallas!*".

Typus: iconotype: Reichenbach (1820): Monogr. Aconit.: 24 (Seitz 1969).

Synonyms: *A. albicans* Host, *A. napellus* subsp. *angustifolium* (Bernh. ex Rchb.) Gáyer, *A. tenuifolium* Host.

Distribution: Julian Alps and Prealps in Italy and Slovenia.

Diagnostic characters: uppermost leaf segment 1–3 (–4) mm wide; flowers light blue to whitish; hood often ± triangular, about as high as broad; inflorescence axis, pedicels and tepals outside glabrous (fo. *angustifolium*) or ± pubescent (fo. *carniolicum*); nectaries hardly bent, their spur ± capitate, always reaching the top of the hood; filaments glabrous; carpels 2–3, glabrous or pilose on the dorsal or ventral side.

Iconography: Reichenbach (1821: t. 15, fig. 2).

Literature: Gáyer (1909); Seitz (1969); Nikolić (1994); Martinčić (1999); Aeschimann et al. (2004).

Hybrids: unknown.

Lower taxa:

a) fo. *angustifolium*

Diagnostic characters: inflorescence axis, pedicels and tepals outside glabrous.

b) fo. *carniolicum* Gáyer, Magyar. Bot. Lap. 8: 184. 1909

Type citation: "Krain, Küstenland, Steiermark. Auf der Zhaun u. Krushza (*Fleischmann* – W); Stiria, Auen a. d. Mur (*Maly* – W. loco ergo incerto)" (Gáyer 1909).

Typus: not designated.

Synonyms: none.

Diagnostic characters: inflorescence axis, pedicels and tepals outside ± pubescent.

5. – Sect. *Euchylodea* Rchb.

Type species: *A. kusnezoffii* Rchb.

Diagnostic characters: nectary lobes inflated; spurs thick, long and hamate curved or short saccate; stipes slightly incurved or nearly erect.

Description: root obconical or elongated, 2–5 cm; stem up to 200–300 cm long, usually intensively branched on the level of inflorescence; caudine leaves with 5 lobes with narrow lanceolate segments; pedicel densely pubescent; bracteoles acute ovate; tepals lilac-blue, outside pubescent; hood hemispherical to slightly elongated; carpels 5, mostly glabrous.

5A. – Ser. *Japonica* (Nakai) Kadota

Type species: *A. japonicum* Thunb.

Diagnostic characters: leaf blades pentagonal to roundish pentagonal, sometimes roundish reniform; lateral lobes shallowly to medially 2-parted or sometimes deeply incised.

Aconitum carmichaelii Debeaux, Actes Soc. linn. Bordeaux 33: 87. 1879

Type citation: "La region montagneuse du Tché-fou, près de Bambou Temple (très-probablement la Pagode neuve) où il a été découvert par le Dr. Carmichael en 1875".

Typus: not designated.

Synonyms: *A. bodinieri* H. Lév. & Vaniot, *A. chinense* Paxton, *A. fischeri* auct., *A. lushanense* Migo, *A. wilsonii* Stapf ex Veitch.

Distribution: China, Vietnam; neophyte in Europe: Scotland, France, Germany, Austria.

Diagnostic characters: Root obconical, 2–4 cm long, 1–1.6 cm broad; stem rigid, upright, 200–300 cm; caudine leaves pentagonal, ± pubescent on the lower surface; inflorescence with many side inflorescences; pedicel densely pubescent; bracteoles acute ovate; tepals lilac-blue, outside pubescent; hood about as high as broad; nectaries glabrous, with an upright claw and a recurved spur; filaments dentate, glabrous; carpels 5, mostly glabrous.

Iconography: Wang et al. (2003).

Literature: Li and Kadota (2001); Lin et al. (2017).

Hybrids: unknown.

ACKNOWLEDGMENTS

The work has been supported by a KBN/NCN grant N303 814440 given to JM and funds of the Institute of Botany of the Jagiellonian University in Kraków.

REFERENCES

- Aeschimann D, Lauber K, Moser DM, Theurillat JP. 2004. *Flora alpina*. Vol. 1. Bern, Stuttgart, Wien.
- Bauhin C. 1671. *Pinax theatri botanici*. Basileae: Impensis Joannis Regis.
- Boroń P, Wróblewska A, Binkiewicz B, Mitka J. 2020. Phylogeny of *Aconitum* subgenus *Aconitum* in Europe. *Acta Soc Bot Pol.* 89: Article 8933. <https://doi.org/10.5586/asbp.8933>
- Bosch M, Simon J, López-Pujol J, Blanché C. 2016. DCDB: an updated online database of chromosome numbers of tribe Delphinieae (Ranunculaceae). *Flora Mediterranea*. 26: 191–201. <http://dx.doi.org/10.7320/FIMedit26.191>
- Cappelletti EM., Poldini L. 1984. Seed morphology in some European aconites (*Aconitum*, *Ranunculaceae*). *Plant Syst Evol.* 145: 193–201. <https://doi.org/10.1007/BF00983948>
- Clapham AR, Tutin TE, Warburg EF. 1957. *Flora of the British Isles, Illustrations* 1. Cambridge.
- Clusius C. 1601. *Rariorvm plantarvm historia. Quae accesserint, proxima pagina docebit*. Vol. 2. Antverpiae: Ex officina Plantiniana Apud Ioannem Moretum.
- Fritsch K. 1895. Beiträge zur Flora der Balkanhalbinsel, mit besonderer Berücksichtigung von Serbien, Dritter Theil. *Verh Zool-Bot Ges Wien.* 45: 367–382.
- Gáyer G. 1906. Die toxicoiden *Aconitum-Arten* in Ungarn. *Magyar Bot Lap.* 5: 122–137.
- Gáyer G. 1909. Vorarbeiten zu einer Monographie der europäischen *Aconitum-Arten*. *Magyar Bot Lap.* 9: 114–206.
- Gáyer G. 1911. De *Aconitis quibusdam alpinis*. *Magyar Bot Lap.* 10: 194–196.
- Gáyer G. 1912. *Aconitum* In: Hegi G, ed. *Illustrierte Flora von Mittel-Europa*. Mit besonderer Berücksichtigung von Deutschland, Oesterreich und der Schweiz. Zum Gebrauche in den Schulen und zum Selbstunterricht. Band III. Dicotyledones (I. Teil). München: J.F. Lehmann Verlag; p. 492–507.

- Götz E. 1967. Die *Aconitum variegatum*-Gruppe und ihre Bastarde in Europa. Feddes Repert. 76 (1–2): 1–62. <https://doi.org/10.1002/fedr.19670760102>
- Hayek A. 1917. Zur Kenntnis der Flora des Berges Žlep bei Ipek. Ann Naturhist Museums Wien. 31: 65–76. www.jstor.org/stable/41768192
- Hong Y, Luo Y, Gao Q, Ren C, Yuan Q, Yang Q-E. 2017. Phylogeny and reclassification of *Aconitum* subgenus *Lycoctonum* (Ranunculaceae). Plos One. 12(1): e0171038. <https://doi.org/10.1371/journal.pone.0171038>
- Ilnicki T, Mitka J. 2009. Chromosome numbers in *Aconitum* sect. *Aconitum* (Ranunculaceae) from the Carpathians. Caryologia. 62: 198–203. <https://doi.org/10.1080/00087114.2004.10589685>
- Ilnicki T, Mitka J. 2011. Chromosome numbers in *Aconitum* sect. *Cammarum* (Ranunculaceae) from the Carpathians. Caryologia. 64: 446–452. <https://doi.org/10.1080/00087114.2011.10589812>
- Jabbour F, De Craene LR, Nadot S, Damerval C. 2009. Establishment of zygomorphy on an ontogenetic spiral and evolution of perianth in the tribe Delphinieae (Ranunculaceae). Annals of Botany. 104: 809–822. <https://doi.org/10.1093/aob/mcp162>
- Jabbour F, Renner SR. 2011a. *Consolida* and *Aconitella* are an annual clade of *Delphinium* (Ranunculaceae) that diversified in the Mediterranean basin and the Irano-Turanian region. Taxon. 60: 1029–1040. <https://doi.org/10.1002/TAX.604007>
- Jabbour F, Renner SR. 2011b. A phylogeny of Delphinieae (Ranunculaceae) shows that *Aconitum* is nested within *Delphinium* and that the Late Miocene transitions to long life cycles in the Himalayas and Southwest China coincide with bursts in diversification. Mol Phyl Evol. 62: 928–942. <https://doi.org/10.1016/j.ympev.2011.12.005>
- Jabbour F, Cossard G, Le Guilloux M, Sannier J, Nadot S, Damerval C. 2014. Specific duplication and dorsoventrally asymmetric expression patterns of *Cycloidea*-like genes in zygomorphic species of Ranunculaceae. Plos One. 9(4): e95727. <https://doi.org/10.1371/journal.pone.0095727>
- Jacquin NJ. 1776. Florae Austriacae, sive plantarum selectarum in Austriae archiducatu sponte crescentium, icones, ad vivum coloratae, et descriptionibus, ac synonymis illustratae. Vol. 4. Viennae Austriae: Typis Leopoldi Joannis Kaliwoda, Aulae imperialis typographi.
- Jeanmonod D, Gamisans J. 2007. Flora Corsica. Aix-en-Provence: Edisud.
- Joachimiak A, Ilnicki T, Mitka J. 1999. Karyological studies on *Aconitum lasiocarpum* (Rchb.) Gáyer (Ranunculaceae). Acta Biol Cracov, Ser Bot. 41: 205–211.
- Kadota Y. 1981. A taxonomic study of *Aconitum* (Ranunculaceae) of the Akaishi Mountain Range in Central Japan. Bull Nat Sci Mus B (Tokyo). 7: 91–112.
- Kadota Y. 1987. A revision of *Aconitum* subgenus *Aconitum* (Ranunculaceae) in East Asia. Utsunomiya: Sanwa Shoyaku Company, Ltd.
- Keener CS, Reveal JL, Dutton E, Ziman S. 1999. A list of suprageneric names in Ranunculaceae (Magnoliophyta). Taxon. 48: 497–506. <https://doi.org/10.2307/1224562>
- Kirschner J, Kirschnerova L, Štěpánek J. 2007. Generally accepted plant names based on material from the Czech Republic and published in 1753–1820. Preslia. 79: 323–365.
- Kita Y, Ito M. 2000. Nuclear ribosomal ITS sequences and phylogeny of East Asian *Aconitum* subg. *Aconitum* (Ranunculaceae), with special reference to extensive polymorphism in individual plants. Plant Syst Evol. 225: 1–13. <https://doi.org/10.1007/BF00985455>
- Kita Y, Ueda K, Kadota Y. 1995. Molecular phylogeny and evolution of the Asian *Aconitum* subg. *Aconitum* (Ranunculaceae). J Plant Res. 108: 429–442.
- Li LQ, Kadota Y. 2001. *Aconitum* L. In: Zhyengi W, Raven PH, Deyuan H, eds. Flora of China. Vol. 6. Caryophyllaceae through Lardizabalaceae. Beijing: Science Press; St. Louis: Missouri Botanical Garden; pp. 149–222. Available online from http://www.efloras.org/flo-rataxon.aspx?flora_id=2&taxon_id=100300
- Lin Q, Yang ZR, Lin Y. 2017. Type specimens in China national herbarium (PE), Volume 8, Angiospermae (5). Henan Science and Technology Press.
- Luferov AN. 2000. Konspekt kavkazskih vidov *Aconitum* (Ranunculaceae). Bot Zhur. 85: 87–96. (in Russian).
- Luo Y, Zhang FM, Young Q. 2005. Phylogeny of *Aconitum* subgenus *Aconitum* (Ranunculaceae) inferred from ITS sequences. Plant Syst Evol. 252: 11–25. <https://doi.org/10.1007/s00606-004-0257-5>
- Martinčič A. 1999. Mala flora Slovenije. Ed. 3. Ljubljana.
- Mattioli PA. 1563. New Kreüterbuch: mit den allerschönsten und artlichsten Figuren aller Gewechss, dergleichen vormals in keiner Sprach nie an Tag kommen. Gedruckt zu Prag: Durch Georgen Melantrich von Auentin, auff sein und Vincenti Valgriss Buchdruckers zu Venedig uncosten.
- Molero J, Puig A. 1990. Seed morphology of Iberian species of the genus *Aconitum*. Collect Bot (Barcelona). 19: 111–127. <https://doi.org/10.3989/COLLECT-BOT.1990.V19.120>
- Mitka J. 2000. Systematyka *Aconitum* subg. *Aconitum* w Karpatach Wschodnich [Systematics of *Aconitum* subg. *Aconitum* in the Eastern Carpathians]. Roczn Biesczadzkie. 9: 79–116. (in Polish).

- Mitka J. 2002. Phenetic and geographic pattern of *Aconitum* sect. *Napellus* (Ranunculaceae) in the Eastern Carpathians – a numerical approach. *Acta Soc Bot Pol.* 71: 35–48. <https://doi.org/10.5586/asbp.2002.005>
- Mitka J. 2003. The genus *Aconitum* in Poland and adjacent countries – a phenetic-geographic study. Kraków: Institute of Botany, Jagiellonian University.
- Mitka J., Boroń P., Sutkowska A. 2013. Holocene history of *Aconitum* in the Polish Western Carpathians and adjacent regions: long-distance migration or cryptic refugia. *Mod Phytomorphol.* 3: 9–18. <https://doi.org/10.5281/ZENODO.161587>
- Mitka J., Binkiewicz B., Stachurska-Swakoń A., Novikov A., Rottensteiner W. 2017. A synopsis of the genus *Aconitum* subg. *Aconitum* in Europe. *Studia Univ Babeş-Bolyai, Sp Iss.* 166–167.
- Mitka J., Boroń P., Wróblewska A., Bąba W. 2015. AFLP analysis reveals infraspecific phylogenetic relationships and population genetic structure of two species of *Aconitum* in Central Europe. *Acta Soc Bot Pol.* 84: 267–276. <https://doi.org/10.5586/asbp.2015.012>
- Mitka J., Starmühler W. 2000. Phenetic variability of *Aconitum lasiocarpum* (Rchb.) Gáyer (Ranunculaceae): extension of taxonomic and geographic borders. *Acta Soc Bot Pol.* 69: 145–155. <https://doi.org/10.5586/asbp.2000.020>
- Mitka J., Sutkowska A., Ilnicki T., Joachimiak A. 2007. Reticulate evolution of high-alpine *Aconitum* (Ranunculaceae) in the Eastern Sudetes and Western Carpathians (Central Europe). *Acta Biol Cracov Ser Bot.* 49: 15–26.
- Molero J., Blanché C. 1984. A propósito de los géneros *Aconitum* L. y *Consolida* (DC.) S.F. Gray en la península Ibérica. *Anal Jard Bot Madrid.* 41: 211–218.
- Molero J., Blanché C. 1986. *Aconitum* L. In: Castroviégo S., ed. *Flora Iberica*, Vol. 1. Lycopodiaceae-Papaveraceae. Madrid: Real Jardín Botánico, C.S.I.C. Pp. 233–242.
- Mucher W. 1990. *Aconitum paniculatum* subsp. *paniculatum* var. *turrachense* var. nova (Ranunculaceae). *Phyton (Horn).* 30 (1): 83–88.
- Mucher W. 1991a. Die Hybriden zwischen den Subspecies von *Aconitum napellus* L. In: Hafellner J., ed. Österreichisches Botanikertreffen in Graz, 1991, Kurzfassungen der Beiträge; p. 49.
- Mucher W. 1991b. Der Bunte Eisenhut, *Aconitum variegatum* L. (Ranunculaceae), in der Steiermark. *Mit Naturwiss Ver Steiermark* 121: 195–198.
- Mucher W. 1991c. *Aconitum napellus* subsp. *lobelii* subsp. nova und *A. napellus* subsp. *koelleanum* comb. nova. *Phyton (Horn).* 31 (1): 129–135.
- Mucher W. 1992. *Aconitum variegatum* ssp. *variegatum* var. *stiriacum* var. nova (Ranunculaceae). *Verh Zool-Bot Ges Österreich.* 129: 143–151.
- Mucher W. 1993a. Systematics and chorology of *Aconitum* ser. *Toxicum* in Europe. *Phyton (Austria).* 33: 51–76.
- Mucher W. 1993b. Die Gattung *Aconitum* in Kärnten. *Carinthia II.* 183/103 (2): 519–527.
- Müller K. 2005. SeqState-primer design and sequence statistics for phylogenetic data sets. *Appl Bioinformatics.* 4: 65–69. <https://doi.org/10.2165/00822942-200504010-00008>
- Nikolić T. 1994. *Flora Croatica. Index Flora Croaticae 1.* Zagreb: Croatian Natural History Museum.
- Novikoff A., Mitka J. 2011. Taxonomy and ecology of the genus *Aconitum* in the Ukrainian Carpathians. *Wulfenia.* 18: 37–61.
- Novikoff A., Mitka J. 2015. Anatomy of stem-node-leaf continuum in *Aconitum* (Ranunculaceae) in the Eastern Carpathians. *Nordic J Bot.* 33: 633–640. <https://doi.org/10.1111/njb.00893>
- Penev I., Simeonovski M. 1970. *Aconitum* L. In: Jordanov D., Kožuharov S., eds. *Flora Reipublicae Popularis Bulgariae*, Vol. 4. Serdicae; pp. 54–61.
- Reichenbach HGL. 1819. Uebersicht der Gattung *Aconitum*. Regensburg.
- Reichenbach HGL. 1820–1821. Monographia generis *Aconiti*. Lipsiae.
- Reichenbach HGL. 1823–1827. Illustratio specierum *Aconi* generis. Lipsiae.
- Reichenbach HGL. 1838–1840. Icones flora germanicae et helveticae. Lipsiae.
- Ross-Craig S. 1948. *Drawings of British Plants*. London.
- Rottensteiner WK. 2018. *Aconitum* subsect. *Burnatii* subseccio nova. *Botanica Serbica.* 42 (suppl. 1): 40.
- Rouy G. 1884. Diagnoses d'espèces nouvelles pour la flore de la péninsule Ibérique. *Le Naturaliste. Journal des échanges et des nouvelles* 6: 405–406.
- Seitz W. 1969. Die Taxonomie der *Aconitum napellus*-Gruppe in Europa. *Feddes Repert.* 80: 1–76. <https://doi.org/10.1002/fedr.19690800102>
- Seitz W. 1970. Nachtrag zur Nomenklatur der *Aconitum napellus*-Gruppe in Europa. *Taxon.* 19: 904–905. <https://doi.org/10.2307/1218306>
- Seitz W., Zinsmeister HD., Abicht M. 1972. Beitrag zur Systematik der Gattung *Aconitum* in Europa. *Bot Jahrb Syst.* 92: 490–507.
- Simon J., Bosch M., Molero J., Blanché C. 2001. A conspect of chromosome numbers in tribe *Delphinieae* (Ranunculaceae). Available online from https://www.researchgate.net/publication/313862132_A_conspect_of_chromosome_numbers_in_tribe_Delphinieae_Ranunculaceae

- Skalický V. 1982. Notulae systematicae, diagnosticae and nomenclatoriae ad *Aconitum* generis investigatio- nem pertinentes. *Preslia*. 54: 115–122.
- Skalický V. 1990. Rod *Aconitum* na Československu. *Zprávy Českoslov Bot Spol.* 25: 1–27. (in Czech).
- Soó R. 1972. Systematisch-nomenklatörische Bemerkun- gen zur Flora Mitteleuropas mit Beziehungen zur südosteuropäischen Flora. *Feddes Repert.* 83(3): 129– 212. <https://doi.org/10.1002/fedr.19720830302>
- Stapf O. 1926. *Aconitum anglicum*. *Curtis's Botanical Magazine*. 151: tab. 9088.
- Starmühler W. 1996a. Beitrag zur Kenntnis des *Aconitum pilipes* (Ranunculaceae) und seiner Hybriden. *Willde- nowia*. 26: 169–182.
- Starmühler W. 1996b. Neuste Erkenntnisse über die Gat- tung *Aconitum* in Siebenbürgen. *Stapfia*. 45: 7–45.
- Starmühler W. 1996c. Systematics and chorology of the genus *Aconitum* in the Bela krajina region (Slovenia). *Hladnikia*. 6: 5–16.
- Starmühler W. 1996d. Ranunculaceae subtrib. Delphinii- nae exsiccatae, Fasc. 1 (Nr. 1–20). *Fritschiana* (Graz). 7: 1–10.
- Starmühler W. 1996e. *Aconita rarissima selecta*. *Fritschiana* (Graz). 7: 11–39.
- Starmühler W. 1997a. Ranunculaceae subtrib. Delphi- niinae exsiccatae, Fasc. 2 (Nr. 21–40). *Fritschiana* (Graz). 10: 1–11.
- Starmühler W. 1997b. Typification of names in the genus *Aconitum*. In: Wisskirchen R, Christensen KI, Dick- oré WB, Gottschlich G, Hanelt P, van Raamsdonk LWD, Schmidt PA, Starmühler W, Wisskirchen R. *Notulae ad Floram Germanicam I*. *Feddes Repert.* 108 (1–2): 102–103. <https://doi.org/10.1002/fedr.19971080108>
- Starmühler W. 1998. Eine Übersicht der Arten aus der Gattung *Aconitum*. *Schweizer Staudengärten*. 27: 24–47.
- Starmühler W. 1999. *Aconita rarissima selecta* (1999). *Fritschiana* (Graz). 18: 11–38.
- Starmühler W. 2000. Fortschritte in der Systematik der Gattung *Aconitum* in Siebenbürgen. In: Helt- mann H, von Killyen H, eds. *Naturwissenschaftliche Forschung in Siebenbürgen VI*, Beiträge zur Geogra- phie, Botanik, Zoologie und Paläontologie. Sieben- bürgisches Archiv. 36: 9–24.
- Starmühler W. 2001. Die Gattung *Aconitum* in Bayern. *Ber Bayer Bot Ges.* 71: 99–118.
- Starmühler W. 2002. *Aconita rarissima selecta* (2002). *Fritschiana* (Graz). 30: 13–42.
- Starmühler W. 2010. Eisenhut (*Aconitum*). In: Speta E, Rákosi L, eds. *Wildpflanzen Siebenbürgens*. Freistadt: Plöchl Druck GmbH; pp. 188–189, 438–443, 602.
- Starmühler W, Mitka J. 2001. Systematics and chorol- ogy of *Aconitum* sect. *Napellus* and its hybrids in the Northern Carpathians and Forest Carpathians. *Thias- zia – J Bot* (Košice). 10: 115–136.
- Sutkowska A, Boroń P, Mitka J. 2013. Natural hybrid zone of *Aconitum* species in the Western Carpathians: Linnaean taxonomy and ISSR fingerprinting. *Acta Biol Cracoviensia Ser Bot.* 55: 114–126. <https://doi.org/10.2478/abcsb-2013-00015>
- Sutkowska A, Boroń P, Warzecha T, Dębowski J, Mitka J. 2017a. Hybridization and introgression among three *Aconitum* (Ranunculaceae) species of different ploidy levels in the Tatra Mountains (Western Carpathians). *Plant Species Biology*: <https://doi.org/10.1111/1442-1984.12162>
- Sutkowska A, Warzecha T, Mitka J. 2017b. Genetic varia- tion of *Aconitum* sect. *Aconitum* at a macrogeographi- cal scale in the Carpathians. *Pol J Ecol.* 65: 57–68. <https://doi.org/10.3161/15052249PJE2017.65.1.006>
- Tamura M. 1993. *Ranunculaceae*. In: K. Kubitzky (ed.) *The families and genera of vascular plants. 2. Flow- ering plants. Dicotyledons. Magnoliid, Hamamelid and Caryophyllid families*. Berlin: Springer Verl; pp. 563–583.
- The Linnaean Plant Name Typification Project. 2006. *Aco- nitum cammarum* Linnaeus. [accessed on 03.10.2020]. Available online from [https://doi.org/10.2307/1223721](https://www.nhm.ac.uk/our- science/data/linnaean-typification/search/detail.dsml?ID=6700&listPageURL=list%2edsml%3fVarqtype%3d starts%2bwith%26CVarqtype%3dstarts%2bwith%26C Genusqtype%3dstarts%2bwith%26CSpeciesqtype%3d starts%2bwith%26Species%3dcamarum%26sort%3d Genus%252cSpecies%26Speciesqtype%3dstarts%2bwit h%26Genus%3dAconitum%26Genusqtype%3dstarts% 2bwith%26CSpqtype%3dstarts%2bwith%</p>
<p>Turland NJ, Barrie FR. 2001. Family name listings modi- fied in Appendix IIB of the Saint Louis Code. <i>Taxon</i>. 50 (3): 897–903. <a href=)
- Tutin TG, Burges NA, Chater AO, Edmondson JR, Hey- wood VH, Moore DM, Valentine DH, Walters SM, Webb DA. 1993. *Flora Europaea*. Vol. 1: Psilotaceae to Plantanaceae. 2nd ed. New York: Cambridge Uni- versity Press.
- Tzvelev NN. 1996. On some genera of the family Ranunculaceae in the Eastern Europe. *Bot Zhurn.* 81: 112– 122. (in Russian).
- Utelli A-B, Roy BA, Baltisberger M. 2000. Molecular and morphological analyses of European *Aconitum* spe- cies (Ranunculaceae). *Plant Syst Evol.* 224: 195–212. <https://doi.org/10.1007/BF00986343>
- Wacławská-Ćwiertnia K, Mitka J. 2016. Typification of Zapałowicz's names in *Aconitum* section *Aco-*

- nitum*. PhytoKeys. 58: 119–126. <https://dx.doi.org/10.3897%2Fphytokeys.58.7110>
- Wang W, Liu Y, Yu SX, Gao TG, Chen ZD. 2013. *Gymnaconitum*, a new genus of Ranunculaceae endemic to the Qinghai-Tibetan Plateau. *Taxon*. 62: 713–722. <https://doi.org/10.12705/624.10>
- Wang W, Fu D, Li LQ, Bartholomew B, Brach AR, Dutton BE, Gilbert MG, Kadota Y, Robinson OR, Tamura M, Warnock MJ, Zhu G, Ziman SN. 2003. Ranunculaceae. In: Wu ZY, Zhu G, eds. Flora of China illustrations. Vol 6. Caryophyllaceae through Lardizabalaceae. Beijing: Science Press and St. Louis: Missouri Botanical Garden Press.
- Wissjulina E. 1939. Novyj vid roda *Aconitum* (Tourn.) L. flory USSR. Zhurn Inst Botan AN URSR. 21–22: 253–254. (in Ukrainian).
- Wojciechowska B, Makulec J. 1969. Seed morphology and anatomy of some *Aconitum* species. Monographiae Botanicae. 29: 137–163.
- Yang QE. 1996. A karyotype study of 15 species in the tribe *Delphineae* (Ranunculaceae) from China. *Acta Phytotaxonomica Sinica*. 34 (1): 39–47.
- Yu M, Yang YX, Shu XY, Huang J, Hou DB. 2016. *Aconitum carmichaelii* Debeaux, cultivated as a medicinal plant in western China. *Genet Resour Crop Evol*. 63: 919–924. <https://doi.org/10.1007/s10722-016-0398-8>
- Zapałowicz H. 1908a. Krytyczny Przegląd Roślinności Galicji (części: XII, XIII) [A critical review of the flora of Galicia (parts: XII, XIII)]. Rozprawy Wydziału Matematyczno-Przyrodniczego Akademii Umiejętności. Dział B. Nauki Biologiczne. 48 (ser. III, t. 8): 41–90, 187–256.
- Zapałowicz H. 1908b. Conspectus floriae Galiciae criticae. Vol. 2. Cracoviae.
- Zajac M, Zajac A. 2009. The geographical elements of native flora of Poland. Kraków: Jagiellonian University.
- Zhao D, Shi Y, Zhu X, Liu L, Ji P, Long C, Shen Y, Kennelly EJ. 2017. Identification of potential biomarkers from *Aconitum carmichaelii*, a traditional Chinese medicine, using a metabolomic approach. *Planta Medica*. 84(06/07): 434–441. <https://doi.org/10.1055/s-0043-121708>
- Zieliński R. 1982a. An electrophoretic and cytological study of hybridization between *Aconitum napellus* subsp. *skerisorae* ($2n = 32$) and *A. variegatum* ($2n = 16$). I. Electrophoretic evidence. *Acta Soc Bot Pol*. 51: 453–464.
- Zieliński R. 1982b. An electrophoretic and cytological study of hybridization between *Aconitum napellus* subsp. *skerisorae* ($2n = 32$) and *A. variegatum* ($2n = 16$). II. Cytological evidence. *Acta Soc Bot Pol*. 51: 465–471.
- Ziman SM, Schiyan NM, Bulakh OV. 2015. Types of the taxa of genus *Aconitum* (Ranunculaceae) described from Ukraine. *Ukr Bot Zhurn*. 72 (4): 325–333 (in Ukrainian). <https://doi.org/10.15407/ukrbotj72.04.325>

APPENDIX

CHECKLIST OF ACONITUM SUBG. ACONITUM IN EUROPE

- A. *xacuminatum* Rchb.** [*A. degenii* × *A. napellus*]
- A. *acutum* Rchb.** [*A. tauricum* × *A. variegatum*]
- A. *adriaticum* Gáyer** = **A. *superbum* Fritsch**
- A. *albicans* Host** = **A. *angustifolium* Bernh.** ex Rchb.
- A. *algoviense* Gáyer** = **A. *xschneebergense* Gáyer** [*A. napellus* × *A. variegatum*]
- A. *amoenum* Rchb.** = **A. *plicatum* Köhler** ex Rchb.
- A. *anglicum* Stapf**
- A. *angustifolium* Bernh.** ex Rchb.
- fo. ***angustifolium***
 - fo. ***carnolicum* Gáyer**
- A. *xaquilonare* A.Kern.** ex Gáyer [*A. variegatum* × *A. vitosanum*]
- A. *xaustriacum* Starm.** [*A. pilipes* × *A. variegatum*]
- A. *autumnale* Clus.** ex Rchb. = **A. *napellus* L.**
- A. *adriaticum* Gáyer** = **A. *superbum* Fritsch**
- A. *xbartokianum* Starm.** [*A. toxicum* × *A. variegatum*]
- nothosubsp. ***bartokianum* Starm.** [*A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *nasutum*]
 - nothosubsp. ***rapaicsianum* Starm.** [*A. toxicum* subsp. *crispulum* × *A. variegatum* subsp. *nasutum*]
 - nothosubsp. ***sooanum* Starm.** [*A. toxicum* subsp. *toxicum* × *A. variegatum* subsp. *variegatum*]
- A. *xbavaricum* Starm.** [*A. napellus* × *A. plicatum*]
- nothosubsp. ***bavaricum* [A. napellus subsp. *napellus* × *A. plicatum*]**
 - nothosubsp. ***lusenense* Starm.** [*A. napellus* subsp. *lobelii* × *A. plicatum*]
- A. *xberdai* Zapał.** [*A. firmum* × *A. variegatum*]
- nothosubsp. ***berdai* [A. firmum subsp. *firmum* × *A. variegatum* subsp. *variegatum*]**
 - nothosubsp. ***walasii* (Mitka)** Mitka [*A. firmum* subsp. *moravicum* × *A. variegatum* subsp. *variegatum*]
- A. *bernhardianum* Rchb., p.p.** = **A. *napellus* L.**
- A. *bernhardianum* Rchb., p.p.** = **A. *plicatum* Köhler** ex Rchb.
- A. *beskidense* (Zapał.)** Gáyer = **A. *lasiocarpum* Rchb.**
- subsp. ***kotulae* (Pawl.) Starm. & Mitka**
- A. *bodinieri* H.Lév. & Vaniot** = **A. *carmichaelii* Debeaux**
- A. *bosniacum* Beck** = **A. *toxicum* Rchb.**
- A. *bucovinense* Zapał.**

- fo. **bucovinense**
fo. **orthotricha** Gáyer
- A. burnatii** Gáyer
subsp. *pentheri* (Hayek) Jalas = **A. pentheri** Hayek
A. callibotryon Rchb. = **A. plicatum** Köhler ex Rchb.
- A. ×cammarum** L. em. Fries [*A. napellus* × *A. variegatum*]
var. *beskidene* Zapał. = **A. lasiocarpum** Rchb.
var. **cammarum**
 fo. **cammarum**
 fo. **bicolor** (Schult.) DC.
var. *pilipes* Rchb. = **A. pilipes** (Rchb.) Gáyer
- A. cammarum* Jacq.
var. *beskidene* Zapał. = **A. lasiocarpum** Rchb.
subsp. **kotulae** (Pawl.) Starm. & Mitka
var. *koscieliskanum* Zapał., p.p. = **A. lasiocarpum** Rchb. subsp. **kotulae** (Pawl.) Starm. & Mitka
- A. carmichaelii** Debeaux
- A. castellanum** (Molero & Blanché) Rottensteiner
- A. caucasicum* N.Busch
subsp. *pubiceps* (Rupr.) N.Busch
var. *tuscheticum* N.Busch = **A. ×tuscheticum** (N.Busch) N.Busch [A. *degenii* × A. *vitosanum*]
- A. cernuum* Rchb., non Wulfen
[var.] α *laxiflorum* Rchb. = **A. degenii** Gáyer subsp. **paniculatum** (Arcang.) Mucher var. *laxiflorum* (Rchb.) Mucher
- A. chinense* Paxton = **A. carmichaelii** Debeaux
- A. compactum* Rchb. = **A. napellus** L.
- A. clusianum** Rchb.
- A. corsicum** Gáyer
- A. ×czarnohorense** (Zapał.) Mitka [A. *bucovinense* × A. *xnanum*]
- A. dasycarpum* Schur = **A. lasiocarpum** Rchb.
- A. degenii** Gáyer
subsp. **degenii**
var. **degenii**
 fo. **degenii**
 fo. **craciunelense** Gáyer
var. **intermedium** (Zapał.) Mitka
nothosubsp. **gandogerii** Mucher [A. *degenii* subsp. *paniculatum* × A. *degenii* subsp. *valesiacum*]
nothosubsp. **lippertianum** Starm. [A. *degenii* subsp. *paniculatum* × A. *degenii* subsp. *rhaeticum*]
subsp. **paniculatum** (Arcang.) Mucher
var. **laxiflorum** (Rchb.) Mucher
var. **turrachense** (Mucher) Mucher
subsp. **rhaeticum** Starm.
subsp. **valesiacum** (Gáyer) Mucher
- A. delphinense* Gáyer = **A. burnatii** Gáyer
- A. diabolicum* Gáyer = **A. toxicum** Rchb.
- A. divergens* Pančić nom. illeg., p.p. = **A. pentheri** Hayek
- subsp. *burnatii* (Gáyer) W.Seitz = **A. burnatii** Gáyer
A. dolomiticum A.Kern. = **A. tauricum** Wulfen in Koelle
A. dolomiticum Hayek, p.p. = **A. tauricum** Wulfen in Koelle nothosubsp. **hayekianum** (Gáyer) Grinč. in Savulescu [*A. tauricum* subsp. *latemarensis* × *A. tauricum* subsp. *tauricum*]
A. dolomiticum Hayek, p.p. = **A. ×teppneri** Mucher ex Starm. nothosubsp. **kernerii** Starm. [*A. napellus* subsp. *napellus* × *A. tauricum* subsp. *latemarensis*]
A. ×dragulescuanum Mucher [*A. degenii* × *A. toxicum*]
nothosubsp. **dragulescuanum** [A. *degenii* subsp. *degenii* × A. *toxicum* subsp. *toxicum*]
nothosubsp. **rintescuanum** Starm. [A. *degenii* subsp. *degenii* × A. *toxicum* subsp. *crispulum*]
A. eminens Koch ex Rchb. = **A. napellus** L.
- A. ×exaltatum** Bernh. ex Rchb. [*A. plicatum* × *A. variegatum*]
var. *hamatum* Rchb. = **A. ×exaltatum** Bernh. ex Rchb. [*A. plicatum* × *A. variegatum*]
A. eustachyum Rchb. = **A. tauricum** Wulfen subsp. **tauricum** var. **eustachyum** (Rchb.) Starm.
- A. firmum** Rchb.
subsp. **firmum**
subsp. **fissurae** Nyár.
nothosubsp. **fussianum** Starm. [A. *firmum* subsp. *firmum* × A. *firmum* subsp. *fissurae*]
subsp. **maninense** (Skalický) Starm. = **A. maninense** (Skalický) Mitka
subsp. **moravicum** Skalický
var. *maninense* Skalický = **A. maninense** (Skalický) Mitka
nothosubsp. *paxii* Starm. in Starmühler & Mitka = **A. ×mariae** Rottensteiner, Mitka & Novikov nothosubsp. *paxii* (Starm.) Mitka, **comb. nov.** [A. *firmum* subsp. *moravicum* × A. *maninense*]
subsp. **skerisorae** (Gáyer) Starm.
nothosubsp. *zapalowiczzii* Starm. = **A. ×zapalowiczzii** (Starm.) Mitka [A. *firmum* × A. *×mariae* nothosubsp. *paxii*]
A. firmum Rchb. subsp. **firmum** × **A. firmum** Rchb.
subsp. **moravicum** Skalický
A. firmum Rchb. subsp. **firmum** × A. *maninense* (Skalický)
Mitka = **A. ×mariae** Rottensteiner, Mitka & Novikov nothospec. (hybr.) nov. [A. *firmum* s.str. × A. *maninense*]
A. fischeri Auct. = **A. carmichaelii** Debeaux
A. flerovii Steinberg in Komarov = **A. firmum** Rchb.
subsp. **fissurae** Nyár.
A. formosum Rchb. = **A. napellus** L. subsp. **formosum** (Rchb.) Gáyer in Hegi
A. ×gayeri Starm. [A. *degenii* × *A. lasiocarpum*]
A. gracile Rchb., p.p. = **A. variegatum** L. subsp. **variegatum**

subsp. *grosserratum* (Gáyer) Grinč.
 fo. *beskidense* Zapał. = **A. lasiocarpum** Rchb.
 subsp. *kotulae* (Pawł.) Starm. & Mitka
A. hamatum (Rchb.) Gáyer = **A. ×exaltatum** Bernh. ex Rchb. [*A. plicatum* × *A. variegatum*]
A. hayekianum Gáyer = **A. tauricum** Wulfen in Koelle
A. ×hebegynum DC. [*A. degenii* × *A. variegatum*]
A. ×hebegynum Auct. Fl. Carpathorum, non DC. = **A. degenii** Gáyer subsp. **degenii** var. **intermedium** (Zapał.) Mitka
A. hebegynum Auct., p.p. = **A. lasiocarpum** Rchb. subsp. **kotulae** (Pawł.) Starm. & Mitka
A. hians Rchb. = **A. plicatum** Köhler ex Rchb.
A. intermedium DC. = **A. ×cammarum** L. em. Fries [*A. napellus* × *A. variegatum*]
A. koehleri Rchb. = **A. plicatum** Köhler ex Rchb.
A. koelleanum Rchb., p.p. = **A. tauricum** Wulfen in Koelle
 var. *firmum* (Rchb.) Rchb. = **A. firmum** Rchb.
A. laetum Rchb. = **A. plicatum** Köhler ex Rchb.
A. lasiocarpum (Rchb.) Gáyer
 subsp. *kotulae* (Pawł.) Starm. & Mitka
 subsp. **lasiocarpum**
A. lasiocarpum (Rchb.) Gáyer × **A. toxicum** Rchb.
A. latemarensis Degen & Gáyer = **A. tauricum** Wulfen
 subsp. **latemarensis** (Degen & Gáyer) Starm. in W.Maurer
A. laxum Gáyer = **A. ×schneebergense** Gáyer [*A. napellus* × *A. variegatum*]
A. laxum Rchb. = **A. napellus** L.
A. ×lengyelii Gáyer = **A. ×berdaui** Zapał. [*A. firmum* × *A. variegatum*]
 nothosubsp. *walasii* Mitka. = **A. ×berdaui** Zapał.
 nothosubsp. *walasii* (Mitka) Mitka [*A. firmum* subsp. *moravicum* × *A. variegatum*]
A. linnaeanum Gáyer = **A. napellus** L.
A. lobelianum Host., non Rchb., p.p. = **A. napellus** L.
 subsp. **lobelii** Mucher
 fo. *brachytrichum* Gáyer. = **A. ×acuminatum** Rchb. [*A. degenii* × *A. napellus*]
A. lobelianum Rchb. = **A. napellus** L.
A. lobelianum (Rchb.) Host. in Gáyer, p.p. = **A. napellus**
 L. subsp. **lobelii** Mucher
 fo. *brachytrichum* Gáyer. = **A. ×acuminatum** Rchb. [*A. degenii* × *A. napellus*]
A. lobelianum Rouy-Fouc. = **A. corsicum** Gáyer
A. lushanense Migo = **A. carmichaelii** Debeaux
A. maninense (Skalický) Mitka
A. ×mariae Rottensteiner, Mitka & Novikov [*A. firmum* s.str. × *A. maninense*]
 nothosubsp. **mariae** [*A. firmum* subsp. *firmum* × *A. maninense*]

nothosubsp. **paxii** (Starm.) Mitka, **comb. nov.** [*A. firmum* subsp. *moravicum* × *A. maninense*]
A. meyeri Rchb. = **A. napellus** L.
A. ×mielichhoferi Rchb. [*A. degenii* × *A. tauricum*]
A. molle Rchb. = **A. degenii** Gáyer
A. multifidum Koch ex Rchb. = **A. plicatum** Köhler ex Rchb., *A. multifidum* Auct. Fl. Carpathorum, non Koch = **A. bucovinense** Zapał.
A. ×nanum (Baumg.) Simonk. [*A. bucovinense* × *A. firmum*]
A. napellus L.
 subsp. *angustifolium* (Bernh. ex Rchb.) Gáyer = **A. angustifolium** Bernh. ex Rchb.
 subsp. *burnatii* (Gáyer) J.-M. Tison. = **A. burnatii** Gáyer
 subsp. *castellanum* Molero and Blanché = **A. castellanum** (Molero & Blanché) Rottenst.
 subsp. *corsicum* (Gáyer) W. Seitz = **A. corsicum** Gáyer
 subsp. *eustachyum* (Rchb.) Gayer in Hegi = **A. tauricum** Wulfen subsp. **tauricum** var. **eustachyum** (Rchb.) Starm.
 subsp. *firmum* (Rchb.) Gáyer, p.p. = **A. firmum** Rchb.
 subsp. **firmum**
 var. *carpathicum* Maloch = **A. firmum** Rchb. subsp. **moravicum** Skalický
 subsp. **formosum** (Rchb.) Gáyer
 subsp. *hians* (Rchb.) Gáyer = **A. plicatum** Köhler ex Rchb.
 nothosubsp. **hinterhuberi** Starm. [*A. napellus* subsp. *formosum* × *A. napellus* subsp. *napellus*]
 subsp. *koelleanum* (Rchb.) Mucher = **A. tauricum** Wulfen in Koelle
 subsp. **lobelii** Mucher
 subsp. **lusitanicum** Rouy
 subsp. **napellus**
 var. *babiogorense* Zapał.
 fo. *babiogorense* = **A. firmum** Rchb.
 fo. *subfissum* Zapał. = **A. firmum** Rchb.
 var. *carpathicum* Zapał.
 fo. *carpathicum* = **A. firmum** Rchb.
 var. *compactum* Rapaics. = **A. corsicum** Gáyer
 var. *czarnohorense* Zapał. = **A. ×czarnohorense** (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *amoenum* Zapał. = **A. ×czarnohorense** (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *czarnohorense* Zapał. = **A. ×czarnohorense** (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *glabratum* Zapał. = **A. ×czarnohorense** (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *hoyerlanum* Zapał. = **A. ×czarnohorense** (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]

- fo. *nanum* Zapał., non Baumg. = *A. ×czarno-horense* (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *rodnense* Zapał. = *A. ×czarnohorense* (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *tenuisectum* Zapał. = *A. ×czarnohorense* (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 fo. *turkulense* Zapał. = *A. ×czarnohorense* (Zapał.) Mitka [*A. bucovinense* × *A. ×nanum*]
 [var.] γ *eustachyum* (Rchb.) Fiori = *A. tauricum* Wulfen subsp. *tauricum* var. *eustachyum* (Rchb.) Starm.
 var. *firmum* (Rchb.) Pawł. = *A. firmum* Rchb.
 [var.] *formosum* (Rchb.) Koch. = *A. napellus* L. subsp. *formosum* (Rchb.) Gáyer in Hegi
 [var.] β *hemisphaericum* Form
 fo. *formosum* (Rchb.) Beck. = *A. napellus* L. subsp. *formosum* (Rchb.) Gáyer in Hegi
 var. *nanum* Baumg. = *A. ×nanum* (Baumg.) Simonk. [*A. bucovinense* × *A. firmum*]
 var. *napellus*
 fo. *koelleanum* (Rchb.) Gürke in Richter-Gürke = *A. tauricum* Wulfen in Koelle
 var. *subtatrense* Zapał.
 fo. *abnorme* Zapał. = *A. firmum* Rchb.
 fo. *latisectum* Zapał. = *A. firmum* Rchb.
 fo. *subtatrense* = *A. firmum* Rchb.
 var. *tatrense* Zapał. = *A. firmum* Rchb.
 var. *tauricum* (Wulfen) Sér. = *A. tauricum* Wulfen in Koelle
 subsp. *napellus* sensu Seitz, 1969 = *A. anglicum* Stapf nothosubsp. *polatschekii* Mucher ex Starm. [*A. napellus* subsp. *formosum* × *A. napellus* subsp. *lobelii*] nothosubsp. *seitzii* Mucher ex Starm. [*A. napellus* subsp. *lobelii* × *A. napellus* subsp. *napellus*]
 subsp. *skerisorae* Seitz, non Gáyer = *A. firmum* Rchb. subsp. *skerisorae* (Gáyer) Starm.
 subsp. *superbum* (Fritsch) W.Seitz = *A. superbum* Fritsch
 subsp. *tauricum* (Wulfen) Gáyer = *A. tauricum* Wulfen in Koelle
***A. napellus* L. × *A. pilipes* (Rchb.) Gáyer**
A. nasutum Fischer ex Rchb., p.p. = *A. variegatum* L. subsp. *nasutum* (Fischer ex Rchb. em. Rupr.) Götz
 subsp. *tuscheticum* (N.Busch) A.N.Luferov = *A. ×tuscheticum* (N.Busch) N.Busch [*A. degenii* × *A. vitosanum*]
A. neomontanum Wulfen = *A. napellus* L.
A. neomontanum Auct., non Koelle = *A. toxicum* Rchb.
A. neubergense Clus. ex Rchb. = *A. napellus* L.
***A. nevadense* Uechtr. ex Gáyer**
A. occidentale Timb. Lagr. f. ex Gáyer = *A. nevadense* Uechtr. ex Gáyer
A. odotandrum Wissjul. = *A. ×berdaui* Zapał. [*A. firmum* × *A. variegatum*]
A. palmatifidum Rchb., p.p. = *A. firmum* Rchb.
 fo. *piliferum* Gáyer = *Aconitum ×mariae* Rottensteiner, Mitka & Novikov nothosubsp. *paxii* (Starm.) Mitka, **comb. nov.** [*A. firmum* subsp. *moravicum* × *A. maninense*]
A. paniculatum Lam., p.p. (*nom. inval.*) = *A. degenii* Gáyer subsp. ***degenii***
 subsp. *lasiocarpum* (Rchb.) Soó = *A. lasiocarpum* (Rchb.) Gáyer
 subsp. *paniculatum*
 var. *paniculatum* (*nom. inval.*) = *A. degenii* Gáyer
 subsp. *paniculatum* (Arcang.) Mucher var. *laxiflorum* (Rchb.) Mucher
 var. *podolicum* Zapał.
 fo. *latilobum* Zapał. = *A. lasiocarpum* (Rchb.) Gáyer subsp. *kotulae* (Pawł.) Starm. & Mitka
 fo. *latiusculum* Zapał. = *A. degenii* var. *intermedium* (Zapał.) Mitka
 var. *prutense* Zapał. = *A. degenii* var. *intermedium* (Zapał.) Mitka
 var. *turrachense* Mucher = *A. degenii* Gáyer subsp. *paniculatum* (Arcang.) Mucher var. *turrachense* (Mucher) Mucher
A. parviflorum Host. = *A. tauricum* Wulfen in Koelle
A. ×pawlowskii* Mitka & Starm. [A. lasiocarpum* × *A. variegatum***]**
A. ×paxii (Starm.) Mitka = *A. ×mariae* Rottensteiner, Mitka & Novikov nothosubsp. *paxii* (Starm.) Mitka, **comb. nov.** [*A. firmum* subsp. *moravicum* × *A. maninense*]
***A. pentheri* Hayek**
***A. pilipes* (Rchb.) Gáyer**
***A. pilipes* (Rchb.) Gáyer × *A. tauricum* Wulfen**
***A. ×pilosiusculum* (Seringe) Gáyer** [*A. degenii* × *A. pilipes*]
***A. plicatum* Köhler ex Rchb.**
 subsp. ***plicatum***
 var. *clusianum* (Rchb.) Mitka & Starm. = *A. clusianum* Rchb.
 subsp. ***sudeticum*** Mitka
A. podolicum (Zapał.) Voroshylov = *A. lasiocarpum* (Rchb.) Gáyer subsp. *kotulae* (Pawł.) Starm. & Mitka
A. pubiceps Rupr.
 var. *tuscheticum* (N.Busch) Grossh. = *A. ×tuscheticum* (N.Busch) N.Busch [*A. degenii* × *A. vitosanum*]
A. pyramidale Mill. ex Rchb. = *A. napellus* L.
A. ranunculifolium Rchb.
 var. *dolomiticum* Evers, p.p. = *A. tauricum* Wulfen in Koelle nothosubsp. ***hayekianum***

(Gáyer) Grinč. in Savulescu [*A. tauricum* subsp. *latemarensis* × *A. tauricum* subsp. *tauricum*]
var. *dolomiticum* Evers, p.p. = *A. ×teppneri*
Mucher ex Starm. nothosubsp. *kerneri* Starm.
[*A. napellus* subsp. *napellus* × *A. tauricum* subsp.
latemarensis]
A. rigidum Rchb. = *A. plicatum* Köhler ex Rchb.
A. romanicum Wołoszczak = *A. firmum* Rchb. subsp.
fissurae Nyár.
A. ×ronnigeri Gáyer = *A. ×mielichhoferi* Rchb. [*A. tauricum* subsp. *latemarensis* × *A. tauricum* subsp. *tauricum*]
A. ×schneebergense Gáyer [*A. napellus* × *A. variegatum*]
A. schurii Beck. = *A. toxicum* Rchb.
fo. *retyezatense* Gáyer = *A. toxicum* Rchb.
A. skerisorae Auct., non Gáyer = *A. firmum* Rchb.
A. skerisorae Gáyer = *A. firmum* Rchb. subsp. *skerisorae* (Gáyer) Starm.
A. speciosum Otto ex Rchb. = *A. ×exaltatum* Bernh. ex Rchb. [*A. plicatum* × *A. variegatum*]
A. sostaricianum Fritsch = *A. superbum* Fritsch
A. stoerkianum Rchb. = *A. ×cammarum* L. em. Fries [*A. napellus* × *A. variegatum*]
A. strictum Bernh. = *A. napellus* L.
A. superbum Fritsch
A. superbum Fritsch × *A. variegatum* L.
A. superbum Fritsch × *A. vitosanum* Gáyer
A. tatrae Borb. = *A. firmum* Rchb.
var. *skerisorae* Soó = *A. firmum* Rchb. subsp.
skerisorae (Gáyer) Starm.
A. taurericum Rchb. = *A. tauricum* Wulfen in Koelle subsp. *tauricum* var. *tauricum* fo. *taurericum* (Rchb.) Gáyer
A. tauricum Auct. = *A. firmum* Rchb.
A. tauricum Wulfen in Koelle
nothosubsp. *hayekianum* Gáyer (Gáyer) Grinč. in Savulescu [*A. tauricum* subsp. *latemarensis* × *A. tauricum* subsp. *tauricum*]
subsp. *latemarensis* (Degen and Gáyer) Starm.
subsp. *nanum* Auct. fl. Carpat. Orient., non (Baumg.) Grinč. = *A. bucovinense* Zapál.
subsp. *nanum* (Baumg.) Grinč. = *A. ×nanum* (Baumg.) Simonk. [*A. bucovinense* × *A. firmum*]
subsp. *nanum* (Baumg.) Gáyer = *A. ×nanum* (Baumg.) Simonk. [*A. bucovinense* × *A. firmum*]
subsp. *taureticum* (Rchb.) Grinč. = *A. tauricum* Wulfen in Koelle subsp. *tauricum* var. *tauricum* fo. *taurericum* (Rchb.) Gáyer
subsp. *tauricum*
var. *eustachyum* (Rchb.) Starm.
var. *latemarensis* (Degen & Gáyer) Mucher = *A. tauricum* Wulfen in Koelle subsp. subsp. *latemarensis* (Degen & Gáyer) Starm. in W.Maurer

var. **tauricum**
fo. **tauricum**
fo. **taurericum** (Rchb.) Gáyer
A. tenuifolium Host. = *A. angustifolium* Bernh. ex Rchb.
A. ×teppneri Mucher ex Starm. [*A. napellus* × *A. tauricum*]
nothosubsp. *goetzii* Mucher ex Starm. [*A. napellus* subsp. *lobelii* × *A. tauricum* subsp. *tauricum*]
nothosubsp. *haderlappii* Starm. [*A. napellus* subsp. *lobelii* × *A. tauricum* subsp. *latemarensis*]
nothosubsp. *kerneri* Starm. [*A. napellus* subsp. *napellus* × *A. tauricum* subsp. *latemarensis*]
nothosubsp. *teppneri* [*A. napellus* subsp. *napellus* × *A. tauricum* subsp. *tauricum*]
A. toxicum Rchb.
subsp. *bucegiense* (Nyár.) Mucher
subsp. *crispulum* (Nyár.) Mucher
nothosubsp. *nyaradyanum* Mucher [*A. toxicum* subsp. *crispulum* × *A. toxicum* subsp. *toxicum*]
subsp. **toxicum**
var. *bosniacum* (Beck) Hayek = *A. toxicum* Rchb.
var. *bucegiense* Nyár. = *A. toxicum* Rchb. subsp. *bucegiense* (Nyár.) Mucher
var. *crispulum* Nyárády in Borza = *A. toxicum* Rchb. subsp. *crispulum* (Nyár.) Mucher
var. β *schurii* Beck = *A. toxicum* Rchb.
var. *toxicum*
fo. *diabolicum* (Gáyer) Grinč. = *A. toxicum* Rchb.
nothosubsp. *ungarianum* Starm. [*A. toxicum* subsp. *bucegiense* × *A. toxicum* subsp. *toxicum*]
A. trichocarpum Rchb. = *A. tauricum* Wulfen in Koelle
A. ×tuscheticum (N.Busch) N.Busch [*A. degenii* × *A. vitosanum*]
A. valesiacum Gáyer = *A. degenii* Gáyer subsp. *valesiacum* (Gáyer) Mucher
A. variegatum L.
subsp. *kotulae* Pawł. = *A. lasiocarpum* (Rchb.) Gáyer
subsp. *kotulae* (Pawł.) Starm. & Mitka
subsp. *nasutum* (Fischer ex Rchb. em. Rupr.) Götz
nothosubsp. *podobnikianum* Mucher [*A. variegatum* subsp. *nasutum* × *A. variegatum* subsp. *variegatum*]
subsp. *pyrenaicum* Vivant & Delay = *A. vivantii* Rottensteiner
subsp. *valesiacum* (Gáyer) Greuter & Burdet = *A. degenii* Gáyer subsp. *valesiacum* (Gáyer) Mucher
subsp. **variegatum**
var. *carnolicum* Starm.
var. *diffusum* (Rchb.) Penev & Simeon. = *A. vitosanum* Gáyer
var. *stiriacum* Mucher

var. *variegatum*

fo. *aquilonare* (Gáyer) M.Niketić = **A.**
×*aquilonare* A.Kern. ex Gáyer [*A. variegatum*
× *A. vitosanum*]
fo. *kotulae* (Pawł.) Skalický = **A. lasiocarpum**
(Rchb.) Gáyer subsp. **kotulae** (Pawł.) Starm. &
Mitka

A. ×virgatum Rchb. ex Gáyer = **A. ×schneebergense**
Gáyer [*A. napellus* × *A. variegatum*]

A. vitosanum Gáyer

A. vivantii Rottensteiner, **nom. nov.**

A. vulgare DC. = **A. napellus** L.

A. wilsonii Stapf ex Veitch. = **A. carmichaelii** Debeaux

A. ×zahlbruckneri Gáyer = **A. ×acutum** Rchb. [*A. tauricum* × *A. variegatum*]

A. ×zapalowiczii (Starm.) Mitka [*A. firmum* × *A. ×mariae* nothosubsp. *paxii*]