



Citation: Troia A. et al. (2020) Monitoring of alien aquatic plants in the inland waters of Sicily (Italy). *Webbia. Journal of Plant Taxonomy and Geography* 75(1): 77-83. doi: 10.36253/jopt-8414

Received: April 2, 2020

Accepted: May 8, 2020

Published: June 30, 2020

Copyright: © 2020 A. Troia, V. Ilardi, E. Oddo. This is an open access, peer-reviewed article published by Firenze University Press (<http://www.fupress.com/webbia>) and distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Competing Interests: The Author(s) declare(s) no conflict of interest.

Editor: Riccardo M. Baldini, University of Florence, Italy

Monitoring of alien aquatic plants in the inland waters of Sicily (Italy)

ANGELO TROIA^{1,*}, VINCENZO ILARDI², ELISABETTA ODDO¹

¹ Dipartimento STEBICEF (Scienze e Tecnologie Biologiche, Chimiche e Farmaceutiche), Università degli Studi di Palermo, Palermo, Italy

² Dipartimento DISTEM (Scienze della Terra e del Mare), Università degli Studi di Palermo, Palermo, Italy

*Corresponding author, email angelo.troia@unipa.it

Abstract. Updated and reliable data on the presence and distribution of alien aquatic plant species in Sicily are lacking, and there is a need to fill this gap for a proper and efficient management of freshwater ecosystems and biodiversity. This paper reviews the available knowledge about alien aquatic vascular plants in the inland waters of Sicily (Italy). The aim is to provide an updated checklist, as a first step in the study of the impact of those plants on the native species and ecosystems of this Mediterranean island. The paper focuses on the strictly aquatic species (hydrophytes), excluding emergent macrophytes. Four species were listed, all of them free-floating and with American origin. Most of them occur within protected areas, and their introduction in the island appears to be anthropogenic. A set of functional traits of the alien species, such as relative growth rate, leaf mass per area, nitrogen and carbon content, were screened. These traits are useful for assessing the species invasive potential compared to native ones.

Keywords: alien species, aquatic plants, biodiversity, functional traits, hydrophytes, protected areas.

INTRODUCTION

Invasive alien species pose a major global threat to the conservation of biodiversity, causing the extinction of native species and modifying ecosystem functions: this is true also for aquatic habitats, particularly susceptible to invasion due to usually high disturbance regimes affecting these habitats and the easy dispersal of water plant propagules. In addition, aquatic environments are also difficult to monitor, and an early detection of introduction of a submerged species is seldom possible (Brundu 2015).

Mediterranean islands are particularly appreciated model systems for studying invasions due to the diversity of alien taxa, long history of species introductions and (usually) detailed floristic records (Hulme 2004; Lloret et al. 2005; Bjarnason et al. 2017; Chiarucci et al. 2017; Pasta et al. 2017).

The island of Sicily is one of the main hotspots of plant biodiversity, in the center of the Mediterranean basin (Médail and Diadema 2009), housing

more than 3,000 wild plant species (Pignatti et al. 2017-2019) and being an ideal stage for research in ecology and evolution (e.g. Geraci et al. 2009, 2019; Minissale and Sciandrello 2016); it hosts different types of freshwater habitats, both lentic (coastal wetlands, temporary ponds, lakes, reservoirs) and lotic (springs, streams, permanent and seasonal watercourses) (Gianguzzi et al. 2016), with many endemic or endangered species occurring in aquatic habitats (cf. Bonanno and Veneziano 2011; Troia et al. 2012; Troia and Lansdown 2016; Sciandrello et al. 2016; Minissale et al. 2017).

In the last years new aquatic alien species, such as *Lemna minuta* Kunth (Marrone and Naselli-Flores 2011), and new populations of already reported ones, such as *Eichhornia crassipes* (Mart.) Solms (Ferro 2013), have been reported. Nevertheless, no recent and updated synthesis on the ecology and the distribution of the Sicilian alien aquatic flora is currently available.

Thus, the aims of this work were i) to compile data on the distribution of alien vascular plant species with special reference to their occurrence in protected areas, as a first step in the analysis of the effects of the alien species on the aquatic habitats of the island; ii) to compile data on some functional traits (such as relative growth rate, leaf area, leaf mass per area, nitrogen and carbon content) of the alien species, in order to allow predictions about their potential invasiveness of natural wetlands.

MATERIALS AND METHODS

In the literature, the term “water plant” has usually been applied rather arbitrarily in the case of plants occurring in and near the water (cf. Brundu 2015); here we refer to the classification of Chambers et al. (2008), framing the aquatic plants in four main groups: emergent macrophytes (plants that are rooted in submersed soils or soils that are periodically inundated, with foliage extending into the air), floating-leaved macrophytes (plants rooted to the lake or stream bottom with leaves that float on the surface of the water), submersed macrophytes (plants that grow completely submerged under the water, with roots or root analogs in, attached to, or closely associated with the substrate), and free-floating macrophytes (plants that typically float on or under the water surface). In particular, in this study we focused only on the vascular plants belonging to the last 3 types (“hydrophytes” according to Raunkiaer 1934), so excluding the not-strictly aquatic plants. In this sense, our three categories correspond to the three groups adopted by the European and Mediterranean Plant Protection Organisation (EPPO) (EPPO 2014).

Inland waters are here defined according to the Water Framework Directive (Directive 2000/60/EC), excluding transitional waters.

Finally, alien species are here defined as “plant taxa in a given area (...) whose presence there is due to intentional or unintentional human involvement, or which have arrived there without the help of people from an area in which they are alien” (Pyšek et al. 2004).

We consulted all available literature regarding the alien flora in Sicily, to prepare a list of the alien aquatic vascular plants reported for the island, with family names and order according to Smith et al. (2006) and Haston et al. (2009). In addition, databases and photographs from popular websites such as Acta Plantarum (www.actaplantarum.org) have been consulted to obtain recent data and cover under-represented geographic areas. Finally, the management plans for the “Natura2000 sites” (protected areas designated according to the European ‘Habitats’ Directive 92/43/CE, which mostly overlap with natural parks and reserves), available on the website of the Sicilia Region (http://www.artasicilia.eu/old_site/web/natura2000/index.html), were consulted, too.

We reviewed the scientific literature published in the last fifteen years (2006-2020) to screen the most relevant physiological traits that may aid in assessing the invasive potential of alien species compared to native ones. Literature search was made in the Scopus and Google Scholar databases, using as keywords the names of the species resulting from the list compiled as described in the previous paragraph.

RESULTS

Four aquatic alien species (Table 1), belonging to 3 genera and 3 different families, including both ferns (2 species) and angiosperms (2 species), are reported for Sicily.

About *Azolla*, the taxonomy of this genus “is difficult and controversial since the vegetative and some reproductive characters used to identify these species are highly variable, depending on the collection site and on the environmental conditions” (Pereira et al. 2011). In addition, nomenclatural reasons linked to the use of the name “*Azolla caroliniana*” created further confusion, so that the two species previously reported for Sicily (Romano et al. 1994) have been considered a single species in the last national Flora (Pignatti et al. 2017-2019). According to some recent studies (e.g. Evrard and van Hove 2004; Lastrucci et al. 2019), two different species occur in Sicily: *Azolla filiculoides* Lam. and *A. cristata* Kaulf. (= *A. caroliniana* sensu Auct.) (Table 1).

Table 1. List of the alien aquatic vascular plants reported for Sicily.

Species	Family	Origin	Number of known populations in Sicily	Number of populations within a protected area	References
<i>Azolla filiculoides</i> Lam.	Salviniaceae	America (N&S)	3	2	Romano et al. 1994; Albano 2010; Giordana 2013
<i>Azolla cristata</i> Kaulf. (= <i>Azolla mexicana</i>) (= <i>Azolla caroliniana</i> Auct. non Willd.)	Salviniaceae	America (N&S)	2	2	Romano et al. 1994; Minissale and Sciandrello 2017
<i>Lemna minuta</i> Kunth	Araceae	America (N)	1	1	Marrone and Naselli-Flores 2011
<i>Eichhornia crassipes</i> (Mart.) Solms	Pontederiaceae	America (S)	3	2	Bartolo et al. 1976; Ferro 2013; Di Gregorio 2014

Table 2. Functional traits of the 4 alien aquatic species reported for Sicily.

	LA mm ²	LMA mg mm ⁻²	RGR g g ⁻¹ d ⁻¹	NC %	CC %	
<i>Azolla filiculoides</i>	0.9-0.92	0.003-0.024	0.08-0.9	2.01-3.5	35.4-42.5	Brouwer et al. 2018; Cheng et al. 2010; Gufu et al. 2019; Lukács et al. 2017; Pierce et al. 2012; Shaltout et al. 2012; Van Kempen et al. 2016
<i>Azolla cristata</i> (= <i>Azolla mexicana</i>) (= <i>Azolla caroliniana</i> sensu Auct.)			0.09-0.18	4.0		Hassan et al. 2020; Lizieri et al. 2011; Roberts et al. 2014; Rofkar et al. 2014
<i>Lemna minuta</i>	2.4-3.07	0.008-0.01	0.16	2.37-2.7	35.3-41	Pierce et al. 2012; Gérard & Triest 2014; Lukács et al. 2017
<i>Eichhornia crassipes</i>	3500-7300	0.05-0.07	0.002-0.045 ⁽²⁾	2.2-2.5	40	Henry-Silva et al. 2008; Fan et al. 2013; Wang et al. 2017; Eid & Shaltout 2017; Zahoor et al. 2018; Upadhyay & Pame 2019; Wauton & William-Ebi 2019

Abbreviations: LA, leaf area; LMA, leaf mass per area; RGR, relative growth rate; NC, nitrogen content; CC, carbon content. ⁽¹⁾ average of different clones. ⁽²⁾ different nutrient availability or growth sites.

The presence in Sicily of another species, which we did not include here, *Halophila stipulacea* Asch. (Hydrocharitaceae) (Biliotti and Abdelahad 1990), is noteworthy; although it is a seagrass, native to the Indian Ocean and reported for the coasts of Sicily, it seems able to colonize also coastal ponds with salt or brackish water (Procaccini et al. 1999), potentially threatening protected areas such as “Saline di Trapani e Paceco”, “Stagnone di Marsala” and similar areas including saltworks and lagoons.

Table 1 shows that 4 out of 4 species are of American origin, all of them coming from warm, subtropical or tropical areas. This latter circumstance can be explained with the Mediterranean climate occurring in Sicily; the former one is probably to be linked to the reasons reported in the next paragraph. Interestingly, also the 10 alien hydrophytes reported for Sardinia (Mayoral et al. 2018) are American species.

Considering the geographic origin of the species, and their use in many human activities, it seems probable that *Homo sapiens* is the main vector of their introduction: all of these species are in fact commonly used as ornamental plants in gardens, fish tanks and aquaria, or as effective living elements in constructed wetlands (see Mazza et al. 2015).

All the aquatic alien species occurring in Sicily are free-floating species. Other alien aquatic species with a different growth-form potentially occurring in Sicily, for example the rooted *Elodea canadensis* Michx. and *Myriophyllum aquaticum* (Vell.) Verdc. (occurring in northern and central Italy), have not been found (till now at least), probably for climatic reasons.

Practically all the sites in which alien aquatic species were reported are located within protected areas (Table 1), i.e. natural parks and reserves or sites of the Natura2000 network (but also in an archaeological

park). These areas include both coastal habitats (such as wetlands and river mouths) and lakes and ponds in the hills, not far from the coast.

Available literature from the last fifteen years often reported different traits for the alien species found in Sicilian wetlands. The functional traits that could be most widely compared among these species are reported in Table 2. We did not find recent data for *Azolla cristata* in our literature survey: considering the nomenclatural matters cited above, we included in Table 2 data for *A. caroliniana* (*sensu lato*). *Eichhornia crassipes* was the species with the highest leaf area and leaf mass per area (LMA) values. The lowest LMA values were reported for *Lemna minuta*, mainly reflecting the different leaf morphology between the species. Relative growth rate in *Eichhornia crassipes* was lower than the maximum RGR reported for the other species. Tissue nitrogen content was higher for the *Azolla* species, as predictable. Available data on carbon content, instead, were quite similar among the different species.

DISCUSSION

Comparing the number of alien aquatic plant species reported for Sicily with the total number of alien aquatic plant species in the EPPO list (Brundu 2015) - even if it does not include some of the species in our list - reveals that about 1/5 of that number have been found in Sicily, on a very limited area compared to the EPPO region currently including 50 countries (www.eppo.int). However in Sardinia, another Mediterranean island geologically different from Sicily but similar for climate and size, the number of alien hydrophytes is more than double (Mayoral et al. 2018).

The alien aquatic species here reported seem to be limited to few localities, so it seems they are not invasive; but really the feeling is that we have too scarce information - and scattered in time - to say something about their real status.

Our results show that aquatic alien species in Sicily have been reported almost exclusively from protected areas. This probably because protected areas are better studied and monitored, compared to other non-protected areas, and so they can function as 'sentinels' for monitoring the spread of invasive aquatic plant species. But this means also that: 1) the presence of (potentially invasive) alien species in protected areas must be managed (see for example threats and priorities for alien plant invasions in protected areas in Foxcroft et al. 2017); 2) Probably, other populations (and other species?) of alien plants could be already present in "secondary" and overlooked aquatic habitats, outside protected areas.

Usually, monitoring of invasive plant species focuses on terrestrial habitats. During the preparation of the management plans of the Natura2000 sites in Sicily, that were made in the same period all over the island about 12 years ago, special attention to alien invasive species was asked by Regional Coordinators: only in one (Albano 2010) of the 25 examined plans a single aquatic invasive plant species is mentioned.

The situation we found in Sicily for aquatic alien species fits well (unfortunately) with the general lack of information on the presence and abundance of invasive aquatic plants in protected areas in all Mediterranean islands, as highlighted by Brundu (2013), who defined this a serious hindrance for management at international levels.

In the case of alien aquatic species in Sicily, we have seen that their introduction is generally linked to their use and release (intentional or not) by man. The release of alien species in natural habitats is generally prohibited by law, but in Italy, as in several other countries, this is not sufficient to avoid the diffusion of potentially invasive plants. While the impact of alien species on the native aquatic systems of Sicily is well known as regards the faunistic aspects (Marrone and Naselli-Flores 2015), it still has to be evidenced for floristic aspects.

The identification of a general set of functional traits favoring invasiveness may be controversial, as traits of invaders depend on many factors, among which reproductive strategies (Violle et al. 2007), ecological and physiological traits of native species and the environmental conditions of invaded habitats (Funk 2013). Generally, native species are the ones best adapted to a given habitat, but the arrival of alien, more competitive species may alter the ecosystem balance, especially when perturbations of that habitat occur. For terrestrial habitats, many studies have reported differences in morphological or physiological traits between native and invasive taxa, while others have not proved the predictability of invasiveness from these traits (Lefler et al. 2014).

In aquatic environments, an important factor determining invasiveness is nutrient availability. For example, experiments comparing competitiveness between *Lemna minuta* and *L. minor* found that the invasive species was dominant only under high levels of nutrient availability (Njambuya et al. 2011), and therefore invasion would increase with eutrophication, as reported also for *E. crassipes* (Coetzee et al. 2017). In mixed aquatic communities, plant density has been reported as another relevant factor affecting invasiveness, either through facilitation or competition, depending on the species (Silveira and Thiébaud 2020).

LMA is a trait central to the pattern of the leaf economic spectrum, which shows a negative correlation with photosynthetic rate across species (John et al. 2017). High values of LMA are typically found in slow growing species, while low values are common for fast growing species (Wright et al. 2004; Reich 2014). Most hydrophytes share low LMA values, a trait related to high resource acquisition strategies that may aid in competition (Pierce et al. 2012), so invasive species with lower LMA than native ones may be at advantage.

It is clear from this first analysis that updated and complete data on the presence of alien aquatic species in Sicily and on their competitiveness with native species are lacking, so there is the need to fill this gap for a proper and efficient management of ecosystems and biodiversity. Conservation of wetlands remains particularly challenging, given the importance of fresh water for human communities and the consequent pressure wetlands and water bodies are prone to, and this type of ecosystems can no longer be considered the “poor cousins” (Kingsford et al. 2016) of the other terrestrial ones.

ACKNOWLEDGEMENTS

We thank Salvatore Pasta (CNR-IBBR Palermo) for his critical reading of a preliminary version of this paper, and two reviewers for their work and useful suggestions.

REFERENCES

- Albano PG (ed.). 2010. Piano di Gestione del sito “La Gurna e Fiume Fiumefreddo”, Ed. 5. Catania: Provincia di Catania.
- Bartolo G, Brullo S, Marcenò C. 1976. Contributo alla flora sicula. Boll Accad Gioenia Sci Nat Catania, s. 4, 12(9-10): 72-78.
- Biliotti M, Abdelahad N. 1990. *Halophila stipulacea* (Forssk.) Aschers. (Hydrocharitaceae): espèce nouvelle pour l'Italie. Posidonia newsletter 3 (2): 23-26
- Bjarnason A, Katsanevakis S, Galanidis A, Vogiatzakis IN, Moustakas A. 2017. Evaluating hypotheses of plant species invasions on Mediterranean islands: inverse patterns between alien and endemic species. Front Ecol Evol. 5: 91. doi: 10.3389/fevo.2017.00091
- Bonanno G, Veneziano V. 2011. Ecology and distribution of a controversial macrophyte in Sicily: *Zannichellia peltata* (Zannichelliaceae). Biologia 66(5): 833-836.
- Brouwer P, Schlupepmann H, Nierop KG, Elderson J, Bijl PK, van der Meer I, de Visser W, Reichart G-J, Smeekens S, van der Werf A. 2018. Growing *Azolla* to produce sustainable protein feed: the effect of differing species and CO₂ concentrations on biomass productivity and chemical composition. J Sci Food Agric. 98(12): 4759-4768.
- Brundu G. 2013. Invasive Alien Plants in Protected Areas in Mediterranean Islands: Knowledge Gaps and Main Threats. In: Foxcroft, L.C., Pyšek, P., Richardson, D.M., Genovesi, P. (eds.) Plant Invasions in Protected Areas. Patterns, Problems and Challenges. Invading Nature - Springer Series in Invasion Ecology, Vol. 7, pp. 395-422 [ISBN: 978-94-007-7749-1 (Printed) 978-94-007-7750-7 (On-line)].
- Brundu G. 2015. Plant invaders in European and Mediterranean inland waters: profiles, distribution, and threats. Hydrobiologia 746: 61-79.
- Chambers PA, Lacoul P, Murphy KJ, Thomaz SM. 2008. Global diversity of aquatic macrophytes in freshwater. Hydrobiologia 595: 9-26.
- Cheng W, Sakai H, Matsushima M, Yagi K, Hasegawa T. 2010. Response of the floating aquatic fern *Azolla filiculoides* to elevated CO₂, temperature and phosphorous levels. Hydrobiologia 656: 5-14.
- Chiarucci A, Fattorini S, Foggi B, Landi S, Lazzaro L, Podani J, Simberloff D. 2017. Plant recording across two centuries reveals dramatic changes in species diversity of a Mediterranean archipelago. Sci Rep 7: 5415. <https://doi.org/10.1038/s41598-017-05114-5>
- Coetzee JA, Hill MP, Ruiz-Téllez T, Starfinger U, Brunel S. 2017. Monographs on invasive plants in Europe N° 2: *Eichhornia crassipes* (Mart.) Solms. Bot Lett. 164 (4): 303-326.
- Di Gregorio G. 2014. Forum Acta Plantarum, Topic_id: 66946: *Eichhornia crassipes* (Mart.) Solms. <https://floraitaliae.actaplantarum.org/>
- Eid EM, Shaltout KH. 2017. Growth dynamics of water hyacinth (*Eichhornia crassipes*): a modeling approach. Rend Lincei Sci Fis Nat 28(1): 169-181.
- EPPO. 2014. PM 9/19 (1) Invasive alien aquatic plants. EPPO Bulletin 44: 457-471.
- Evrard C, van Hove C. 2004. Taxonomy of the American *Azolla* species (Azollaceae): a critical review. Syst Geogr Pl. 74: 301-318.
- Fan S, Liu C, Yu D, Xie D. 2013. Differences in leaf nitrogen content, photosynthesis, and resource-use efficiency between *Eichhornia crassipes* and a native plant *Monochoria vaginalis* in response to altered sediment nutrient levels. Hydrobiologia 711: 129-137.
- Ferro G. 2013. La Riserva Naturale Fiume Ciane e Saline di Siracusa. In: Cencini C., Corbetta F. (Eds.) Il manuale del bravo conservatore (Saggi di Ecologia applicata). Bologna: Edagricole, pp. 437-438.

- Foxcroft LC, Pyšek P, Richardson DM, Genovesi P, MacFadyen S. 2017. Plant invasion science in protected areas: progress and priorities. *Biol Invasions*. DOI 10.1007/s10530-016-1367-z
- Funk JL. 2013. The physiology of invasive plants in low-resource environments. *Conserv Physiol* 1. DOI: 10.1093/conphys/cot026
- Geraci A, Raimondo FM, Troia A. 2009. Genetic diversity and local population structure in *Ambrosina bassii* (Araceae, Ambrosineae), a Mediterranean relict species. *Biochem Syst Ecol*. 37: 737–746.
- Geraci A, Inzerillo S, Oddo E. 2019. Physio-morphological traits and drought stress responses in three wild Mediterranean taxa of Brassicaceae. *Acta Physiol Plant*. 41: 106. <https://doi.org/10.1007/s11738-019-2899-5>
- Gérard J, Triest L. 2014. The effect of phosphorus reduction and competition on invasive lemnids: life traits and nutrient uptake. *ISRN Bot*. Article ID 514294, <http://dx.doi.org/10.1155/2014/514294>
- Gianguzzi L, Papini F, Cusimano D. 2016. Phytosociological survey vegetation map of Sicily (Mediterranean region). *J Maps*. 12(5): 845-851.
- Giordana F. 2013. Forum Acta Plantarum, Topic_id: 55430: *Azolla filiculoides* Lam. <https://floraitaliae.actaplantarum.org/>
- Gufu GD, Manea A, Leishman MR. 2019. Growth, reproduction and functional trait responses of three freshwater plant species to elevated carbon dioxide. *Aquatic Bot*. 154: 18-23.
- Hassan A, Mohamed HE, Moustafa E. 2020. Nutrient starvation enhances the phenolic compounds and antioxidant activity in *Azolla caroliniana* plant. *Egypt J Bot*. 60 (1): 239-247.
- Haston E, Richardson JE, Stevens PF, Chase MW, Harris DJ. 2009. The Linear Angiosperm Phylogeny Group (LAPG) III: a linear sequence of the families in APG III. *Bot J Linn Soc*. 161: 128-131.
- Henry-Silva GG, Camargo AFM, Pezzato MM. 2008. Growth of free-floating aquatic macrophytes in different concentrations of nutrients. *Hydrobiologia* 610: 153-160.
- Hulme PE. 2004. Invasions, islands and impacts: a Mediterranean perspective. In: *Island Ecology* (ed. J.M. Fernández Palacios), pp. 337–361. Asociación Española de Ecología Terrestre, La Laguna, Spain.
- John GP, Scoffoni C, Buckley TN, Villar R, Poorter H, Sack L. 2017. The anatomical and compositional basis of leaf mass per area. *Ecol Lett*. 20(4): 412-425.
- Kingsford RT, Basset A, Jackson L. 2016. Wetlands: conservation's poor cousins. *Aquat Conserv*. 26(5): 892-916.
- Lastrucci L, Fiorini G, Lunardi L, Viciani D. 2019. Herbarium survey on the genus *Azolla* (Salviniaceae) in Italy: distributive and taxonomic implications. *Pl Biosyst*. 153: 710-719.
- Leffler AJ, James JJ, Monaco TA, Sheley RL. 2014. A new perspective on trait differences between native and invasive exotic plants. *Ecology* 95: 298-305
- Lizieri C, Aguiar R, Kuki KN. 2011. Manganese accumulation and its effects on three tropical aquatic macrophytes: *Azolla caroliniana*, *Salvinia minima* and *Spirodela polyrrhiza*. *Rodriguésia* 62(4): 909-917.
- Lloret F, Médail F, Brundu G, Camarda I, Moragues E, Rita J, Lambdon P, Hulme PE. 2005. Species attributes and invasion success by alien plants in Mediterranean islands. *J Ecol*. 93: 512-520.
- Lukács BA, Vojtkó AE, Mesterházy A, Molnár VA, Süveges K, Végvári Z, Brusa G, Cerabolini BE. 2017. Growth-form and spatiality driving the functional difference of native and alien aquatic plants in Europe. *Ecol Evol*. 7(3): 950-963.
- Marrone F, Naselli-Flores L. 2011. Primo reperto di una lenticchia d'acqua alloctona in Sicilia: *Lemna minuta* Kunth (Araceae Lemnoideae). *Naturalista Sicil*. 35: 179-185.
- Marrone F, Naselli-Flores L. 2015. A review on the animal xenodiversity in Sicilian inland waters (Italy). *Adv Oceanogr Limnol*. 6 (1-2): 2-12.
- Mayoral O, Mascia F, Podda L, Laguna E, Fraga P, Rita J, Frigau L, Bacchetta G. 2018. Alien plant diversity in Mediterranean wetlands: a comparative study within Valencian, Balearic and Sardinian Floras. *Not Bot Horti Agrob*. 46(2): 317-326.
- Mazza G, Aquiloni L, Inghilesi AF, Giuliani C, Lazzaro L, Ferretti G, Lastrucci L, Foggi B, Tricarico E. 2015. Aliens just a click away: the online aquarium trade in Italy. *Manag Biol Invasion*. 6 (3): 253-261.
- Médail F, Diadema K. 2009. Glacial refugia influence plant diversity patterns in the Mediterranean Basin. *J Biogeogr*. 36: 1333–1345.
- Minissale P, Sciandrello S. 2016. Ecological features affect patterns of plant communities in Mediterranean temporary rock pools. *Pl Biosyst*. 150 (1): 171–179.
- Minissale P, Sciandrello S. 2017. The wild vascular flora of the Archaeological Park of Neapolis in Syracuse and surrounding areas (Sicily, Italy). *Biodivers J*. 8(1): 87–104.
- Minissale P, Molina JA & Sciandrello S. 2017. *Pilularia minuta* Durieu (Marsileaceae) discovered in south-eastern-Sicily: new insights on its ecology, distribution and conservation status. *Bot Lett*. 164: 197-208.
- Njambuya J, Stiers I, Triest L. 2011. Competition between *Lemna minuta* and *Lemna minor* at different nutrient concentrations. *Aquatic Bot*. 94: 158-164

- Pasta S, Ardenghi NM, Badalamenti E, La Mantia T, Livreri Console S, Parolo G. 2017. The alien vascular flora of Linosa (Pelagie Islands, Strait of Sicily): update and management proposals. *Willdenowia*. 47(2): 135-144.
- Pereira AL, Martins M, Margarida Oliveira M, Carrapico F. 2011. Morphological and genetic diversity of the family Azollaceae inferred from vegetative characters and RAPD markers. *Pl Syst Evol*. 297: 213-226.
- Pierce S, Brusa G, Sartori M, Cerabolini BEL. 2012. Combined use of leaf size and economic traits allows direct comparison of hydrophyte and terrestrial herbaceous adaptive strategies. *Ann Bot (Oxford)*. 109: 1047-1053
- Pignatti S, La Rosa M, Guarino R. 2017-2019. *Flora d'Italia*. 2nd edition. 4 Voll. Milano-Bologna: Edagricole – New Business Media.
- Procaccini G, Acunto S, Famà P, Maltagliati F. 1999. Structural, morphological and genetic variability in *Halophila stipulacea* (Hydrocharitaceae) populations in the Western Mediterranean. *Mar Biol*. 135: 181-189.
- Pyšek P, Richardson DM, Rejmánek M, Webster GL, Williamson M, Kirschner J. 2004. Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon*. 53: 131-143.
- Raunkiaer CC. 1934. *The Life Forms of Plants and Statistical Plant Geography*. Oxford: Oxford University Press.
- Reich PB. 2014. The world-wide 'fast-slow' plant economics spectrum: a traits manifesto. *J Ecol*. 102(2): 275-301.
- Roberts AE, Boylen CW, Nierzwicki-Bauer SA. 2014. Effects of lead accumulation on the *Azolla caroliniana*-*Anabaena* association. *Ecotox Environ Safe*. 102: 100-104.
- Rofkar JR, Dwyer DE, Bobak DM. 2014. Uptake and toxicity of arsenic, copper, and silicon in *Azolla caroliniana* and *Lemna minor*. *Int J Phytoremediat*. 16(2): 155-166.
- Romano S, Ottonello D, Marcenò C. 1994. Contributo alla floristica siciliana: nuovi rinvenimenti ed ulteriori dati distributivi di alcune entità indigene ed esotiche. *Naturalista Sicil*. 18(1-2): 3-14.
- Sciandrello S, Puglisi M, Privitera M, Minissale P. 2016. Diversity and spatial patterns of plant communities in volcanic temporary ponds of Sicily (Italy). *Biologia* 71(7): 793-803.
- Shaltout KH, El-Komi TM, Eid EM. 2012. Seasonal variation in the phytomass, chemical composition and nutritional value of *Azolla filiculoides* Lam. along the water courses in the Nile Delta, Egypt. *Feddes Reperert*. 123: 37-49.
- Silveira MJ, Thiébaud G. 2020. Effect of density and neighbours on interactions between invasive plants of similar growth form. *Aquatic Ecol*. 54: 463-474.
- Smith AR, Pryer KM, Schuettpelz E, Korall P, Schneider H, Wolf PG. 2006. A classification for extant ferns. *Taxon*. 55 (3): 705-731.
- Troia A, Lansdown R. 2016. The first confirmed population of the globally endangered *Pilularia minuta* (Marsileaceae) in Sicily. *Webbia*. 71(2): 283-286.
- Troia A, Bazan G, Schicchi R. 2012. Micromorphological approach to the systematics of Mediterranean *Isoetes* species (Isoëtaceae, Lycopodiophyta): analysis of the megaspore surface. *Grana*. 51: 35-43.
- Upadhyay RK, Pame P. 2019. Lead phyto-toxicity induced by accumulation and uptake potentially inhibits morpho-physiological depression and alterations in an aquatic model plant, *Eichhornia crassipes*. *Eurasia J BioSci*. 13(2): 1565-1573.
- Van Kempen MML, Smolders AJP, Bogemann GM, Lamers LPM, Roelofs JGM. 2016. Interacting effects of atmospheric CO₂ enrichment and solar radiation on growth of the aquatic fern *Azolla filiculoides*. *Freshwater Biol*. 61: 596-606.
- Violle C, Navas M-L, Vile D, Kazakou E, Fortunel C et al. 2007. Let the concept of trait be functional! *Oikos* 116: 882-892
- Wang T, Hu J, Liu C, Yu D. 2017. Soil type can determine invasion success of *Eichhornia crassipes*. *Hydrobiologia*. 788(1): 281-291.
- Wauton I, William-Ebi D. 2019. Characterization of water hyacinth (*Eichhornia crassipes*) for the production of thermochemical fuels. *J Multidiscip Eng Sci Stud*. 5(7): 2661-65.
- Wright IJ, Reich PB, Westoby M, Ackerly DD, Baruch Z, Bongers F et al. 2004. The worldwide leaf economics spectrum. *Nature*. 428(6985): 821-827.
- Zahoor A, Ahmad F, Hameed M, Basra SMA. 2018. Structural and functional aspects of photosynthetic response in *Eichhornia crassipes* (Mart.) Solms under cadmium stress. *Pakistan J Bot*. 50(2): 489-493.